

The total irregularity of a graph

Hosam Abdo, Stephan Brandt, D. Dimitrov

► **To cite this version:**

Hosam Abdo, Stephan Brandt, D. Dimitrov. The total irregularity of a graph. *Discrete Mathematics and Theoretical Computer Science, DMTCS*, 2014, Vol. 16 no. 1 (in progress) (1), pp.201–206. hal-01179217

HAL Id: hal-01179217

<https://hal.inria.fr/hal-01179217>

Submitted on 22 Jul 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The total irregularity of a graph

Hosam Abdo^{1*}Stephan Brandt^{2†}Darko Dimitrov^{1‡}¹Institut für Informatik, Freie Universität Berlin, Germany²Department of Mathematics and Computer Science, University of Southern Denmark, Odense, Denmarkreceived 1st Oct. 2012, revised 10th Feb. 2014, accepted 18th Mar. 2014.

In this note a new measure of irregularity of a graph G is introduced. It is named the *total irregularity* of a graph and is defined as $\text{irr}_t(G) = \frac{1}{2} \sum_{u,v \in V(G)} |d_G(u) - d_G(v)|$, where $d_G(u)$ denotes the degree of a vertex $u \in V(G)$. All graphs with maximal total irregularity are determined. It is also shown that among all trees of the same order the star has the maximal total irregularity.

Keywords: the irregularity of a graph, the total irregularity of a graph, extremal graphs

1 Introduction

We consider only finite undirected graphs without loops or multiple edges. For terminology and notation not defined here we refer the reader to Bondy and Murty's textbook [6].

In this note we investigate irregularity measures of graphs. A graph is *regular* if all its vertices have the same degree, otherwise it is *irregular*. However, it is of interest to measure how irregular it is. Several approaches have been proposed that characterize how irregular a graph is.

For a graph G , Albertson [4] defines the *imbalance* of an edge $e = uv \in E(G)$ as $|d_G(u) - d_G(v)|$ and the *irregularity* of G as

$$\text{irr}(G) = \sum_{uv \in E(G)} |d_G(u) - d_G(v)|, \quad (1)$$

where $d_G(v)$ is the degree of $v \in V(G)$. He presented upper bounds on the irregularity of graphs, bipartite graphs, and triangle-free graphs, as well as a sharp upper bound for trees. He showed that if a graph has maximal irregularity among all graphs of order n , then it is the join of a clique and an independent vertex set. Some claims about bipartite graphs given in [4] have been formally proved in [16]. Related to the work of Albertson is the work of Hansen and Mélot [15], who characterized the graphs with n vertices

*Email: abdo@mi.fu-berlin.de

†Email: stephan.brandt@imada.sdu.dk

‡Email: darko@mi.fu-berlin.de

and m edges with maximal irregularity. For more results on imbalance, the irregularity of a graph, and other approaches, that measure how irregular a graph is, we refer the reader to [2, 3, 5, 7–12, 17–20].

In the sequel we introduce and consider an irregularity measure that is related to the irregularity measure (1). As well as (1), the new measure also captures the irregularity only by the difference of vertex degrees. For a graph G it is defined as

$$\text{irr}_t(G) = \frac{1}{2} \sum_{(u,v) \in V^2(G)} |d_G(u) - d_G(v)|. \quad (2)$$

For an ordered degree sequence of $V(G) = \{v_1, v_2, \dots, v_n\}$ with $d(v_1) \leq d(v_2) \leq \dots \leq d(v_n)$, (2) can be expressed in the form

$$\text{irr}_t(G) = \sum_{i>j} (d(v_i) - d(v_j)). \quad (3)$$

Because of the obvious connection with the irregularity, we called $\text{irr}_t(G)$ the *total irregularity* of a graph. Note that for every graph the total irregularity is an even number since the number of vertices of odd degree is even, and it is completely determined by its degree sequence – graphs with the same degree sequence have the same total irregularity, while this is not always true with the irregularity of a graph (see Figure 1 for such an example). Since the total irregularity of a graph depends only on its




Fig. 1: Two non-isomorphic graphs G_1 and G_2 with the same degree sequence 1, 1, 1, 1, 2, 3, 3. They have different irregularities ($\text{irr}(G_1) = 10$ and $\text{irr}(G_2) = 8$), but the same total irregularity ($\text{irr}_t(G_1) = \text{irr}_t(G_2) = 22$).

degree sequence, it can be applied as irregularity measure when the adjacency information of the vertices is unknown. Another motivation to introduce irr_t as an irregularity measure is the fact that the graphs with maximal irr , among graphs of same order, are bidegreed graphs, more precisely the so-called *clique-star graphs*, graphs obtained by join of a clique and isolated vertices [1, 4]. Also there are a graphs with large irr that have very small degree sets, the property that one will not expect from very irregular graphs. On the contrary, as we show in this note, the graphs with maximal irr_t have large degree sets, and some of them have even the largest possible ones.

Both measures are zero if and only if G is regular, and $\text{irr}_t(G)$ is an upper bound of $\text{irr}(G)$. Very recently, these two irregularity measures were compared in [13], where it was shown that for a connected graph G with n vertices, $\text{irr}_t(G) \leq n^2 \text{irr}(G)/4$. Moreover, if G is a tree, then it was shown that $\text{irr}_t(G) \leq (n-2)\text{irr}(G)$.

Since the most irregular graphs with respect to irr are the join of a complete graph and an independent set, they have only two different vertex degrees. We will show below that the most irregular graphs with respect to irr_t are graphs that have many different vertex degrees, and the graphs with the maximal number of different vertex degrees belong to the graphs with maximal total irregularity.

2 Graphs with maximal total irregularity

It is easy to see that there is no graph of order n with n different vertex degrees. Asking for graphs with $n - 1$ different degrees there are two possible sets of degrees: $\{1, 2, \dots, n - 1\}$ and $\{0, 1, \dots, n - 2\}$ since the graph cannot contain vertices of degree 0 and $n - 1$ at the same time. A simple induction proof verifies that for every n there is a unique graph with the respective set of degrees up to isomorphism, see Figure 2. These graphs have been called "half-complete" in [14]. The graph with a vertex of degree $n - 1$ is denoted H_n and the graph with a vertex of degree 0 is its complement \overline{H}_n (this can be seen by mirroring in Figure 2 one of the graphs H_n, \overline{H}_n along a horizontal line). The graphs H_n and \overline{H}_n can be obtained by labelling the vertices v_1, \dots, v_n and adding all edges $v_i v_j$ where $i + j \geq n + 1$ ($i + j > n + 1$, respectively). Note that the two vertices of H_n with equal degrees have indices $\lfloor \frac{n}{2} \rfloor$ and $\lfloor \frac{n}{2} \rfloor + 1$.




Fig. 2: The graphs H_n (with dashed edges) and \overline{H}_n (without dashed edges) for even and odd n , respectively.

Take the graph H_n with vertices labelled as above and let $S = \{v_i v_j \mid i + j = n + 1\}$, the set of dashed edges in Figure 2. Then $\overline{H}_n = H_n - S$. Now we can formulate our main result.

Theorem 2.1 A graph G of order n has maximal total irregularity, if and only if it is isomorphic to a graph $H_n - S'$, where $\emptyset \subseteq S' \subseteq S$.

Observe that the set S' of $G = H_n - S'$ can be determined only from the degree sequence of G . This follows from the fact that the vertex labelling of G inherited from H_n has the property that $d_G(v_1) \leq d_G(v_2) \leq \dots \leq d_G(v_n)$. Therefore for $j < i, i + j = n + 1$, we have $v_i v_j \in S'$ if and only if $d_G(v_j) = j - 1$. In particular, every two graphs $G_1 = H_n - S_1$ and $G_2 = H_n - S_2$ with $\emptyset \subseteq S_1 \neq S_2 \subseteq S$ are not isomorphic. Therefore, since $|S| = \lfloor \frac{n}{2} \rfloor$ there are exactly $2^{\lfloor \frac{n}{2} \rfloor}$ non-isomorphic graphs of order n with maximal total irregularity.

Note that all these graphs have the property that they are subgraphs of H_n and supergraphs of \overline{H}_n , but the converse does not hold. E.g., the path on four vertices is a subgraph of H_4 and a supergraph of \overline{H}_4 , but its total irregularity is 4, while the maximal total irregularity is 6.

Proof: For an ordered degree sequence, consider an edge $v_i v_j, i > j$. Adding $v_i v_j$ to G or deleting $v_i v_j$ from G only changes the summands $d(u) - d(w)$ in (3) where $|\{u, w\} \cap \{v_i, v_j\}| = 1$. Adding the edge $v_i v_j$ to G increases all summands $d(v_i) - d(v_k)$ with $i > k$ and $i \neq j$ by one while the summands $d(v_k) - d(v_i)$ for $k > i$ are decreased by one (or increased by one, if $d_G(v_k) = d_G(v_i)$). Similarly, all summands $d(v_j) - d(v_k)$ with $j > k$ are increased by one, while the summands $d(v_k) - d(v_i)$ for $k > i$

and $j \neq i$ are decreased by one (or increased by one, if $d_G(v_k) = d_G(v_j)$). Therefore,

$$\begin{aligned} \text{irr}_t(G + v_i v_j) &\geq \text{irr}_t(G) + (i - 2) - (n - i) + (j - 1) - (n - j - 1) \\ &= \text{irr}_t(G) + 2(i + j - 1) - 2n \end{aligned}$$

which exceeds $\text{irr}_t(G)$ if $i + j > n + 1$.

Deleting an edge $v_i v_j$ from G has the effect that all summands $d(v_k) - d(v_i)$ with $k > i$ are increased by one while the summands $d(v_i) - d(v_k)$ with $k < i$, $k \neq j$ are decreased by one (or increased by one, if $d_G(v_k) = d_G(v_i)$). All summands $d(v_k) - d(v_j)$ with $k > j$, $k \neq i$, are increased by one, while the summands $d(v_j) - d(v_k)$ with $k < j$, are decreased by one (or increased by one, if $d_G(v_k) = d_G(v_j)$). Therefore,

$$\begin{aligned} \text{irr}_t(G - v_i v_j) &\geq \text{irr}_t(G) + (n - i) - (i - 2) + (n - j - 1) - (j - 1) \\ &= \text{irr}_t(G) + 2n - 2(i + j - 1), \end{aligned} \quad (4)$$

with equality if and only if $d_G(v_j) > d_G(v_{j-1})$ and $d_G(v_i) > d_G(v_{i-1})$ ($d_G(v_i) > d_G(v_{i-2})$ if $j = i - 1$). Therefore, if G has maximal irregularity then G has all edges $v_i v_j$ with $i + j > n + 1$ and no edge with $i + j < n + 1$. Any such graph can be expressed in the form $G = H_n - S'$ with $\emptyset \subseteq S' \subseteq S$. It is left to show that every such graph has the same total irregularity. Let $G = H_n - S'$ where $\emptyset \subseteq S' \subseteq S$ and let $v_i v_j \in S \setminus S'$. Then v_i and v_j have both their original degree from H_n , but $d(v_{i-1})$ and $d(v_{j-1})$ can only have decreased. Since $d_{H_n}(v_j) > d_{H_n}(v_{j-1})$ and $d_{H_n}(v_i) > d_{H_n}(v_{i-1})$ if $j < i - 1$ and $d_{H_n}(v_i) = d_{H_n}(v_j) > d_{H_n}(v_{j-1})$ if $j = i - 1$, the inequality (4) holds with equality and $\text{irr}_t(G - v_i v_j) = \text{irr}_t(G)$. \square

Corollary 2.2 *For any simple undirected graph G with n vertices it holds that*

$$\text{irr}_t(G) \leq \begin{cases} \frac{1}{12}(2n^3 - 3n^2 - 2n + 3) & n \text{ odd,} \\ \frac{1}{12}(2n^3 - 3n^2 - 2n) & n \text{ even.} \end{cases}$$

Proof: The bounds follow with Theorem 2.1 from the total irregularity of H_n . The only two vertices of H_n with equal degrees are $v_{\lfloor \frac{n}{2} \rfloor}$ and $v_{\lfloor \frac{n}{2} \rfloor + 1}$. Since, $1 = d(v_1) < \dots < d(v_{\lfloor \frac{n}{2} \rfloor}) = d(v_{\lfloor \frac{n}{2} \rfloor + 1}) < \dots < d(v_n) = n - 1$, we obtain

$$\begin{aligned} \text{irr}_t(H_n) &= \sum_{i>j} (d(v_i) - d(v_j)) = \sum_{i=2}^{n-1} \sum_{j=1}^{i-1} (d(v_i) - d(v_j)) \\ &= \sum_{i=2}^{n-1} \sum_{j=1}^{i-1} (i - j) + \sum_{j=1}^{\lfloor \frac{n}{2} \rfloor - 1} \left(\lfloor \frac{n}{2} \rfloor - j \right) + \sum_{i=\lfloor \frac{n}{2} \rfloor + 1}^{n-1} \left(i - \lfloor \frac{n}{2} \rfloor \right) \\ &= \sum_{i=2}^{n-1} \sum_{j=1}^{i-1} j + \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor - 1} i + \sum_{i=1}^{\lceil \frac{n}{2} \rceil - 1} i = \binom{n}{3} + \binom{\lfloor \frac{n}{2} \rfloor}{2} + \binom{\lceil \frac{n}{2} \rceil}{2}, \end{aligned}$$

which immediately gives $\text{irr}_t(H_n) = \frac{1}{12}(2n^3 - 3n^2 - 2n + 3)$ for odd n , and $\text{irr}_t(H_n) = \frac{1}{12}(2n^3 - 3n^2 - 2n)$ for even n . \square

Proposition 2.3 *If T is a tree with n vertices, then $\text{irr}_t(T) \leq (n-1)(n-2)$. Moreover, equality holds if and only if T is a star.*

Proof:

Let T be a tree that is not a star, with u being a vertex with maximal degree. Consider a pendant vertex v that is adjacent to a vertex $w \neq u$. We remove the edge vw from T and add the edge uv . After this replacement, only the degrees of u and w are affected, namely, $d(u)$ increases by one while $d(w)$ decreases by one. Thus, the summands $|d(u) - d(w)|$ and $|d(w) - d(u)|$ in (2) both increase by two. Each of the $2(n-2)$ summands $|d(u) - d(x)|$ and $|d(x) - d(u)|$ with $x \neq u, w$ increases by one while each of $2(n-2)$ summands $|d(w) - d(x)|$ and $|d(x) - d(w)|$ with $x \neq u, w$ increases or decreases by one. Thus, we have $\text{irr}_t(T - vw + uv) \geq \text{irr}_t(T) + 2 + n - 2 - (n - 2) = \text{irr}_t(T) + 2$. Therefore the largest total irregularity among the trees of order n is only attained by the star, whose total irregularity equals $(n-1)(n-2)$. \square

References

- [1] H. Abdo, N. Cohen, and D. Dimitrov. Graphs with maximal irregularity. *Filomat*, 2014, in press.
- [2] Y. Alavi, A. Boals, G. Chartrand, P. Erdős, and O. R. Oellermann. k -path irregular graphs. *Congr. Numer.*, 65:201–210, 1988.
- [3] Y. Alavi, G. Chartrand, F. R. K. Chung, P. Erdős, R. L. Graham, and O. R. Oellermann. Highly irregular graphs. *J. Graph Theory*, 11:235–249, 1987.
- [4] M. O. Albertson. The irregularity of a graph. *Ars Comb.*, 46:219–225, 1997.
- [5] F. K. Bell. On the maximal index of connected graphs. *Linear Algebra Appl.*, 144:135–151, 1991.
- [6] A. Bondy and U. S. R. Murty. *Graph Theory*. Springer Verlag, Berlin, 2008.
- [7] Y. Caro and R. Yuster. Graphs with large variance. *Ars Comb.*, 57:151–162, 2000.
- [8] G. Chartrand, P. Erdős, and O. R. Oellermann. How to define an irregular graph. *Coll. Math. J.*, 19:36–42, 1988.
- [9] G. Chartrand, K. S. Holbert, O. R. Oellermann, and H. C. Swart. F -degrees in graphs. *Ars Comb.*, 24:133–148, 1987.
- [10] G. Chartrand, M. S. Jacobson, J. Lehel, O. Oellerman, S. Ruiz, and F. Saba. Irregular networks. *Congr. Numer.*, 64:197–210, 1988.
- [11] L. Collatz and U. Sinogowitz. Spektren endlicher Graphen. *Abh. Math. Sem. Univ. Hamburg*, 21:63–77, 1957.

- [12] D. Cvetković and P. Rowlinson. On connected graphs with maximal index. *Publications de l'Institut Mathématique (Beograd)*, 44:29–34, 1988.
- [13] D. Dimitrov and R. Škrekovski. Comparing the irregularity and the total irregularity of graphs. *Ars Math. Contemp.*, 2014, in press.
- [14] P. C. Fishburn. Packing graphs with odd and even trees. *J. Graph Theory*, 7:369–383, 1983.
- [15] P. Hansen and H. Mélot. Variable neighborhood search for extremal graphs. 9. bounding the irregularity of a graph. *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.*, 69:253–264, 2005.
- [16] M. A. Henning and D. Rautenbach. On the irregularity of bipartite graphs. *Discrete Math.*, 307:1467–1472, 2007.
- [17] D. E. Jackson and R. Entringer. Totally segregated graphs. *Congress. Numer.*, 55:159–165, 1986.
- [18] D. Rautenbach. Propagation of mean degrees. *Electr. J. Comb.*, 11:N11, 2004. N11.
- [19] D. Rautenbach and I. Schiermeyer. Extremal problems for imbalanced edges. *Graphs Comb.*, 22:103–111, 2006.
- [20] D. Rautenbach and L. Volkmann. How local irregularity gets global in a graph. *J. Graph Theory*, 41:18–23, 2002.