

A characterization of trees with equal 2-dominating and 2-independence numbers

Christoph Brause^{1,2} Michael A. Henning^{2*} Marcin Krzywkowski^{2,3†}

¹ Institute of Discrete Mathematics and Algebra, TU Bergakademie Freiberg, Germany

² Department of Pure and Applied Mathematics, University of Johannesburg, South Africa

³ Faculty of Electronics, Telecommunications and Informatics, Gdansk University of Technology, Poland

received 25th Apr. 2016, accepted 12th Jan. 2017.

A set S of vertices in a graph G is a 2-dominating set if every vertex of G not in S is adjacent to at least two vertices in S , and S is a 2-independent set if every vertex in S is adjacent to at most one vertex of S . The 2-dominating number $\gamma_2(G)$ is the minimum cardinality of a 2-dominating set in G , and the 2-independence number $\alpha_2(G)$ is the maximum cardinality of a 2-independent set in G . Chellali and Meddah [*Trees with equal 2-dominating and 2-independence numbers*, *Discussiones Mathematicae Graph Theory* 32 (2012), 263–270] provided a constructive characterization of trees with equal 2-dominating and 2-independence numbers. Their characterization is in terms of global properties of a tree, and involves properties of minimum 2-dominating and maximum 2-independent sets in the tree at each stage of the construction. We provide a constructive characterization that relies only on local properties of the tree at each stage of the construction.

Keywords: 2-dominating, 2-dominating number, 2-independence, 2-independence number, tree

1 Introduction

We continue the study of 2-dominating and 2-independence in trees. For $k \geq 1$, a k -dominating set of a graph G is a set S of vertices of G such that every vertex outside S has at least k neighbors in S , while S is a k -independent set if every vertex in S is adjacent to at most $k - 1$ vertices of S . The k -dominating number of G , denoted by $\gamma_k(G)$, is the minimum cardinality of a k -dominating set of G , and the k -independence number of G , denoted by $\alpha_k(G)$, is the maximum cardinality of a k -independent set of G . In particular, we note that for $k = 1$, a 1-dominating set and a 1-independent set are the classical dominating and independent sets, respectively. Thus, the 1-dominating number of G , $\gamma_1(G)$, is the dominating number $\gamma(G)$ and the 1-independence number of G , $\alpha_1(G)$, is the independence number $\alpha(G)$. A k -dominating set of G of minimum cardinality is called a $\gamma_k(G)$ -set, and a k -independent set of G of maximum cardinality is called an $\alpha_k(G)$ -set.

The concepts of k -domination and k -independence in graphs were introduced by Fink and Jacobson [9] in 1985 and is now very well studied in the literature (see for example [1–8, 10, 11]). We refer the reader to

*Research supported in part by the South African National Research Foundation and the University of Johannesburg

†Research fellow of the Claude Leon Foundation at the University of Johannesburg. Research partially supported by the Polish National Science Centre grant 2011/02/A/ST6/00201.

the two books on domination by Haynes, Hedetniemi, and Slater [12, 13], as well as to the excellent survey on k -domination and k -independence in graphs by Chellali, Favaron, Hansberg, and Volkmann [2].

Fink and Jacobson [9] proved that $\gamma_2(G) \leq \alpha_2(G)$ for every graph G . Recently, Chellali and Meddah [3] gave a constructive characterization of trees T satisfying $\gamma_2(T) = \alpha_2(T)$. Their characterization is in terms of global properties of a tree, and involves properties of minimum 2-dominating and maximum 2-independent sets in the tree at each stage of the construction. We provide a constructive characterization that relies only on local properties of the tree at each stage of the construction.

1.1 Notation

For notation and graph theory terminology not defined herein, we refer the reader to [14]. Let G be a graph with vertex set $V(G)$ of order $n(G) = |V(G)|$ and edge set $E(G)$ of size $m(G) = |E(G)|$. A path on n vertices is denoted by P_n . For two vertices u and v in a connected graph G , the distance $d_G(u, v)$ between u and v is the length of a shortest (u, v) -path in G . The maximum distance among all pairs of vertices of G is called the *diameter* of G , which is denoted by $\text{diam}(G)$. A path of length $\text{diam}(G)$ between two vertices at maximum distance apart in G is a *diametrical path* of G . A vertex of degree one is called a *leaf* and its neighbor a *support vertex*. We denote the set of leaves of a tree T by $L(T)$. A *star* is a tree $K_{1,k}$ for some $k \geq 1$, while for $r, s \geq 1$, a *double star* $S_{r,s}$ is a tree with exactly two vertices that are not leaves, one of which is adjacent to r leaves and the other to s leaves.

The *open neighborhood* of a vertex $v \in V(G)$ is the set $N_G(v) = \{u \in V(G) : uv \in E(G)\}$, and the *closed neighborhood* of v is $N_G[v] = N_G(v) \cup \{v\}$. The degree of v is $d_G(v) = |N_G(v)|$. The *open neighborhood* of a set of vertices $S \subseteq V(G)$ is the set $N_G(S) = \bigcup_{v \in S} N_G(v)$, and the *closed neighborhood* of S is $N_G[S] = N_G(S) \cup S$.

For a set $S \subseteq V(G)$, we let $G[S]$ denote the subgraph induced by S . The graph obtained from G by removing the vertices of S along with all edges incident to vertices in S is denoted by $G - S$. If $S = \{v\}$, then we simply denote $G - S$ by $G - v$. We define the *boundary* of S , denoted by $\partial(S)$, to be the set of vertices of S that have a neighbor in $V(G) \setminus S$.

A *rooted tree* T distinguishes one vertex r called the *root*. For each vertex $v \neq r$ of T , the *parent* of v is the neighbor of v on the unique (r, v) -path, while a *child* of v is any other neighbor of v . The set of all children of v we denote by $C(v)$. A *descendant* of v is a vertex $u \neq v$ such that the unique (r, u) -path contains v . Thus every child of v is a descendant of v . We let $D(v)$ denote the set of all descendants of v , and we define $D[v] = D(v) \cup \{v\}$. The *maximal subtree* at v is the subtree of T induced by $D[v]$, and it is denoted by T_v^r . If the root r is clear from the context, then we simply denote the maximal subtree at v by T_v .

1.2 Known Results

Fink and Jacobson [9] proved that $\gamma_2(G) \leq \alpha_2(G)$ for every graph G , and conjectured that for every graph G and integer $k \geq 1$ we have $\gamma_k(G) \leq \alpha_k(G)$. Their conjecture was proven by Favaron [6] by the following stronger result.

Theorem 1 ([6]). *For every graph G and integer $k \geq 1$, the graph G contains a set that is both k -dominating and k -independent, and therefore $\gamma_k(G) \leq \alpha_k(G)$.*

A graph G that satisfies $\gamma_k(G) = \alpha_k(G)$ we call a (γ_k, α_k) -graph. Recently, Chellali and Meddah [3] gave a constructive characterization of (γ_2, α_2) -trees. For this purpose, they defined a family \mathcal{O} of trees $T = T_i$ that can be obtained as follows. Let \mathcal{O} be the family of trees that T that can be obtained from a sequence T_1, T_2, \dots, T_k ($k \geq 1$) of trees, where T_1 is a star $K_{1,p}$ ($p \geq 1$), $T = T_k$, and, if $k \geq 2$, then T_{i+1} is obtained recursively from T_i by one of the following operations:

- **Operation \mathcal{R}_1 :** Add a star $K_{1,p}$, $p \geq 2$, centered at a vertex u and join u by an edge to a vertex of T_i .
- **Operation \mathcal{R}_2 :** Add a double star $S_{1,p}$ with support vertices u and v , where $|L_v| = p$ and join v by an edge to a vertex w of T_i with the condition that if $\gamma_2(T_i - w) = \gamma_2(T_i) - 1$, then no neighbor of w in T_i belongs to a $\gamma_2(T_i - w)$ -set.
- **Operation \mathcal{R}_3 :** Add a path $P_2 = u'v$ and join u' by an edge to a leaf v of T_i that belongs to every $\alpha_2(T_i)$ -set and satisfies in addition $\alpha_2(T_i - v) + 1 = \alpha_2(T_i)$.
- **Operation \mathcal{R}_4 :** Add a path $P_3 = u'vw$ and join v by an edge to a vertex w that belongs to a $\gamma_2(T_i)$ -set and satisfies further $\gamma_2(T_i - w) \leq \gamma_2(T_i)$, with the condition that if $\gamma_2(T_i - w) = \gamma_2(T_i) - 1$, then no neighbor of w in T_i belongs to a $\gamma_2(T_i - w)$ -set.

We are now in a position to state the result due to Chellali and Meddah [3].

Theorem 2 ([3]). *A tree T is a (γ_2, α_2) -tree if and only if $T \cong K_1$ or $T \in \mathcal{O}$.*

2 Main Result

The Chellali and Meddah [3] characterization of (γ_2, α_2) -trees presented in Theorem 2 is a pleasing and important result. However, the characterization is not fully satisfactory in the sense that it is dependant on global properties of the tree at each stage of the construction. For example, in Operation \mathcal{R}_2 one needs to check that the tree T_i and the vertex w satisfy the condition that if $\gamma_2(T_i - w) = \gamma_2(T_i) - 1$, then no neighbor of w in T_i belongs to a $\gamma_2(T_i - w)$ -set. Operations \mathcal{R}_3 and \mathcal{R}_4 also require to check global properties involving minimum 2-dominating and maximum 2-independent sets in the tree. Motivated by the Chellali-Meddah construction of (γ_2, α_2) -trees, our aim is to obtain a constructive characterization that relies only on local properties of the tree at each stage of the construction. We describe such a family \mathcal{T} of (γ_2, α_2) -trees in Section 3. Our main result is the following constructive characterization of (γ_2, α_2) -trees. A proof of Theorem 3 is presented in Section 4.

Theorem 3. *A tree is T a (γ_2, α_2) -tree if and only if $T \in \mathcal{T}$.*

3 The Family \mathcal{T}

In this section, we define a family \mathcal{T} of (γ_2, α_2) -trees. For this purpose, we first define two sets of trees $A = \{T_1, \dots, T_{15}\}$ and $B = \{B_1, \dots, B_{10}\}$ shown in Figure 1 and Figure 2, respectively. We call each tree that belongs to $A \cup B$ a *special tree*.

For each special tree, we 2-color the vertices with the colors white and black as illustrated in Figures 1 and 2 to indicate the roles they play in the tree. We note that exactly one vertex in each special tree is white. Given a special tree T , we denote the set of black vertices by $V_B(T)$. We also specify certain vertices of each special tree T , which we name $v(T)$, $v_1(T)$, $v_2(T)$ and $w(T)$. If a special tree T is clear from context, then we simply refer to these specified vertices as v , v_1 , v_2 , and w . We remark that some special trees occur more than once in Figures 1 and 2. However, for simplicity in the proofs that follow, we assign different names to these special trees.

Let $T_{\text{pdi}} \in A \cup B$ be a special tree and let T be a tree. If T contains a subset U of vertices such that $T[U] \cong T_{\text{pdi}}$ and the degree of every black vertex in $V_B(T_{\text{pdi}})$ equals its degree in T , then we say that the tree T contains T_{pdi} as a *prescribed-degree-induced subtree*, abbreviated *PDI-subtree*. In particular, we note that if T_{pdi} is a PDI-subtree of a tree T , then the degree sequence of the vertices of $V_B(T_{\text{pdi}})$ in T equals the degree sequence of the vertices of $V_B(T_{\text{pdi}})$ in T_{pdi} .

We are now in position to define our family \mathcal{T} .




Fig. 1: The set $A = \{T_1, \dots, T_{15}\}$ of special trees

Definition 4. Let \mathcal{T} be the family of trees that:

- (i) contains all trees of order at most 4,
 - (ii) is closed under the four Operations \mathcal{O}_1 , \mathcal{O}_2 , \mathcal{O}_3 , and \mathcal{O}_4 that are listed below, which extend the tree T' to a tree T by attaching a tree to the vertex $v \in V(T')$, called the *attacher* of T' , and
 - (iii) is closed under the Operations \mathcal{O}_5 , and \mathcal{O}_6 listed below, which extend the tree T' to a tree T by attaching trees to the vertices v_1 and v_2 of T' , called the *attachers* of T' .
- **Operation \mathcal{O}_1 :** Let $T_{\text{pdi}} \in \{T_1, T_2, T_8\}$ be a PDI-subtree of T' and let $v = v(T_{\text{pdi}})$. Add a new vertex u and the edge vu .
 - **Operation \mathcal{O}_2 :** Let $T_{\text{pdi}} \in \{T_4, T_{11}, T_{12}, T_{13}, T_{15}\}$ be a PDI-subtree of T' and let $v = v(T_{\text{pdi}})$. Add a path u_1u_2 to T' and the edge vu_1 .
 - **Operation \mathcal{O}_3 :** Let v be an arbitrary vertex of T' . Add a path $u_1u_2u_3$ to T' and the edge vu_2 .
 - **Operation \mathcal{O}_4 :** Let $T_{\text{pdi}} \in \{T_1, T_2, T_3, T_5, T_6, T_7, T_9, T_{10}\}$ be a PDI-subtree of T' and let $v = v(T_{\text{pdi}})$. Add a path $u_1u_2u_3$ to T' and the edge vu_1 .
 - **Operation \mathcal{O}_5 :** Let $T_{\text{pdi}} \cong T_6$ be a PDI-subtree of T' and let $v_1 = v_1(T_{\text{pdi}})$, $v_2 = v_2(T_{\text{pdi}})$. Add a path u_1u_3 to T' and the edge v_1u_1 , and add a new vertex u_2 and the edge v_2u_2 .
 - **Operation \mathcal{O}_6 :** Let $T_{\text{pdi}} \cong T_{14}$ be a PDI-subtree of T' and let $v_1 = v_1(T_{\text{pdi}})$, $v_2 = v_2(T_{\text{pdi}})$. Remove the edge v_1v_2 , and add a path $u_1u_2u_3$ and the edges v_1u_1 and v_2u_2 .

For $i \in \{1, 2, 4, 5, 6\}$, if T is obtained from T' by applying Operation \mathcal{O}_i to a PDI-subtree T_{pdi} of T' , then we let $X = V(T) \setminus V(T')$ and $T_{\text{pdi}}^{\mathcal{O}_i} = T[V(T_{\text{pdi}}) \cup X]$. Further, we color all vertices of X in $T_{\text{pdi}}^{\mathcal{O}_i}$ black, while the colors of all vertices in the set $V(T_{\text{pdi}})$ remain unchanged.




Fig. 2: The set $B = \{B_1, \dots, B_{10}\}$ of special trees

We shall need the following properties of special trees.

Observation 5. *If T is a special tree, then the vertices of T covered by a square in Figures 3 and 4 form a $\gamma_2(T)$ -set, and the vertices covered by a diamond form an $\alpha_2(T)$ -set.*

The following observation follows readily from the facts that in a rooted tree one can construct a minimum 2-dominating set by “pushing” vertices in the direction of the root, in the sense that if we can replace a vertex in a 2-dominating set by its parent, then we do so; further, we can construct a maximum 2-independent set by “pushing” vertices away from the root as far as possible, in the sense that if we can replace a vertex in a 2-independent set by its children, then we do so.

Observation 6. *Let T' be a tree that contains a PDI-subtree T_{pdi} , and let D' be a $\gamma_2(T')$ -set and S' an $\alpha_2(T')$ -set. Let $D_{\text{pdi}} = D' \cap V(T_{\text{pdi}})$ and $D_{\text{pdi}}^B = D' \cap V_B(T_{\text{pdi}})$, and let $S_{\text{pdi}} = S' \cap V(T_{\text{pdi}})$ and $S_{\text{pdi}}^B = S' \cap V_B(T_{\text{pdi}})$. Then the sets D' and S' can be chosen so that the following hold.*

- (a) *If $T_{\text{pdi}} \in \{T_1, T_2, T_4, T_8, T_{11}, T_{15}, B_3, B_7\}$, then the sets D_{pdi} and S_{pdi} consist of the square and diamond vertices, respectively, of T_{pdi} illustrated in Figures 3 and 4.*
- (b) *If $T_{\text{pdi}} \in \{T_5, T_6, T_9, T_{14}, B_1\}$, then the sets D_{pdi} and S_{pdi}^B consist of the square and diamond vertices, respectively, of T_{pdi} illustrated in Figures 3 and 4.*
- (c) *If $T_{\text{pdi}} \in \{T_{12}, T_{13}, B_8, B_9, B_{10}\}$, then the sets D_{pdi}^B and S_{pdi} consist of the square and diamond vertices, respectively, of T_{pdi} illustrated in Figures 3 and 4.*
- (d) *If $T_{\text{pdi}} \in \{T_{10}, B_2, B_4, B_5, B_6\}$, then the sets D_{pdi}^B and S_{pdi}^B consist of the square and diamond vertices, respectively, of T_{pdi} illustrated in Figures 3 and 4.*
- (e) *If $T_{\text{pdi}} \in \{T_3, T_4, T_7, T_{11}, T_{12}, T_{13}, T_{15}\}$ and $v = v(T_{\text{pdi}})$, then $\alpha_2(T_{\text{pdi}} - v) = \alpha_2(T_{\text{pdi}}) - 1$.*

4 Proof of Theorem 3

In this section, we will prove our main result, namely Theorem 3. We first will present some preliminary results that we will need for our proof.

Observation 7. *Every leaf of a graph G is in every $\gamma_2(G)$ -set and there is an $\alpha_2(G)$ -set containing all leafs of G .*

We now prove that performing Operations $\mathcal{O}_1, \mathcal{O}_2, \dots, \mathcal{O}_6$ maintains the difference between the 2-domination and the 2-independence numbers.




Fig. 3: The set $A = \{T_1, \dots, T_{15}\}$ of special trees

Lemma 8. *If T is obtained from an arbitrary tree T' by applying one of the Operations \mathcal{O}_i for some $i \in [6]$, then $\alpha_2(T) - \gamma_2(T) = \alpha_2(T') - \gamma_2(T')$.*

Proof: Let $c = \alpha_2(T') - \gamma_2(T')$. By Theorem 1, we have $c \geq 0$. We prove that $\alpha_2(T) - \gamma_2(T) = c$. Let D' be a $\gamma_2(T')$ -set and let S' be an $\alpha_2(T')$ -set. Let v be the attacher in T' if T is obtained from T' using Operation \mathcal{O}_1 , \mathcal{O}_2 , \mathcal{O}_3 , or \mathcal{O}_4 , and let v_1 and v_2 be the attachers in T' if T is obtained from T' using Operation \mathcal{O}_5 or \mathcal{O}_6 . If T is obtained from T' by Operation \mathcal{O}_1 , \mathcal{O}_2 , \mathcal{O}_4 , \mathcal{O}_5 , or \mathcal{O}_6 , then let T_{pdi} be the PDI-subtree of T' used to construct the tree T , where $v = v(T_{\text{pdi}})$, $v_1 = v_1(T_{\text{pdi}})$ and $v_2 = v_2(T_{\text{pdi}})$. Further, let $D_{\text{pdi}} = D' \cap V(T_{\text{pdi}})$ and $D_{\text{pdi}}^B = D' \cap V_B(T_{\text{pdi}})$, and let $S_{\text{pdi}} = S' \cap V(T_{\text{pdi}})$ and $S_{\text{pdi}}^B = S' \cap V_B(T_{\text{pdi}})$. By Observation 6, the sets D' and S' can be chosen so that properties (a)-(e) in the statement of the observation hold. Let D be a $\gamma_2(T)$ -set and let S be an $\alpha_2(T)$ -set. We consider six cases, depending on the operation applied to T' in order to obtain the tree T . In all cases, we show that $\alpha_2(T) - \gamma_2(T) = c$.

Case 1. T is obtained from T' by Operation \mathcal{O}_1 . In this case, $T_{\text{pdi}} \in \{T_1, T_2, T_8\}$. Let u be the vertex added to T' and uv be the edge added to T' to obtain T . By Observation 7, we have $u \in D$. By Observation 6(a), the sets D and S can be chosen so that $D \cap V(T_{\text{pdi}})$ and $S \cap V(T_{\text{pdi}})$ are the sets of square and diamond vertices, respectively, of T_{pdi} illustrated in Figure 3, noting that in this case the vertex $v = v(T_{\text{pdi}})$ is the white vertex of T_{pdi} . This implies that either $v \in D$, or $v \notin D$ and v is dominated twice by the vertices of $D \setminus \{u\}$. In both cases, the set $D \setminus \{u\}$ is a 2-dominating set of T' . Therefore, $\gamma_2(T') \leq \gamma_2(T) - 1$. It is easy to observe that every 2-dominating set of T' can be extended to a 2-dominating set of T by adding to it the vertex u , implying that $\gamma_2(T) \leq \gamma_2(T') + 1$. Consequently, $\gamma_2(T) = \gamma_2(T') + 1$. Further, we note that $v \notin S$. Therefore $u \in S$, and $S \setminus \{u\}$ is a 2-independent set of T' implying that $\alpha_2(T') \geq \alpha_2(T) - 1$. By




Fig. 4: The set $B = \{B_1, \dots, B_{10}\}$ of special trees

Observation 6(a) we note that the vertex v does not belong to the $\alpha_2(T')$ -set S' . Thus, S' can be extended to a 2-independent set of T by adding to it the vertex u , implying that $\alpha_2(T) \geq \alpha_2(T') + 1$. Consequently, $\alpha_2(T) = \alpha_2(T') + 1$. Thus, $\alpha_2(T) - \gamma_2(T) = \alpha_2(T') - \gamma_2(T') = c$.

Case 2. T is obtained from T' by Operation \mathcal{O}_2 . In this case, $T_{\text{pdi}} \in \{T_4, T_{11}, T_{12}, T_{13}, T_{15}\}$. Let u_1u_2 be the path added to T' and vu_1 the edge added to T' to obtain T . Since u_3 is a leaf of T , we note that $u_3 \in D$. By Observation 6(a) and 6(c), both sets D' and S' contain the vertex $v = v(T_{\text{pdi}})$. The set $D' \cup \{u_2\}$ and $S' \cup \{u_2\}$ are therefore 2-dominating and 2-independent sets, respectively, of T , implying that $\gamma_2(T) \leq |D'| + 1 = \gamma_2(T') + 1$ and $\alpha_2(T) \geq |S'| + 1 = \alpha_2(T') + 1$. We now consider the sets D and S . Necessarily, $u_2 \in D$. If $u_1 \in D$, then we can replace it with the vertex v . If $u_1 \notin D$, then $v \in D$ in order to dominate u_1 twice. Hence, we may choose D so that $D \cap \{v, u_1, u_2\} = \{v, u_2\}$. Therefore, $D \setminus \{u_2\}$ is a 2-dominating set in T' , and so $\gamma_2(T') \leq \gamma_2(T) - 1$. Consequently, $\gamma_2(T) = \gamma_2(T') + 1$. We can always choose S so that $u_2 \in S$. If $u_1 \notin S$, then $S \setminus \{u_1\}$ is a 2-independent set in T' , implying that $\alpha_2(T') \geq |S| - 1 = \alpha_2(T) - 1$. Suppose that $u_1 \in S$. Then, $v \notin S$, and so $S \setminus \{u_1, u_2\}$ is a 2-independent set of $T' - v$. By Observation 6(e), $\alpha_2(T') = \alpha_2(T' - v) + 1 \geq (|S| - 2) + 1 = |S| - 1 = \alpha_2(T) - 1$. In both cases, $\alpha_2(T') \geq \alpha_2(T) - 1$. Consequently, $\alpha_2(T) = \alpha_2(T') + 1$. Thus, $\alpha_2(T) - \gamma_2(T) = \alpha_2(T') - \gamma_2(T') = c$.

Case 3. T is obtained from T' by Operation \mathcal{O}_3 . In this case, the attacher v is an arbitrary vertex of T' . Let $u_1u_2u_3$ be the path added to T' and vu_2 the edge added to T' to obtain T . Since u_1 and u_3 are leaves of T , we note that $\{u_1, u_3\} \subset D$. If $u_2 \in D$, then we can simply replace u_2 in D by the vertex v . Hence, we may assume that $u_2 \notin D$. The set $D \setminus \{u_1, u_3\}$ is therefore a 2-dominating set of T' , and so $\gamma_2(T') \leq \gamma_2(T) - 2$. Every 2-dominating set of T' can be extended to a 2-dominating set of T by adding to it the leaves u_1 and u_3 , implying that $\gamma_2(T) \leq \gamma_2(T') + 2$. Consequently, $\gamma_2(T) = \gamma_2(T') + 2$. Every 2-independent set of T' can be extended to a 2-independent set of T by adding to it the leaves u_1 and u_3 , implying that $\alpha_2(T) \geq \alpha_2(T') + 2$. Suppose that $u_2 \in S$. Then, at most one of u_1 and u_3 belong to S . Renaming u_1 and u_3 if necessary, we may assume that $u_1 \notin S$. In this case, we can simply replace u_2 in S with u_1 . Hence, we may assume that $u_2 \notin S$, and so $\{u_1, u_3\} \subset S$. The set $S \setminus \{u_1, u_3\}$ is therefore a 2-independent set of T' , and so $\alpha_2(T') \geq \alpha_2(T) - 2$. Consequently, $\alpha_2(T) = \alpha_2(T') + 2$. Thus, $\alpha_2(T) - \gamma_2(T) = \alpha_2(T') - \gamma_2(T') = c$.

Case 4. T is obtained from T' by Operation \mathcal{O}_4 . In this case, $T_{\text{pdi}} \in \{T_1, T_2, T_3, T_5, T_6, T_7, T_9, T_{10}, T_{14}\}$. Let $P: u_1u_2u_3$ be the path added to T' and vu_1 the edge added to T' to obtain T . Every 2-dominating set of T' can be extended to a 2-dominating set of T by adding to it vertices u_2 and u_3 , implying that $\gamma_2(T) \leq \gamma_2(T') + 2$. Since u_3 is a leaf of T , we have $u_3 \in D$. If $u_2 \in D$, then we can simply replace u_2 in D by u_1 . If $u_2 \notin D$, then $u_1 \in D$ in order to dominate the vertex u_2 twice. Thus, we may assume that $D \cap \{u_1, u_2, u_3\} = \{u_1, u_3\}$. Suppose $T_{\text{pdi}} \in \{T_1, T_2, T_5, T_6, T_9, T_{10}, T_{14}\}$. By Observation 6(a), 6(b), and 6(d), the set D can be chosen so that $D \cap V(T_{\text{pdi}})$ are the square vertices of T_{pdi} illustrated in Figure

3, noting that in this case the vertex $v = v(T_{\text{pdi}})$ is the white vertex of T_{pdi} . This implies that either $v \in D$ or $v \notin D$ and v is dominated twice by vertices of $D \setminus \{u_1, u_3\}$. In both cases, the set $D \setminus \{u_1, u_3\}$ is a 2-dominating set of T' . Therefore, $\gamma_2(T') \leq |D| - 2 = \gamma_2(T) - 2$. Consequently, $\gamma_2(T) = \gamma_2(T') + 2$. Suppose $T_{\text{pdi}} \in \{T_3, T_7\}$. If $v \notin D$, then since the two neighbors of v in T_{pdi} have degree at most 2, the set D must contain both neighbors of v in T_{pdi} , for otherwise a neighbor of v not in D would not be dominated twice by vertices of D , a contradiction. Therefore, either $v \in D$ or $v \notin D$ and both neighbors of v belong to D . In both cases, the set $D \setminus \{u_1, u_3\}$ is a 2-dominating set of T' . Therefore, $\gamma_2(T') \leq \gamma_2(T) - 2$. Consequently, $\gamma_2(T) = \gamma_2(T') + 2$. If $u_1 \in S$, then at most one of u_2 and u_3 belong to S , and in this case we can simply replace the two vertices of P that belong to S with the vertices u_2 and u_3 . If $u_1 \notin S$, then $\{u_2, u_3\} \subset S$. Hence, we may assume that $S \cap \{u_1, u_2, u_3\} = \{u_2, u_3\}$. The set $S \setminus \{u_2, u_3\}$ is therefore a 2-independent set of T' , and so $\alpha_2(T') \geq \alpha_2(T) - 2$. Every 2-independent set of T' can be extended to a 2-independent set of T by adding to it the vertices u_2 and u_3 , implying that $\alpha_2(T) \geq \alpha_2(T') + 2$. Consequently, $\alpha_2(T) = \alpha_2(T') + 2$. Thus, $\alpha_2(T) - \gamma_2(T) = \alpha_2(T') - \gamma_2(T') = c$.

Case 5. T is obtained from T' by Operation \mathcal{O}_5 . In this case, $T_{\text{pdi}} \cong T_6$. Let u_1u_3 be the path added to T' and u_2 the new vertex added to T' , and let v_1u_1 and v_2u_2 be the two edges added to T' to obtain T . Since u_2 and u_3 are leaves of T , we note that $\{u_2, u_3\} \subset D$. If $u_1 \in D$, then we can simply replace u_1 in D by v_1 . If $u_1 \notin D$, then $v_1 \in D$ in order to dominate u_1 twice. Therefore, we can choose D so that $D \cap \{u_1, u_2, u_3, v_1\} = \{u_2, u_3, v_1\}$. Let $v_1w_1w_2w_3v_2$ be the (v_1, v_2) -path. If $w_1 \in D$, we can replace w_1 in D by w_2 . If $w_1 \notin D$, then $w_2 \in D$ in order to dominate w_1 twice. Hence, we can choose D so that $D \cap \{w_1, w_2\} = \{w_2\}$. If $w_3 \in D$, we can replace w_3 in D by v_2 . If $w_3 \notin D$, then $v_2 \in D$ in order to dominate w_3 twice. Hence, we can choose D so that $D \cap \{w_3, v_2\} = \{v_2\}$. The set $D \setminus \{u_2, u_3\}$ is therefore a 2-dominating set of T' , implying that $\gamma_2(T') \leq |D| - 2 = \gamma_2(T) - 2$. By Observation 6(b), the set D' contains the vertex v_1 , and can therefore be extended to a 2-dominating set of T by adding to it the leaves u_2 and u_3 , implying that $\gamma_2(T) \leq |D'| + 2 = \gamma_2(T') + 2$. Consequently, $\gamma_2(T) = \gamma_2(T') + 2$. If $v_1 \in S$, then at most one of u_1 and u_3 belong to S , and in this case we can replace the vertices in the set $\{u_1, u_3, v_1\}$ that belong to S with the vertices u_1 and u_3 . If $v_1 \notin S$, then $\{u_1, u_3\} \subset S$. Hence, we may assume that $S \cap \{u_1, u_3, v_1\} = \{u_1, u_3\}$. If $w_1 \notin S$, then $w_3 \in S$ and we can replace w_3 in the set S with w_1 . Hence, we may assume that $w_1 \in S$. If $w_2 \notin S$, then $w_3 \in S$ and we can replace w_3 in the set S with w_2 . Hence, we may assume that $w_2 \in S$ and $w_3 \notin S$. If $u_2 \notin S$, then $v_2 \in S$ and we can replace v_2 in the set S with u_2 . Hence, we may assume that $u_2 \in S$. With these assumptions, we note that both cases $v_2 \in S$ and $v_2 \notin S$ may possibly occur. However in both cases, the set $(S \setminus \{u_1, u_2, u_3, w_2\}) \cup \{v_1, w_3\}$ is a 2-independent set of T' , and so $\alpha_2(T') \geq |S| - 2 = \alpha_2(T) - 2$. By Observation 6(b), we note that $S' \cap \{v_1, w_1, w_2, w_3\} = \{v_1, w_1, w_3\}$. We note that both cases $v_2 \in S'$ and $v_2 \notin S'$ may possibly occur. However in both cases, the set $(S' \setminus \{v_1, w_3\}) \cup \{u_1, u_2, u_3, w_2\}$ is a 2-independent set of T , implying that $\alpha_2(T) \geq |S'| + 2 = \alpha_2(T') + 2$. Consequently, $\alpha_2(T) = \alpha_2(T') + 2$. Thus, $\alpha_2(T) - \gamma_2(T) = \alpha_2(T') - \gamma_2(T') = c$.

Case 6. T is obtained from T' by Operation \mathcal{O}_6 . In this case, $T_{\text{pdi}} = T_{14}$. Let $u_1u_2u_3$ be the path added to T' , and let v_1u_1 and v_2u_2 be the two edges added to T' to obtain T . Since u_3 is a leaf of T , we note that $u_3 \in D$. Further, since $(V(T_{\text{pdi}}) \setminus v_1) \cup \{u_2\}$ induces a PDI-subgraph T_{14} , $u_2 \in D$ by Observation 6. In order to dominate some vertex u_1 , $u_1 \in D$, $v_1 \in D$ twice. We can replace u_1 in the set D by v_1 if $u_1 \in D$. Hence, we may assume that $v_1 \in D$. But now, v_1 dominates v_2 in T' . Therefore, $D \setminus \{u_2, u_3\}$ is a 2-dominating set of T' and $\gamma_2(T') \leq |D| - 2 = \gamma_2(T) - 2$. By Observation 6(b), $v_1 \in D'$. Now, $D' \cup \{u_2, u_3\}$ is a 2-dominating set of T , implying $\gamma_2(T) \leq \gamma_2(T') + 2$. Consequently, $\gamma_2(T) = \gamma_2(T') + 2$. We note that by definition, $|S \cap \{u_1, u_2, u_3\}| \leq 2$. Hence, $S \setminus \{u_1, u_2, u_3\}$ is a 2-independent set of T' , and further $\alpha_2(T') \geq \alpha_2(T) - 2$. By Observation 6(b), we note that $v_2 \notin S'$. Therefore, $S' \cup \{u_2, u_3\}$ is a 2-independent set of T , implying $\alpha_2(T) \geq |S'| + 2 = \alpha_2(T') + 2$. Consequently, $\alpha_2(T) = \alpha_2(T') + 2$.

Thus, $\alpha_2(T) - \gamma_2(T) = \alpha_2(T') - \gamma_2(T') = c$. \square

We are now in position to prove the following lemma.

Lemma 9. *Every tree of the family \mathcal{T} is a (γ_2, α_2) -tree.*

Proof: We proceed by induction on the order $n \geq 1$ of a tree $T \in \mathcal{T}$. Every tree of order at most 4 is a (γ_2, α_2) -tree. This establishes the base. For the inductive hypothesis, let $n \geq 5$ and assume that every tree of order less than n that belongs to the family \mathcal{T} is a (γ_2, α_2) -tree. Let T be a tree of order n that belongs to the family \mathcal{T} . Then there exists a sequence T_0, T_1, \dots, T_k of trees such that T_0 is a tree of order at most 4, $T_k = T$, and for $i \in [k]$, the tree T_i can be obtained from the tree T_{i-1} by one of the Operations $\mathcal{O}_1, \mathcal{O}_2, \dots, \mathcal{O}_6$. Let $T' = T_{k-1}$ and note that $T' \in \mathcal{T}$ and T' has order less than n . Applying the inductive hypothesis to the tree T' , we have that T' is a (γ_2, α_2) -tree. Thus, $\gamma_2(T') = \alpha_2(T')$. The tree T can be obtained from T' by applying one of the Operations $\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3, \mathcal{O}_4, \mathcal{O}_5$, or \mathcal{O}_6 . Therefore, by Lemma 8, $\alpha_2(T) - \gamma_2(T) = \alpha_2(T') - \gamma_2(T') = 0$; or, equivalently, $\gamma_2(T) = \alpha_2(T)$. Therefore, T is a (γ_2, α_2) -tree. \square

Lemma 10. *Every (γ_2, α_2) -tree belongs to the family \mathcal{T} .*

Proof: We show that if T is a (γ_2, α_2) -tree, then $T \in \mathcal{T}$. We proceed by induction on the order $n \geq 1$ of a (γ_2, α_2) -tree T . If $n \leq 4$, then $T \in \mathcal{T}$. This establishes the base case. For the inductive hypothesis, let $n \geq 5$ and assume that every (γ_2, α_2) -tree of order less than n belongs to the family \mathcal{T} . Let T be a (γ_2, α_2) -tree of order n . We show that $T \in \mathcal{T}$. If T is a star, then this is immediate since T can be obtained from a path P_3 by repeated applications of Operation \mathcal{O}_1 . Hence, we may assume that $\text{diam}(T) \geq 3$.

We will frequently use the following three facts throughout the remaining proof.

Fact 10.1. *If T contains a set U of vertices such that T can be obtained from the tree $T - U$ by applying Operation \mathcal{O}_i for some $i \in [5]$, then $T \in \mathcal{T}$.*

Proof: Let U be a set of vertices of T , and let $T' = T - U$. If T can be obtained from the tree T' by applying Operation \mathcal{O}_i for some $i \in [5]$, then, by Lemma 8, $\alpha_2(T) - \gamma_2(T) = \alpha_2(T') - \gamma_2(T')$. By supposition, T is a (γ_2, α_2) -tree, and so $\alpha_2(T) - \gamma_2(T) = 0$. Therefore, $\alpha_2(T') - \gamma_2(T') = 0$, and so T' is a (γ_2, α_2) -tree. Applying the inductive hypothesis to T' , the tree $T' \in \mathcal{T}$. Since T can be restored by applying Operation \mathcal{O}_i to the tree $T' \in \mathcal{T}$, the tree $T \in \mathcal{T}$. \square

As a consequence of Fact 10.1, we have the following result.

Fact 10.2. *If T contains a PDI-subtree $T_{\text{pdi}}^{\mathcal{O}_i}$ for some $i \in \{1, 2, 4, 5, 6\}$, then $T \in \mathcal{T}$.*

Proof: Clearly, the result is true for $i \in \{1, 2, 4, 5\}$ by Fact 10.1. Hence, we may assume that T contains a PDI-subtree $T_{14}^{\mathcal{O}_6}$. Therefore, by definition, there is a tree T' such that T is obtained by applying Operation \mathcal{O}_6 to T' . Moreover, by Lemma 8, $\alpha_2(T') - \gamma_2(T') = \alpha_2(T) - \gamma_2(T)$, and so we conclude that T' is a (γ_2, α_2) -tree. By induction, $T' \in \mathcal{T}$, implying $T \in \mathcal{T}$. \square

Fact 10.3. *Let U be a set of vertices in T , and let $T' = T[U]$. Let D' be a 2-dominating set in T' and let S' be a 2-independent set in T' . If $|D'| < |S'|$ and $\partial(U) \cap S' = \emptyset$, then we have a contradiction to the choice of T .*

Proof: Suppose that $|D'| < |S'|$ and $\partial(U) \cap S' = \emptyset$. Every 2-dominating set of $T - U$ can be extended to a 2-dominating set of T by adding to it the set D' , implying that $\gamma_2(T) \leq \gamma_2(T - U) + |D'|$. Since no vertex in S' is adjacent to a vertex in $V(T) \setminus U$, every 2-independent set in $T - U$ can be extended to a 2-independent set in T by adding to it the set S' . Thus, $\alpha_2(T) \geq \alpha_2(T - U) + |S'|$. By Theorem 1, $\gamma_2(T - U) \leq \alpha_2(T - U)$.

Thus, $\alpha_2(T) = \gamma_2(T) \leq \gamma_2(T - U) + |D'| \leq \alpha_2(T - U) + |D'| < \alpha_2(T - U) + |S'| \leq \alpha_2(T)$, a contradiction. \square

We proceed further with the following series of claims.

Claim 10.4. *If T has a support vertex adjacent to at least three leaves, then $T \in \mathcal{T}$.*

Proof: Let v be a support vertex adjacent to at least three leaves. Let u be an arbitrary leaf adjacent to v and let $U = \{u\}$. Since T can be obtained from the tree $T - U$ by applying Operation \mathcal{O}_1 with the vertex v as the attacher in $T - U$, Fact 10.1 implies that $T \in \mathcal{T}$. \square

By Claim 10.4, we may assume that every support vertex of T is adjacent to at most two leaves, for otherwise the desired result follows. By our earlier assumptions, $\text{diam}(T) \geq 3$. Let P be a longest path in T and suppose that P is an (r_1, r_2) -path. Necessarily, r_1 and r_2 are leaves in T . Renaming r_1 and r_2 if necessary, we may assume that the degree of the support vertex adjacent to r_1 is at most the degree of the support vertex adjacent to r_2 . We now let $r = r_1$ and root the tree T at the vertex r .

We call a vertex of degree at least 2 in T a *large vertex*. Let \mathcal{L} be the set of large vertices in T . For each vertex $w \in \mathcal{L}$, let $\ell(w)$ be a leaf at maximum distance from w in T that belongs to the maximal subtree, T_w , at w . In particular, we note that w belongs to the $(r, \ell(w))$ -path.

Claim 10.5. *If $w \in \mathcal{L}$ and T_w is a PDI-subtree T_1 in T , then $T \in \mathcal{T}$.*

Proof: Suppose that $w \in \mathcal{L}$ and $D[w]$ induces the PDI-subtree T_1 . Let $U = D[w]$. In this case, U consists of w and its two children. Since T can be obtained from the tree $T - U$ by applying Operation \mathcal{O}_3 with the parent of w in T as the attacher in $T - U$, Fact 10.1 implies that $T \in \mathcal{T}$. \square

By Claim 10.5, we may assume that if $w \in \mathcal{L}$, then T_w is not a PDI-subtree T_1 in T , for otherwise the desired result follows. We define $\mathcal{B}_0 = \{B_1\}$, $\mathcal{B}_1 = \{B_2\}$, $\mathcal{B}_2 = \{B_3\}$, $\mathcal{B}_3 = \{B_4, B_5, B_6\}$, $\mathcal{B}_4 = \{B_7, B_8\}$, $\mathcal{B}_5 = \{B_9, B_{10}\}$, and $\mathcal{B}_i = \emptyset$ for $i \geq 6$. If T_{pdi} is a PDI-subtree of T and T_{pdi} is isomorphic to a tree in the family \mathcal{B}_i for some $i \geq 0$, then we say that T_{pdi} is a *PDI-subtree of \mathcal{B}_i in T* .

Claim 10.6. *For every vertex $w \in \mathcal{L}$, the subtree T_w is a PDI-subtree of $\mathcal{B}_{d(w, \ell(w))}$ in T or $T \in \mathcal{T}$.*

Proof: We proceed by induction on the distance, $d(w, \ell(w))$, from w to the leaf $\ell(w)$. Suppose that $d(w, \ell(w)) = 1$. In this case, every child of w is a leaf. Since every support vertex in T is adjacent to at most two leaves, T_w is a PDI-subtree B_2 or T_1 . By Claim 10.5, we have $T_w \cong B_2$, implying that T_w is a PDI-subtree of \mathcal{B}_1 in T , where w is the white vertex of B_2 .

Case 1. Assume that $d(w, \ell(w)) = 2$. For each child z of w , we note by induction that T_z is a PDI-subtree of $\mathcal{B}_0 \cup \mathcal{B}_1$ in T . Further, since $d(w, \ell(w)) = 2$, at least one child of w , say x_1 , satisfies $T_{x_1} \cong B_2$. Let y_1 be the leaf adjacent to x_1 . If $d_T(w) = 2$, then $T_w \cong B_3$, implying that T_w is a PDI-subtree of \mathcal{B}_3 in T , where w is the white vertex of B_3 . Hence, we may assume that $d_T(w) \geq 3$, for otherwise the desired result follows. Let x_2 be a child of w different from x_1 .

Now suppose that $T_{x_2} \cong B_2$. Let y_2 be the leaf adjacent to x_2 , and let $U = \{w, x_1, x_2, y_1, y_2\}$. Let $T' = T[U]$. Then, $D' = \{w, y_1, y_2\}$ is a 2-dominating set in T' and $S' = \{x_1, x_2, y_1, y_2\}$ is a 2-independent set in T' . Since $|D'| < |S'|$ and $\partial(U) \cap S' = \emptyset$, we have a contradiction by Fact 10.3. Hence, $T_{x_2} \cong B_1$. We now let $U = \{x_2\}$ and note that $T[\{w, x_1, y_1\}] \cong T_2$ is a PDI-subtree of $T - U$. Thus, T contains a PDI-subtree $T_2^{\mathcal{O}_1}$ with the vertex w as the attacher in this subtree, and so, by Fact 10.2, $T \in \mathcal{T}$.

Case 2. Assume that $d(w, \ell(w)) = 3$. For each child z of w , we note by induction that T_z is a PDI-subgraph of $\mathcal{B}_0 \cup \mathcal{B}_1 \cup \mathcal{B}_2$ in T . Further, since $d(w, \ell(w)) = 3$, at least one child of w , say x_1 , satisfies $T_{x_1} \cong B_3$. If $d_T(w) = 2$, then T_w is a PDI-subtree $B_4 \in \mathcal{B}_3$ in T where w is the white vertex of B_4 . Hence, we may assume that $d_T(w) \geq 3$, for otherwise the desired result follows. If w has a child x_2 such that T_{x_2} is a PDI-subtree B_2 in T where x_2 is the white vertex of B_2 , then let $U = D[x_1]$. Now $T[D(x_2) \cup \{w\}] \cong T_2$

is a PDI-subtree of $T - U$. Therefore, T contains a PDI-subtree $T_2^{\mathcal{O}_3}$ with the vertex w as the attacher in this subtree, and so, by Fact 10.2, $T \in \mathcal{T}$. Thus for each child x of w we have that T_x is a PDI-subtree B_1 or B_3 .

Now assume that w has at least three children, say x_1, x_2 and x_3 . If T_{x_2} and T_{x_3} are two PDI-subtrees B_1 , then let $U = D[x_1]$. Now $T[D(x_2) \cup D(x_3) \cup \{w\}] \cong T_1$ is a PDI-subtree of $T - U$. Thus, T contains a PDI-subtree $T_1^{\mathcal{O}_4}$ with the vertex w as the attacher in this subtree, and so, by Fact 10.2, $T \in \mathcal{T}$. If T_{x_2} and T_{x_3} are two PDI-subtrees such that $T_{x_2} \cong B_1$ and $T_{x_3} \cong B_3$, or $T_{x_2}, T_{x_3} \cong B_3$, then, by defining $U = D[x_1]$, $T[D[x_2] \cup D[x_3] \cup \{w\}]$ is a PDI-subtree T_5 or T_{10} , respectively. Now T contains a PDI-subtree $T_5^{\mathcal{O}_4}$ or $T_{10}^{\mathcal{O}_4}$ with the vertex w as the attacher in this subtree, and so, by Fact 10.2, $T \in \mathcal{T}$. Therefore, w has degree 3 in T and there are at most two children x_1 and x_2 such that T_{x_1} and T_{x_2} are PDI-subtrees satisfying $T_{x_1} \cong B_3$, and $T_{x_2} \cong B_1$ or $T_{x_2} \cong B_3$. Hence, $T_w \cong B_5 \in \mathcal{B}_3$ or $T_w \cong B_6 \in \mathcal{B}_3$, where w is the white vertex in both cases. Hence, T_w is a PDI-subtree of \mathcal{B}_3 in T .

Case 3. Assume that $d(w, \ell(w)) = 4$. For each child z of w , we note by induction that T_z is a PDI-subtree of $\mathcal{B}_0 \cup \mathcal{B}_1 \cup \mathcal{B}_2 \cup \mathcal{B}_3$ in T . Further, since $d(w, \ell(w)) = 4$, at least one child of w , say x_1 , satisfy that T_{x_1} is isomorphic to B_4, B_5 , or B_6 .

Assume that $d_T(w) = 2$. If $T_{x_1} \cong B_4$, then $T_w \cong B_7$, implying that T_w is a PDI-subtree of \mathcal{B}_4 in T , where w is the white vertex of B_7 . Therefore, $T_w \in \{B_5, B_6\}$. Further, since $w \in \mathcal{L}$, we may assume that w is the child of y , and that z is a child of x_1 of degree 2. We note that z has distance 2 to a leaf in T_{x_1} . Now let $U = D[z]$. Depending on whether $T_{x_1} \cong B_5$ or $T_{x_1} \cong B_6$, $T_y - U$ is a PDI-subtree T_3 or T_7 , respectively. Thus, T contains a PDI-subtree $T_3^{\mathcal{O}_4}$ or $T_7^{\mathcal{O}_4}$ with the vertex x_1 as the attacher in this subtree, and so, by Fact 10.2, $T \in \mathcal{T}$. Hence, we may assume that $d_T(w) \geq 3$, for otherwise the desired result follows. Therefore, let x_2 be a child of w different from x_1 .

Recall, for any child z of w , T_z is a PDI-subtree of $\mathcal{B}_0 \cup \mathcal{B}_1 \cup \mathcal{B}_2 \cup \mathcal{B}_3$ in T , and T_{x_1} is a PDI-subtree of $\{B_4, B_5, B_6\}$.

If there is a child, renaming vertices if necessary, say x_2 , of w such that T_{x_2} is a PDI-subtree of $\{B_2, B_4, B_5, B_6\}$ in T , then let $U = \{w\} \cup D[x_1] \cup D[x_2]$. By a simple case analysis and Observation 6, one can readily observe that there is a 2-dominating set D' and a 2-independent set S' in $T[U]$ such that $|D'| < |S'|$ and $\partial(U) \cap S' = \emptyset$. Therefore, we have a contradiction by Fact 10.3, implying that there is a child, say x_2 , distinct from x_1 such that T_{x_2} is a PDI-subtree of $\{B_1, B_3\}$ in T .

Assume that $T_{x_1} \cong B_4$. If $T_{x_2} \cong B_3$, then let $U = D[x_2]$. Thus, $T[D[x_1] \cup \{w\}]$ is a PDI-subtree T_6 , implying that T contains a PDI-subtree $T_6^{\mathcal{O}_4}$ with the vertex w as the attacher in this subtree, and so, by Fact 10.2, $T \in \mathcal{T}$. Therefore, any child of w , distinct from x_1 , is a leaf. If w has at least three children, say x_1, x_2 , and x_3 , then let $U = \{x_3\}$. Now $T[D[x_1] \cup \{x_2, w\}]$ is a PDI-subtree T_8 . Hence, T contains a PDI-subtree $T_8^{\mathcal{O}_1}$ with the vertex w as the attacher in this subtree, and so, by Fact 10.2, $T \in \mathcal{T}$. Therefore, w has two children, namely x_1 and a leaf x_2 . Further, let U be the set of those two vertices in $D(x_1)$ which have largest and second largest distance to x_1 in T . Now $T[(D[x_1] \setminus U) \cup \{x_2, w\}]$ is a PDI-subtree T_4 implying that T contains a PDI-subtree $T_4^{\mathcal{O}_2}$, and so, by Fact 10.2, $T \in \mathcal{T}$.

Assume that $T_{x_1} \cong B_5$. If $T_{x_2} \cong B_3$, then let $U = D(x_2)$ implying that $T[D[x_1] \cup \{x_2, w\}]$ is a PDI-subtree T_{13} in T . Therefore, T contains a PDI-subtree $T_{13}^{\mathcal{O}_2}$ and so, by Fact 10.2, $T \in \mathcal{T}$. Hence, any child of w , distinct from x_1 , is a leaf. If w has at least three children, say x_1, x_2 , and x_3 , then let $U = V(T_w)$. By Observation 6, one can readily observe that there is a 2-dominating set D' and a 2-independent set S' in $T[U]$ such that $|D'| < |S'|$ and $\partial(U) \cap S' = \emptyset$, implying a contradiction by Fact 10.3. Therefore, w has only two children, say x_1 and x_2 , for otherwise the desired result follows. Now $T_w \cong B_8$ and so T_w is a PDI-subtree of \mathcal{B}_4 in T , where w is the white vertex of B_8 .

Assume that $T_{x_1} \cong B_6$. If $T_{x_2} \cong B_1$, then let U be the set of vertices consisting of $\ell(x_1)$ and its support vertex. Now $T[(D[x_1] \setminus U) \cup \{x_2, w\}]$ is a PDI-subtree T_{12} in T . Therefore, T contains a PDI-subtree $T_{12}^{\mathcal{O}_2}$ and so, by Fact 10.2, $T \in \mathcal{T}$. If $T_{x_2} \cong B_3$, then let $U = D(x_1) \cup D[x_2]$. Trivially, $T[U]$ is $3P_3$. Therefore,

any $\gamma_2(T[U])$ -set D' contains all leaves of this forest. Furthermore, any $\gamma_2(T-U)$ -set D^* contains x_1 since it is a leaf in $T-U$. Moreover, there is an $\alpha_2(T[U])$ -set S' such that the leaves in $T[U]$, which are joined by bridges to x_1 or x_2 , are not in S' . Therefore, $S' \cup S^*$ is a 2-independent set for any $\alpha_2(T-U)$ -set S^* . Furthermore, $D' \cup (D^* \setminus \{x_1\})$ is a 2-dominating set, since x_1 and w are dominated by two vertices of D' and one vertex of D' and D^* , respectively. Therefore, since $\gamma_2(T-U) \leq \alpha_2(T-U)$, by Theorem 1, $\gamma_2(T) \leq \gamma_2(T-U) + 5 < \alpha_2(T-U) + 6 \leq \alpha_2(T)$, contradicting the choice of T .

Case 4. Assume that $d(w, \ell(w)) = 5$. For each child z of w , we note by induction that T_z is a PDI-subgraph of $\mathcal{B}_0 \cup \mathcal{B}_1 \cup \mathcal{B}_2 \cup \mathcal{B}_3 \cup \mathcal{B}_4$ in T . Further, since $d(w, \ell(w)) = 5$, at least one child of w , say x_1 , satisfies that T_{x_1} is isomorphic to B_7 or B_8 .

Firstly, assume that x_1 has degree 2. If $T_{x_1} \cong B_7$, then $T_w \cong B_9 \in \mathcal{B}_5$, implying that T_w is a PDI-subtree of \mathcal{B}_4 in T , where w is the white vertex of B_9 . Therefore, $T_{x_1} \cong B_8$. Let $U = D[x_1]$. Now there is a 2-dominating set D' and a 2-independent set S' in $T[U] - x_1$ such that $|D'| < |S'|$. Further, we note that w is a leaf in $T-U$. Therefore, there is a $\gamma_2(T_U)$ -set D^* and an $\alpha_2(T_U)$ -set S^* containing w . Clearly, $S^* \cup S'$ is a 2-independent set in T and $D^* \cup D'$ is a 2-dominating set for $T - x_1$ by definition. Moreover, x_1 is dominated by its leaf neighbor and the vertex w , implying that $D^* \cup D'$ is a 2-dominating set in T . Since $|D^*| \leq |S^*|$, by Theorem 1, $\gamma_2(T) \leq |D^* \cup D'| = |D^*| + |D'| < |S^*| + |S'| = |S^* \cup S'| \leq \alpha_2(T)$, a contradiction to the choice of T . Hence, we may assume that $d_T(w) \geq 3$, for otherwise the desired result follows. Therefore, let x_2 be a child of w different from x_1 .

Assume that $T_{x_1} \cong B_7$. Let $U = D[x_1] \cup D[x_2] \cup \{w\}$. By some simple case analysis and Observation 6, one can readily observe that there is a $\gamma_2(T[U])$ -set D' and an $\alpha_2(T)$ -set S' such that $|D'| < |S'|$ and $\partial(U) \cap S' = \emptyset$ for $T_{x_2} \in \mathcal{B}_5$, a contradiction by Fact 10.3.

Assume $T_{x_1} \cong B_8$. If T_{x_2} is a PDI-subtree of $\{B_2, B_4, B_5, B_6, B_7, B_8\}$ in T , then let $U = \{w\} \cup D[x_1] \cup D[x_2]$. Again, by some simple case analysis and Observation 6, one can readily observe that there is a $\gamma_2(T[U])$ -set D' and an $\alpha_2(T)$ -set S' such that $|D'| < |S'|$ and $\partial(U) \cap S' = \emptyset$, a contradiction by Fact 10.3. If $T_{x_2} \cong B_3$, then let $U = D[x_2]$. Now $T[D[x_1] \cup \{w\}]$ is a PDI-subtree T_{14} . Hence, T contains a PDI-subtree $T_{14}^{\mathcal{O}_4}$ with the vertex w as the attacher in this subtree, and so, by Fact 10.2, $T \in \mathcal{T}$. It remains to consider the case that all children of w different from x_1 are leaves of T . Recall that any support vertex is adjacent to at most two leaves. Further, if w is adjacent to exactly one leaf, then $T_w \cong B_{10}$, implying that T_w is a PDI-subtree of \mathcal{B}_5 in T , where w is the white vertex of B_{10} . On the other hand, if w is adjacent to two leaves, then let U be the set containing $\ell(w)$ and its support vertex. Now $T[D[w] - U]$ is a PDI-subtree T_{15} in T . Therefore, T contains a PDI-subtree $T_{15}^{\mathcal{O}_2}$, and so, by Fact 10.2, $T \in \mathcal{T}$.

Case 5. Assume that $d(w, \ell(w)) = 6$. For each child z of w , we note by induction that T_z is a PDI-subtree of $\mathcal{B}_0 \cup \mathcal{B}_1 \cup \mathcal{B}_2 \cup \mathcal{B}_3 \cup \mathcal{B}_4 \cup \mathcal{B}_5$ in T . Further, since $d(w, \ell(w)) = 6$, at least one child of w , say x_1 , satisfies that T_{x_1} is isomorphic to B_9 or B_{10} .

Assume that $T_{x_1} \cong B_9$. If w has degree 2, then let $U = D[x_1]$. Now there is a 2-dominating set D' and a 2-independent set S' in $T[U - x_1]$ such that $|D'| \leq |S'|$. Moreover, since w is a leaf in $T-U$, there is a $\gamma_2(T-U)$ -set D^* in $T-U$ and an $\alpha_2(T)$ -set S^* containing w . Hence, x_1 is dominated by a vertex of D' and a vertex of D^* . Therefore, $D' \cup D^*$ is 2-dominating in T and $S' \cup S^*$ is 2-independent in T . It follows, by Theorem 1, $\gamma_2(T) \leq |D' \cup D^*| = |D'| + |D^*| < |S'| + |S^*| = |S' \cup S^*| \leq \alpha_2(T)$, a contradiction. Hence, there is a child of w , say x_2 , distinct from x_1 . If T_{x_2} is a PDI-subtree of $\{B_2, B_4, B_5, B_6, B_6, B_7, B_8, B_9\}$ in T , then let $U = D[x_1] \cup D[x_2] \cup \{w\}$. By Observation 6, one can readily observe that in all cases there is a 2-dominating set D' and a 2-independent set S' in $T[U]$ such that $|D'| < |S'|$ and $\partial(S') \cap U = \emptyset$, a contradiction by Fact 10.3. If $T_{x_2} \cong B_1$, then let U be the set of three vertices consisting of $\ell(x_1)$, its support vertex and x_2 . Then $T[(D[x_1] \cup \{w\}) \setminus U]$ is a PDI-subtree T_6 in T . Therefore, T contains a PDI-subtree $T_6^{\mathcal{O}_5}$, and so, by Fact 10.2, $T \in \mathcal{T}$. If $T_z \cong B_3$, then let $U = D[x_2]$. Now $T[D[x_1] \cup \{w\}]$ is a PDI-subtree T_9 in T . Again, $T \in \mathcal{T}$, by Fact 10.2 and since T contains a PDI-subtree $T_9^{\mathcal{O}_4}$.

Assume that $T_{x_1} \cong B_{10}$. If w has degree 2, then let y be the parent of w . Now $D[w] \cup \{y\}$ induces a PDI-subtree $T_{14}^{\mathcal{O}_6}$, and so $T \in \mathcal{T}$ by Fact 10.2. If w has degree at least 3, then there is a child of w , say x_2 , distinct from x_1 . Further, T_{x_2} is a PDI-subtree of $\{B_1, B_2, \dots, B_{10}\}$ in T . Let $U = D[x_1] \cup D[x_2] \cup \{w\}$. By Observation 6, one can readily observe that there is a 2-dominating set D' and a 2-independent set S' in $T - U$ such that $|D'| < |S'|$ and $\partial(S') \cap U = \emptyset$, a contradiction by Fact 10.3.

Note that the proof for $d(w, \ell(w)) = 6$ immediately implies that $\mathcal{B}_i = \emptyset$ for $i \geq 6$. This completes the proof of Claim 10.6 □

We now return to the proof of Lemma 10 for the last time. Recall that T is a (γ_2, α_2) -tree. Further, let w be the support vertex of r . By Claim 10.6, we deduce that T_w is a PDI-subtree of $\mathcal{B}_{d(w, \ell(w))}$ in T or $T \in \mathcal{T}$. Since the desired result follows in the latter case, we assume that T_w is a PDI-subtree of $\mathcal{B}_{d(w, \ell(w))}$. Moreover, $d(w, \ell(w)) \leq 5$ since $\mathcal{B}_i = \emptyset$ for $i \geq 6$. Further, r is an end-vertex of a diametrical path which implies that $\text{diam}(T) \leq 6$. Now let $T_w \cong B_i$. Then, all vertices in $V(T) \setminus V(T_w)$ are leaves adjacent to w . Note that there is at least one leaf, namely the vertex r , and, by Claim 10.4, there are at most two leaves. If $T_w \cong B_1$ or $T_w \cong B_2$, then T is a star, which can be obtained by applying Operation \mathcal{O}_1 to a path P_3 . If T_w is a PDI-subtree of $\{B_3, B_4, B_7, B_9\}$ in T , then, by the choice of r , we have exactly one leaf outside $V(T_w)$ being adjacent to w . Thus, T is a path. Depending on the length of the path, we have $T \in \mathcal{T}$ or $\gamma_2(T) < \alpha_2(T)$ which gives the result or contradicts the choice of T , respectively. If T_w is a PDI-subtree of $\{B_5, B_8, B_{10}\}$ in T , then the degree of w is at least 3, but the support vertex of $\ell(w)$ has degree 2, a contradiction to the choice of r_1 . If $T_w \cong B_6$, then we have a contradiction since r does not lie on a diametrical path. □

As an immediate consequence of Lemmas 9 and 10, we obtain the following characterization of trees with equal 2-domination and 2-independence numbers.

Theorem 3. *A tree is a (γ_2, α_2) -tree if and only if $T \in \mathcal{T}$.*

Proof: Let T be a tree. If $T \in \mathcal{T}$, then by Lemma 9, T is a (γ_2, α_2) -tree. This establishes the sufficiency. If T is a (γ_2, α_2) -tree, then, by Lemma 10, we have $T \in \mathcal{T}$. This proves the necessity. □

References

- [1] Y. Caro and Y. Roditty. A note on the k -domination number of a graph. *Int. J. Math. Math. Sci.*, 13:205–206, 1990.
- [2] M. Chellali, O. Favaron, A. Hansberg, and L. Volkmann. k -domination and k -independence in graphs: A survey. *Graphs Combin.*, 28:1–55, 2012.
- [3] M. Chellali and N. Meddah. Trees with equal 2-domination and 2-independence numbers. *Discuss. Math. Graph Theory*, 32:263–270, 2012.
- [4] E. DeLaViña, C. E. Larson, R. Pepper, and B. Waller. Graffiti.pc on the 2-domination number of a graph. *Proceedings of the Forty First Southeastern International Conference on Combinatorics, Graph Theory and Computing, Congress. Numerantium*, 203:15–32, 2010.
- [5] W. J. Desormeaux, M. A. Henning, D. Rall, and A. Yeo. Relating the annihilation number and the 2-domination number of a tree. *Discrete Math.*, 319:15–23, 2014.
- [6] O. Favaron. On a conjecture of Fink and Jacobson concerning k -domination and k -dependence. *J. Combin. Theory Series B*, 39:101–102, 1985.

- [7] O. Favaron. k -domination and k -independence in graphs. *Ars Combin.*, 25:159–167, 1988.
- [8] O. Favaron, A. Hansberg, and L. Volkmann. On k -domination and minimum degree in graphs. *J. Graph Theory*, 57:33–40, 2008.
- [9] J. F. Fink and M. S. Jacobson. n -domination in graphs. In Y. Alavi and A. J. Schwenk, editors, *Graph Theory with Applications to Algorithms and Computer Science*, pages 283–300. Wiley, New York, 1985.
- [10] J. Fujisawa, A. Hansberg, T. Kubo, A. Saito, M. Sugita, and L. Volkmann. Independence and 2-domination in bipartite graphs. *Australas. J. Combin.*, 40:265–268, 2008.
- [11] A. Hansberg and L. Volkmann. On graphs with equal domination and 2-domination numbers. *Discrete Math.*, 308:2277–2281, 2008.
- [12] T. W. Haynes, S. T. Hedetniemi, and P. J. Slater. *Fundamentals of Domination in Graphs*. Marcel Dekker, Inc. New York, 1998.
- [13] T. W. Haynes, S. T. Hedetniemi, and P. J. Slater, editors. *Domination in Graphs: Advanced Topics*. Marcel Dekker, Inc. New York, 1998.
- [14] M. A. Henning and A. Yeo. *Total domination in graphs*. Springer Monographs in Mathematics, 2013. ISBN: 978-1-4614-6524-9 (Print) 978-1-4614-6525-6 (Online).