

Cell-paths in mono- and bichromatic line arrangements in the plane*

Oswin Aichholzer¹ Jean Cardinal² Thomas Hackl¹ Ferran Hurtado³
Matias Korman⁴ Alexander Pilz¹ Rodrigo I. Silveira⁵
Ryuhei Uehara⁶ Pavel Valtr⁷ Birgit Vogtenhuber¹ Emo Welzl⁸

¹Graz University of Technology, Austria.

²Université Libre de Bruxelles, Belgium.

³Universitat Politècnica de Catalunya, Spain.

⁴National Institute of Informatics (NII), Japan; JST, ERATO, Kawarabayashi Large Graph Project.

⁵Dept. de Matemática & CIDMA, Universidade de Aveiro, Portugal, and Universitat Politècnica de Catalunya, Spain.

⁶Japan Advanced Institute of Science and Technology, Japan.

⁷Charles University, Czech Republic.

⁸ETH Zürich, Switzerland.

received 21st May 2014, revised 13th Nov. 2014, accepted 2nd Dec. 2014.

We prove that the dual graph of any arrangement of n lines in general position always contains a path of length at least $n^2/4$. Further, we show that in every arrangement of n red and blue lines — in general position and not all of the same color — there is a simple path through at least n cells where red and blue lines are crossed alternately.

Keywords: Discrete geometry, Line arrangement, Planar graphs, Hamiltonicity

1 Introduction

Given an arrangement of n red and blue lines in the Euclidean plane, we consider sequences of cells of the arrangement such that consecutive cells share an edge and no cell appears more than once in the sequence. We refer to such sequences as *cell-paths*, or simply *paths*. A path is called *alternating* if the common edges of consecutive cells alternate in color. The *length* of a path is defined to be one less than the number of cells involved. Cell-paths can also be seen as paths in the dual graph of the arrangement, in which there is a vertex for every cell in the arrangement, and an edge connects two vertices when the corresponding cells are adjacent.

*This paper is dedicated to the memory of Ferran Hurtado, to whom we are grateful for constantly proposing beautiful problems, like the ones studied in this paper.

In this paper we study the existence of long paths in both monochromatic and bichromatic arrangements. First we consider the following question: What is the largest number $f(n)$ such that every arrangement of n lines in general position (no three lines share a point and no two lines are parallel) has a cell-path of length at least $f(n)$? It is not hard to argue that $f(n) \in \Omega(n)$, and a quadratic upper bound trivially follows from the number of cells in an arrangement. In Section 2, we prove that $n^2/4 \leq f(n) \leq n^2/3 + O(n)$. We also show that there always exists a path of length at least the number of cells of size four or more.

Originally, our research on cell-paths was motivated by the following bichromatic variant of the question: What is the largest number $p(n)$ such that every arrangement of n blue and red lines in general position and not all of the same color has an alternating path of length at least $p(n)$? Naturally, this problem is more constrained and it is non-trivial to prove even the existence of, say, a linear-size cell-path. In Section 3, we answer this question in the affirmative, proving that $p(n) \geq n$. We also give a simple example showing $p(n) \leq 2n - O(1)$. Hence, there is indeed a significant difference between the monochromatic and the bichromatic setting.

Previous work. Arrangements of lines have been thoroughly studied for decades [7, 10, 13, 14, 15, 23]. For example, properties of monotone paths in the arrangement have been considered (see, e.g., [9]). Substantial emphasis has been put into studying degenerate arrangements in which, e.g., the number of vertices decreases dramatically. Further, the kind of cells one may obtain as well as their extremal number were investigated (for example how many triangles appear in any simple arrangement). In another direction, one can study the graph having as vertices the intersection points of lines, which are adjacent if they are consecutive in one of the lines [6], and investigate its basic properties as a graph, such as edge-colorings or whether it can be decomposed into Hamiltonian cycles (in projective space) [11]. For later use, recall that the cells in any arrangement can be 2-colored chessboard-like, i.e., no two cells with the same color are adjacent [24] (see also [18]).

Not many problems on colored arrangements of lines were considered in the early times, in contrast to the rich (and still growing) literature on combinatorial problems on red and blue points [7, 21]. The first publications considered bichromatic sets of lines and studied the number and distribution of the intersection points of lines with the same color [16, 17, 27]. There is a recent line of research on problems in which lines have to be colored to achieve some property, or are already colored and one looks at the kind of cells that appear, regarding the color of their sides [3, 4, 5]. Our problem on alternating paths adds to this trend.

The present work on bichromatic line arrangements was inspired by a well known (still open) problem on points: Consider a set R of n red points and a set B of n blue points in convex position. Then it is easy to see that a crossing-free alternating Hamiltonian path on $R \cup B$ does not always exist. What is the longest crossing-free alternating path spanned by these points? Akiyama and Urrutia [2] gave an $O(n^2)$ algorithm to find an alternating Hamiltonian path, if it exists. Erdős (see [22]) popularized the problem of studying the value $\ell(n)$ such that a plane alternating path of length at least $\ell(n)$ always exists for any such pair R and B . Abellanas et al. [1], and independently Kynčl et al. [22], proved that $\ell(n) \leq \frac{4}{3}n + O(\sqrt{n})$. Hajnal and Mészáros [20] proved that $\ell(n) \geq n + \Omega(\sqrt{n})$. The gap is still to be closed. Other variations and related problems appear in Mészáros' PhD thesis [25].

Our results are also related to a long-standing open problem about paths in planar graphs. In 1963, Moon and Moser [26] showed that there exist three-connected planar graphs with n vertices in which the longest simple path has length at most $cn^{\log 2 / \log 3}$, where c is some constant. It is conjectured (see [19])

that this is a lower bound as well, hence that every three-connected planar graph contains a path of this length. Our result on $f(n)$ shows that considering dual graphs of arrangements instead, we always get a path of length $\Omega(n^2)$.

Finally note that during the preparation of the final version of the manuscript, we heard about a recent unpublished work on this topic due to Hoffmann, Kleist, and Miltzow, showing that $f(n) = n^2/3 - O(n)$, closing the gap above. They also provide additional results on the bichromatic version.

2 Long paths in arrangements

We begin by considering the following problem: given a set S of $n \geq 2$ lines in general position, our aim is to find bounds on the length of the longest simple path in the dual graph of the arrangement. Let $A(S)$ be the arrangement associated with S , and let G be the dual graph of $A(S)$. Recall that the number of vertices of G is $N = \binom{n+1}{2} + 1$. We define $f(S)$ as the length of the longest simple path in G , and let $f(n) = \min_{|S|=n} f(S)$. In this section we show:

Theorem 2.1 $f(n) = \Theta(n^2)$.

The upper bound on f follows from the fact that the number of vertices of G is quadratic. In Proposition 2.13 below we improve this trivial upper bound with a better constant.

Similarly, we provide two distinct lower bounds for f (Theorem 2.10 and Theorem 2.11). The first one uses Tutte's theorem to show that a graph very similar to G is Hamiltonian, whereas the proof of the second one is elementary and directly gives a simple algorithm to find the path. The difference between them is that the bound on the first one is proportional to the number of large cells (having four or more edges), while the second one depends only on n . Thus, clearly the first one is more useful for the particular cases in which the number of such cells is big.

2.1 A lower bound using Tutte's Theorem

The first lower bound construction uses Tutte's theorem stating that every four-connected planar graph is Hamiltonian. Thus, our aim is to perform local transformations to G to make it four-connected while preserving planarity and removing as few vertices of G as possible.

Recall that, since every vertex of $A(S)$ has degree 4, G is a bipartite quasi-quadrangulation (i.e., every face of G has size four, except for the unbounded one). It is easy to check that G is two-connected and planar. We consider the natural embedding of G given by S , in which every vertex of G is located in the corresponding face of $A(S)$, every edge of G intersects exactly one edge of $A(S)$, and the face corresponding to the unbounded cells of $A(S)$ is the outer face. Recall that a *vertex cutset* of a graph is a set of vertices whose removal disconnects the graph.

Let C be a simple cycle of G . By Jordan's theorem, the removal of C from G decomposes the remaining vertices into two subsets, which we call *outer* and *inner*, where the outer one is the component that contains the unbounded cells of the arrangement. Given a vertex $v \in C$, we define its *inner degree* as the number of neighbors of v that belong to the inner component.

Lemma 2.2 *In any cycle C of G of length $2k$, the inner degree of any vertex of C is at most $k - 2$ and the number of vertices in the inner part is at most $(k - 1)(k - 2)/2$.*

Proof: Consider the set $S(C) \subseteq S$ of lines associated with edges of C . Since C is a cycle, every line in $S(C)$ is intersected by C an even number of times, and at least twice. Hence, there are $|S(C)| \leq k$ such lines. As any vertex v of C is incident to two edges on C , there are exactly $|S(C)| - 2$ lines left that could correspond to edges incident to v and in the inner part of C . Further, the number of vertices in the inner part of C is at most the number of bounded cells of the arrangement formed by $S(C)$ and thus at most $(k - 1)(k - 2)/2$. \square

Let P be a simple path of G whose first and last vertex are incident to the outer face, while all other vertices are interior vertices of G . Then the removal of P splits the remaining vertices of G into two subsets as well. We refer to P as a *separating path* and to the induced subsets as *separated vertex sets*. The proof of the following lemma is similar to the one of Lemma 2.2.

Lemma 2.3 *For any separating path P of G with k vertices, one of the separated vertex sets of G has cardinality at most $(k - 1)(k - 2)/2$.*

Lemma 2.4 *Let C be a simple cycle of G , and let $I(C)$ be the set of vertices in its interior.*

If $I(C) \neq \emptyset$, there exists a simple cycle C' with the same set $I(C)$ in its interior such that no two consecutive vertices of C' have inner degree zero.

For any separating path P that splits the remaining vertices of G into V_1 and V_2 , there exists a separating path P' such that (1) for each side of P' , at least one out of any two consecutive vertices of P' has an emanating edge to that side, (2) P' splits the remaining vertices of G into $V_1' \supseteq V_1$ and $V_2' \supseteq V_2$, and (3) the first and the last vertex of P' have emanating edges to both sides of P' .

Proof: Suppose $I(C) \neq \emptyset$. As all faces of G in the interior of C have size four, at most two consecutive vertices of C can have inner degree zero. Moreover, the only possibility of having two consecutive such vertices is that C uses three consecutive edges of a face in its interior. Iteratively replacing these edges by the fourth edge of this face wherever such a situation occurs, we obtain a simple cycle C' . C' has exactly $I(C)$ in its interior as well, and no two consecutive vertices of C' have inner degree zero.

Similarly, in a separating path P , two consecutive vertices without emanating edges on one side can occur only if P uses three consecutive edges of a face on that side. Replacing all such occurrences on both sides gives the claimed properties. \square

Lemma 2.5 *All vertex cutsets of size two of G consist of the two neighbors of a degree-two vertex.*

Proof: Consider a vertex cutset $C = \{c_1, c_2\}$ of size two and the at least two resulting sets V_1, V_2 of the remaining vertices of G . First let one of the sets, w.l.o.g. V_1 , contain only inner vertices of G . It follows that V_1 is the interior of a cycle of G and the vertices of this cycle are not contained in V_1 . By Lemma 2.4, there exists a cycle, also with V_1 in its interior such that no two consecutive vertices of that cycle have inner degree zero. Further, for V_1 not to be empty, this cycle has to have a length of at least 6 by Lemma 2.2. Hence, at least three vertices of this cycle have inner degree larger than zero and C is not a cutset.

If both V_1 and V_2 contain vertices of the outer face, then there exists a separating path in G . By Lemma 2.4 (3), c_1 and c_2 have to be the end points (first and last vertex) of this separating path. Therefore, by Lemma 2.4 (1), this separating path contains at most three vertices. Thus, by Lemma 2.3, $\min\{|V_1|, |V_2|\} \leq 1$, implying that c_1 and c_2 are the two neighbors of a degree-two vertex or C is not a cutset. \square




Fig. 1: The Y- Δ transformation.

Lemma 2.6 *If C is a vertex cutset of size three of G where one of the separated sets does not contain any vertex of the outer face, then C consists of the three neighbors of a degree-three vertex.*

Proof: Consider a minimal cutset C of size three and let V_1 be a connected component that contains only interior vertices of G . Recall the first part of the previous proof. By Lemma 2.4, there exists a separating cycle with V_1 in its interior such that no two consecutive vertices of that cycle have inner degree zero. If V_1 contains only one vertex, then this separating cycle has to have a length of at least 6 by Lemma 2.2. Hence, at least three vertices of this cycle have inner degree larger than zero. Thus, C can be a cutset, if V_1 contains only one degree-three vertex. However, if V_1 contains more than one vertex, then the separating cycle has to have a length of at least 8 by Lemma 2.2. Therefore, at least four vertices of this cycle have inner degree larger than zero and C is not a cutset. \square

In order to construct a long path in G , we will use the following well-known result.

Theorem 2.7 (Tutte [28]) *Every four-connected planar graph is Hamiltonian.*

Given a degree-three vertex x in a graph, adjacent to u, v, w , the corresponding Y- Δ transformation consists of removing vertex x , adding edges uv, uw , and vw , and removing any parallel edges. This is illustrated in Figure 1. We define a new graph G' by applying the following two transformations to G , in the given order (see Figure 2):

1. Add an extra vertex v_∞ , and make it adjacent to all vertices of G dual to the unbounded cells of $A(S)$.
2. For all vertices of degree three, perform a Y- Δ transformation. Observe that after adding v_∞ , no two vertices of degree three are adjacent. Hence, the transformation is well-defined.

Lemma 2.8 *G' is a four-connected planar graph, hence it is Hamiltonian.*

Proof: We need to show that G' does not have any cutset of size three or less. We first verify the following claim: if C is a cutset of G' , then $C \setminus \{v_\infty\}$ is a cutset of G . To verify this, we need to show that the vertices of degree three that are eliminated by the Y- Δ transformations do not reconnect the separated vertex sets in G . Each such Y- Δ transformation involves three vertices u, v, w that are pairwise adjacent in G' . Hence, the degree-three vertex of G that was eliminated can be assigned to the same side of the partition in G as the vertices $\{u, v, w\} \setminus C$.

Next, we rule out the existence of a cutset C of size two in G' , thereby showing that G' is at least three-connected. From our observation, $C \setminus \{v_\infty\}$ would be a cutset of G . If v_∞ belongs to C then $C \setminus \{v_\infty\}$ has size one, which is impossible since G is known to be two-connected. Otherwise, we have a cutset of size two in G , which from Lemma 2.5 must consist of the two neighbors of a degree-two vertex




Fig. 2: Transforming the dual graph G into G' .

in G . This vertex, however, after adding v_∞ became of degree three, thus it must have been eliminated by a Y- Δ transformation, hence again C cannot be a cutset in G' .

Now suppose that C is a cutset of size exactly three in G' , and suppose first that C does not contain v_∞ . From our observation, C is also a cutset of G . By Lemma 2.6, the separated sets either all contain vertices of the outer face or only one separated set, containing exactly one vertex of degree three, is cut off by its neighbors. Degree-three vertices of G are eliminated by the Y- Δ transformations and all vertices of the outer face in G are connected by v_∞ in G' . Thus, C cannot be a cutset in G' and we have a contradiction. Hence, we can assume that $v_\infty \in C$, and that $C \setminus \{v_\infty\}$ is a cutset of size two of G . But this case was already ruled out above, hence G' cannot have a cutset of size at most three, and therefore is four-connected. \square

Note that using Chiba and Nishizeki's algorithm [8], a Hamiltonian cycle in G' can be found in linear time. In order to complete the bound, we must provide a lower bound on the number of vertices of G' . Let $N_{>3}$ be the number of vertices of G that have degree larger than 3. Observe that the transformations we have done to G have only affected the vertices of degree two or three. That is, any vertex that has degree 4 or more in G will also be present in G' . In particular, we have that G' has at least $N_{>3}$ vertices.

Lemma 2.9 $N_{>3} \geq n^2/6 - 5n/6 + 1$.

Proof: Recall that G has exactly $N = \binom{n+1}{2} + 1$ vertices. G has no interior vertices of degree two. Furthermore, it is known that the maximum number of degree-three interior vertices in G , that is, the number of bounded triangular cells in the arrangement $A(S)$, is at most $n(n-2)/3$ [12]. Also, the number of exterior vertices of degree two or three in G is at most $2n$. Hence, $N_{>3}$ is at least $n(n+1)/2 + 1 - (n(n-2)/3) - 2n = n^2/6 - 5n/6 + 1$. \square

Theorem 2.10 $f(n) \geq N_{>3}$.

Proof: Consider a Hamiltonian cycle in G' . This cycle can be transformed into a simple path of length at least $N_{>3}$ in G , by eliminating v_∞ and replacing every portion of the cycle using one or two edges




Fig. 3: Obtaining a long path in G from a Hamiltonian cycle in G' (left), using the operations shown on the right.

of a triangle obtained from a Y - Δ transformation by two edges incident to the degree-three vertex (see Figure 3). □

2.2 An improved $\Omega(n^2)$ lower bound

In this section we present an alternative lower bound that performs better than the previous one when $N_{>3}$ is small.

Theorem 2.11 $f(n) \geq n^2/4 + 3n/4$.

Proof: Consider the arrangement $A(S)$ to have a fixed direction, with distinguished topmost and bottommost infinite cells. In order to describe the lower bound construction, we will use the simplified wiring diagram W of $A(S)$ with respect to this direction, as depicted in Fig. 4. We refer the reader to the book by Stefan Felsner for precise definitions and discussions of wiring diagrams [10].

Recall that the horizontal lines in W represent the levels of $A(S)$ (in the given fixed direction). We number the levels from top to bottom with L_1, \dots, L_n . Let R_0 and R_n be the two (one-cell) *outer* rows above L_1 and below L_n , respectively. Let R_i , $1 \leq i \leq n - 1$, be an *inner* row of cells between two levels L_i and L_{i+1} .

Note that the union over the cells in all inner rows does not include the topmost cell (in R_0) and the bottommost cell (in R_n). Further, note that vertical segments in W represent line intersections in $A(S)$, hence two cells are adjacent in the arrangement exactly if they share a horizontal segment in W .

Observation 2.12 *Each interior cell in any inner row R_i , $1 \leq i \leq n - 1$, has at least one neighbor in R_{i-1} , at least one neighbor in R_{i+1} , and altogether at least three neighbors in $R_{i-1} \cup R_{i+1}$*

Proof: If an interior cell had only two neighbors, then it would be bounded by two lines intersecting at least twice, see Fig. 5. □

Let $U_i = R_i \cup R_{i+1}$ for $0 \leq i \leq n - 1$. The subgraph H_i of the dual graph induced by U_i is a caterpillar (a star if $i = 0$ or $i = n - 1$). Let P_i be the longest path in H_i connecting the left infinite cells with the




Fig. 4: An example of an arrangement and its presentations as a wiring diagram and as a simplified wiring diagram. The dual graph is shown in the arrangement and the simplified wiring diagram.




Fig. 5: An interior cell C cannot have two neighbors only.

right infinite cells. See Fig. 6 for an example. It is easy to see that all vertices dual to infinite cells of U_i are in P_i .

We show that the union of the paths P_i contains the dual vertices of all cells of W . Consider an interior cell C in R_i . The vertex dual to C lies on P_i if and only if C has at least two neighbors in R_{i+1} . Similarly, the vertex dual to C lies on P_{i-1} if and only if C has at least two neighbors in R_{i-1} . It follows by Observation 2.12 that the vertex dual to C is in P_i or in P_{i-1} (or in both of them).

Our aim is first to show that we can combine (parts of) the paths P_i into four different longer paths, each of them connecting the topmost cell to the bottommost cell. Then we will prove that at least one of these four paths fulfills the bound claimed in the theorem.




Fig. 6: Example showing the longest paths P_1 (dark dashed) and P_2 (dark dotted) in the subset $R_1 \cup R_2 \cup R_3$ of the example in Fig. 4.

For $i = 1, \dots, n - 1$, let C_i^L be the left infinite cell in R_i and let C_i^R be the right infinite cell in R_i . Denote the topmost cell by $C_0^L = C_0^R$ and the bottommost cell by $C_n^L = C_n^R$.




Fig. 7: The (sub)paths P_i^L and P_i^R of the paths P_i ($i = \{1, 2\}$) of the example in Fig. 6. The non-contained parts of P_1 and P_2 are shown in gray, dashed and dotted, respectively. The infinite cells C_i^L and C_i^R are shaded.

For $1 \leq i \leq n - 2$, let $P_i^L \subseteq P_i$ be the path in H_i from C_i^L to C_{i+1}^R , and let $P_i^R \subseteq P_i$ be the path in H_i from C_i^R to C_{i+1}^L . See Fig. 7 for an example. Note that each of P_i^L and P_i^R traverses all interior cells of P_i plus at least two of its infinite cells. Consider the four infinite cells $C_i^L, C_{i+1}^L, C_i^R, C_{i+1}^R$ of U_i . Observe that one of the two paths P_i^L, P_i^R contains both C_i^L and C_{i+1}^L (in some order), while the other path contains exactly one of the two cells. An analogous statement holds for C_i^R, C_{i+1}^R . Hence, out of the four infinite cells of U_i ($1 \leq i \leq n - 2$), two are contained exactly once and two are contained exactly twice in P_i^L and P_i^R together. Thus, summing the number of infinite cells contained in P_i^L and P_i^R and averaging over the four infinite cells of U_i , we count each such infinite cell at least $3/2$ times.

For $i \in \{0, n - 1\}$, $P_0^L = P_0^R = P_0$ is the path $C_1^L C_0^L C_1^R = C_1^R C_0^R C_1^L$, and $P_{n-1}^L = P_{n-1}^R = P_{n-1}$ is the path $C_{n-1}^L C_n^L C_{n-1}^R = C_{n-1}^R C_n^R C_{n-1}^L$. Trivially, every infinite cell of U_0 and U_{n-1} is visited twice by P_0^L together with P_0^R , and P_{n-1}^L together with P_{n-1}^R , respectively.

We partition the paths P_i^L, P_i^R into four different sets $\mathcal{Q}_k, k \in \{0, 1, 2, 3\}$, defined by:

$$\mathcal{Q}_k := \{P_j^L : (j - k - 1) \equiv 0 \pmod{4}\} \cup \{P_j^R : (j - k + 1) \equiv 0 \pmod{4}\}.$$

See Fig. 8 for an example. Observe that in each \mathcal{Q}_k the paths are disjoint. Moreover, they can easily be connected to form one long path by adding certain edges connecting neighboring infinite cells. See the shaded cells in Fig. 8 for examples.

We define these sets E_k of connecting edges for each \mathcal{Q}_k :

$$E_k := \{C_j^L C_{j+1}^L : (j - k) \equiv 0 \pmod{4}\} \cup \{C_j^R C_{j+1}^R : (j + 2 - k) \equiv 0 \pmod{4}\}.$$

Combining the edges in E_k with the edges in the paths in \mathcal{Q}_k , we get a path Q_k , for each $k \in \{0, 1, 2, 3\}$. What remains to show is that at least one of the four paths Q_k has the desired length.

Recall that P_i^L and P_i^R contain all interior cells of P_i . By Observation 2.12, each interior cell lying in some row R_i has at least two neighbors above or below and therefore lies on both paths P_i^L and P_i^R , or on both paths P_{i-1}^L and P_{i-1}^R (or even on all four paths). Thus, each interior cell lies on at least two of the four paths Q_k . Further, all infinite cells (except for the topmost and bottommost one) in $U_i, 0 \leq i \leq n - 1$, are counted at least $3/2$ times (on average) in P_i^L and P_i^R together. For the topmost and bottommost cell, observe that both are contained in each of the four paths Q_k .




Fig. 8: The four sets \mathcal{Q}_k , $k = 0, 1, 2, 3$, of paths, for the example in Fig. 4. For each set \mathcal{Q}_k , the infinite cells that need to be connected to obtain one long path are shown shaded.

Altogether, it follows that the sum of the number of the cells that are traversed by each Q_k for $k \in \{0, 1, 2, 3\}$ is at least

$$2 \cdot \left(\binom{n+1}{2} + 1 - 2n \right) + 2 \cdot \frac{3}{2} \cdot (2n - 2) + 8 = n^2 + 3n + 4.$$

Thus, by the pigeonhole principle, at least one out of the four paths Q_k visits at least $\left\lceil \frac{n^2+3n+4}{4} \right\rceil$ cells. \square

There exists a family of arrangements with a fixed direction, see Fig. 9 (left), such that the technique described in the proof of Theorem 2.11 results in a path visiting only $\frac{n^2}{4} + n + 2$ cells (if n is a multiple of 4). On the other hand, for the same arrangement with a different direction, see Fig. 9 (right), this technique results in a path visiting $\frac{n^2}{2} - \frac{n}{2} + 4$ cells. An interesting open question is whether we can always find a direction such that we get a better bound.

The proof of Theorem 2.11 clearly also holds for cell-paths in pseudoline arrangements. Since the lower bound of $n(n-2)/3$ of bounded triangular faces as well holds for pseudoline arrangements [12], it can be observed that also Theorem 2.10 remains true for pseudoline arrangements.

2.3 Upper bound

We show that the previous lower bound on the length of the longest simple path in G is within a factor $4/3$ of the optimum.

Proposition 2.13 $f(n) \leq n^2/3 + O(n)$.




Fig. 9: An example where the above proof gives a path of length $\sim n^2/4$, instantiated for $n = 8$. (left) With the given direction of the arrangement our technique yields a path (Q_0 and Q_3) visiting 26 cells. (right) The same arrangement rotated by 90° : the constructed path (Q_1 and Q_3) visits 32 cells.

Proof: It is well-known that the cells of any line arrangement can be properly two-colored, hence that G is a bipartite graph. We will refer to the colors as black and white. Füredi and Palásti [13] give an example of an arrangement of n lines in which there are $n^2/3 + O(1)$ black cells and $n^2/6 + O(n)$ white cells. Hence, there are roughly twice as many black cells as white cells, which is known to be asymptotically tight [18]. Now observe that any simple path or cycle in G will alternately traverse white and black cells, hence cannot have length greater than $n^2/3 + O(n)$. \square

3 Long alternating paths in bichromatic arrangements

In this section we prove the existence of long alternating paths in bichromatic arrangements. First observe that general position is important to allow a positive answer: Assume that $n \geq 4$ and that at least two lines are red and two lines are blue. Take all the n lines in the arrangement to go through a common point, so that all of the red lines have slope between 0 and 1, and all of the blue lines have positive slope larger than 1. Then every alternating path has length at most two. This holds since each cell on an alternating path, except for the first and the last one, has to be *bichromatic*, i.e., has to have a red and a blue edge on its boundary. But the constructed arrangement has only four bichromatic cells and no two of them share an edge.

Further, consider an arrangement of n lines, all blue except for exactly one of them red, in general position. Then the length of the longest alternating path is $2n - O(1)$: In the arrangement (as edge set) there are n red edges which can be used at most once in a path, so in an alternating path at most $2n + 1$ edges can be used; hence the upper bound. For the lower bound we go through the $2n$ cells along the red line, crossing the red line in every other step, and a blue edge for entering a red-line-incident cell not yet visited in the steps in between; this results in a path of length $2n - 1$.

The following lemma directly implies our main result for the stated problem on bichromatic arrangements.

Lemma 3.1 *Any pair of bichromatic cells in an arrangement of red and blue lines in general position is connected by an alternating path.*

Before giving the proof of this statement, we discuss an implication. Consider two antipodal bichromatic infinite cells in an arrangement (“*antipodal*” means that the cells are separated by all n lines). As long as there is at least one line of each color, such a pair of cells has to exist and, by Lemma 3.1, is connected by an alternating path. Clearly, such a path has to cross every line at least once.

Theorem 3.2 *In a set of n blue and red lines — in general position and not all of the same color — there is an alternating path of length n .*

By the example with exactly one red line, the bound in the theorem is asymptotically tight. However, if we require the same number of red and blue lines, we do not know whether longer alternating paths always exist.

3.1 Existence of long alternating paths

In this section we give a proof of Lemma 3.1 (that is, any pair of bichromatic cells in an arrangement of red and blue lines in general position is connected by an alternating path). Recall that the graph underlying our problem is the dual graph G of the arrangement: the $\binom{n+1}{2} + 1$ cells are the vertices of the graph and two vertices are adjacent if their corresponding cells share an edge in the arrangement.

In order to capture the ‘alternating’ property, we can orient the edges of G as follows. Recall that G can be properly 2-colored [24] and denote the two colors by “r-out” and “b-out”, respectively. Now direct the edges of G by directing red edges (i.e., edges dual to red edges in the arrangement) from color r-out to color b-out and blue edges from color b-out to color r-out. This is illustrated in Figure 10. As the partition of cells of $A(S)$ into two color classes is unique, the construction of the directed version of G is unique up to global direction change. Further, it is an easy exercise to verify that every (undirected) alternating path in the arrangement can be directed in one way so that it appears as a directed path in this oriented version of the dual of the arrangement — and vice versa, every directed path is clearly alternating.

Let us fix an arbitrary bichromatic cell z in the arrangement and consider the set of all cells that can be reached from z by a directed path in the just defined directed graph. Let $\text{reach}(z)$ be the closure of the union of all the cells that are reachable by a directed path starting from z in the oriented graph.

Here comes the crucial observation: Let us call a vertex in the arrangement *bichromatic* if it is the intersection of a red and a blue line. Then the four cells incident to a bichromatic vertex form a directed cycle in the oriented graph and thus either all four of them are contained in $\text{reach}(z)$ or none of them is. (Here we implicitly use the fact that the existence of a directed walk — i.e., with repetitions of vertices allowed — from vertex x to vertex y implies the existence of a directed path from x to y .) Therefore, every bichromatic vertex is interior either to $\text{reach}(z)$ or to its complement.

We have now established the following fact.

Lemma 3.3 *Let E be the set of edges of the arrangement that separate $\text{reach}(z)$ from its complement. Then no red edge in E shares a vertex with a blue edge in E .*

We will show that bichromatic vertices cannot be interior to the complement of $\text{reach}(z)$. Thus all of them have to be interior to $\text{reach}(z)$ and therefore, all bichromatic cells are in $\text{reach}(z)$, from which Lemma 3.1 follows. To this end consider a set $X \subseteq E$ of separating edges that form a cycle or a maximal path in the graph of separating edges. A maximal path has to start and end with an edge that extends to




Fig. 10: Orientation of the dual graph G of a bichromatic line arrangement. Cells of b-out-colored vertices are drawn as shaded, while cells of r-out-colored vertex are drawn white. Every bichromatic vertex of the arrangement yields a four-cycle in the oriented graph.

infinity (otherwise such path can be extended). Consequently, the union of edges in X separates the plane into two parts, one part of which contains the seed cell z . Suppose the edges in X are all blue (recall that they all have the same color). Then it is not possible that both sides contain a point on a red line, as otherwise, since we can travel on red lines between these two points, a red line would have to cross the union of edges in X , which we know is not possible, since all of them are blue. Thus, the side that does not contain z must be completely monochromatic, i.e., all cells there are bounded by edges of the same color.

Since every point p in the complement of $\text{reach}(z)$ must have such a cycle or path X separating p from $\text{reach}(z)$, it follows that bichromatic vertices cannot be interior to the complement of $\text{reach}(z)$, as claimed. As argued before, this implies Lemma 3.1.

3.2 Discussion

- The linear bound on the length of the alternating path is probably not tight if an equal number of red and blue lines is required. However, as already stated at the beginning of Section 3, when abandoning the general position assumption, it is easy to construct an example with the same number of red and blue lines where the maximum alternating path has only length two. Obviously, the same bound still holds if no three lines intersect in a common point, but all lines are allowed to be parallel - just place all blue lines vertically followed by all red lines vertically. On the other hand, these arrangements have only one vertex in the first construction and no vertex at all in the second case. Another degenerate example arrangement but with a quadratic number of vertices (and cells) whose longest alternating path has length only $O(n)$ is depicted in Fig. 11. This example has an equal number of red and blue lines, half of the lines are vertical parallel and half of them are horizontal parallel.
- A closer inspection of the given proof shows that we have actually established the following.

Theorem 3.4 *Let C be a set of red and blue simple closed or biinfinite curves, each of which separates the plane into two parts. If the union of red curves is connected, the union of blue curves*




Fig. 11: An arrangement not in general position, but with a quadratic number of vertices, in which the length of the longest alternating path is in $O(n)$ (an example of a long path is shown).

is connected, and no point is contained in more than two of the curves, then any pair of bichromatic cells in the arrangement is connected by an alternating path.

- Further, Lemma 3.1 can be generalized to higher dimensions: Consider two antipodal bichromatic cells in a $(d + 1)$ -dimensional arrangement. Intersect these two cells (and the arrangement) with a hyperplane H . The intersection of the arrangement with H gives a d -dimensional bichromatic arrangement, in which the antipodal cells are connected by induction.

Acknowledgements

This work was initiated during the ComPoSe-Meeting “Workshop on combinatorics of colored point sets” held on Feb. 4–8, 2013 at the University of Seville, Spain. We thank the other participants — J.M. Díaz-Báñez, P. Kamousi, D. Orden, P. Ramos, M. Saumell, W. Steiger, and I. Ventura — for helpful comments. We further thank S. Felsner for valuable discussions and for the example shown in Fig. 9.

Oswin Aichholzer, Alexander Pilz, and Birgit Vogtenhuber were supported by the ESF EUROCORES programme EuroGIGA—CRP ‘ComPoSe’, Austrian Science Fund (FWF): I648-N18. Jean Cardinal was supported by the ESF EUROCORES programme EuroGIGA—CRP ‘ComPoSe’ and F.R.S.-FNRS Grant R70.01.11F. Thomas Hackl was supported by the Austrian Science Fund (FWF): P23629-N18 ‘Combinatorial Problems on Geometric Graphs’. Ferran Hurtado, Matias Korman, and Rodrigo I. Silveira were partially supported by projects MINECO MTM2012-30951, Gen. Cat. DGR2009SGR1040, and ESF EUROCORES programme EuroGIGA—CRP ‘ComPoSe’: MICINN Project EUI-EURC-2011-4306. Matias Korman received support of the Secretary for Universities and Research of the Ministry of Economy and Knowledge of the Government of Catalonia and the European Union. Rodrigo I. Silveira was funded

by Portuguese funds through CIDMA (Center for Research and Development in Mathematics and Applications) and FCT (Fundação para a Ciência e a Tecnologia), within project PEst-OE/MAT/UI4106/2014, and by FCT grant SFRH/BPD/88455/2012. Emo Welzl was supported by EuroCores/EuroGiga/ComPoSe SNF 20GG21 134318/1.

References

- [1] M. Abellanas, A. García, F. Hurtado, and J. Tejel. Caminos alternantes. *Actas X Encuentros de Geometría Computacional* (in Spanish), Sevilla, 2003, 7–12.
- [2] J. Akiyama, J. Urrutia. Simple alternating path problem. *Discrete Math.* 84:101–103, 1990.
- [3] E. Ackerman, J. Pach, R. Pinchasi, R. Radoičić, G. Tóth. A note on coloring line arrangements. *Electr. J. Comb.* 21(2):P2.23.
- [4] S. Bereg, F. Hurtado, M. Kano, M. Korman, D. Lara, C. Seara, R.I. Silveira, J. Urrutia and K. Verbeek. Balanced partitions of 3-colored geometric sets in the plane. Manuscript, 2012. Abstract in *29th EuroCG*, 2013.
- [5] P. Bose, J. Cardinal, S. Collette, F. Hurtado, S. Langerman, M. Korman, and P. Taslakian. Coloring and guarding arrangements. *Discr. Math. & Theor. Comp. Sci.*, 15(3):139-154, 2013.
- [6] P. Bose, H. Everett, and S. Wismath. Properties of Arrangements. *Int. J. Comput. Geom.*, 13:447-462, 2003.
- [7] P. Brass, W. Moser and J. Pach. *Research Problems in Discrete Geometry*. Vieweg Verlag, 2004.
- [8] N. Chiba and T. Nishizeki. The hamiltonian cycle problem is linear-time solvable for 4-connected planar graphs. *J. Algorithms*, 10(2):187-211, 1989.
- [9] A. Dumitrescu. On some monotone path problems in line arrangements. *Comput. Geom.*, 32(1):13–25, 2005.
- [10] S. Felsner. *Geometric Graphs and Arrangements*. Springer, Berlin, 2005.
- [11] S. Felsner, F. Hurtado, M. Noy and I. Streinu. Hamiltonicity and colorings of arrangement graphs. *Discrete Appl. Math.*, 154:2470-2483, 2006.
- [12] S. Felsner and K. Kriegel. Triangles in Euclidean Arrangements. *Discrete Comput. Geom.*, 22:429-438, 1999.
- [13] Z. Füredi and I. Palásti. Arrangements of lines with a large number of triangles. *Proc. Amer. Math. Soc.*, 92:561-566, 1984.
- [14] B. Grünbaum. *Arrangements and Spreads*. Regional Conf. Ser. Math., Amer. Math. Soc., 1972.
- [15] B. Grünbaum. How many triangles? *Geombinatorics*, 8:154-159, 1998.
- [16] B. Grünbaum. Arrangements of colored lines. Abstract 720-50-5, *Notices Amer. Math. Soc.*, 22(1975), A-200.

- [17] B. Grünbaum. Monochromatic intersection points in families of colored lines. *Geombinatorics*, 9:3-9, 1999.
- [18] B. Grünbaum. Two-coloring the faces of arrangements. *Period. Math. Hungar.*, 11(3):181-185, 1980.
- [19] B. Grünbaum and H. Walther. Shortness exponents of families of graphs. *J. Comb.Theory A*, 14:364-385, 1973.
- [20] P. Hajnal, V. Mészáros. A note on noncrossing path in colored convex sets. Accepted to *Discr. Math. & Theor. Comp. Sci.*, 2011.
- [21] A. Kaneko and M. Kano. Discrete geometry on red and blue points in the plane – a survey. In *Discrete and Computational Geometry, The Goodman-Pollack Festschrift*, pp. 551-570, 2003.
- [22] J. Kynčl, J. Pach and G. Tóth. Long alternating paths in bicolored point sets. *Discrete Math.* 308:4315–4322, 2008.
- [23] J. Leañós, M. Lomelí, C. Merino, G. Salazar, and J. Urrutia. Simple Euclidean arrangements with no (≥ 5)-gons. *Discrete Comput. Geom.*, 38:595–603, 2007.
- [24] E. Lucas. *Récréations mathématiques* IV. Paris, 1894
- [25] V. Mészáros. Extremal problems on planar point sets. PhD Thesis, University of Szeged, 2011.
- [26] J.W. Moon and L. Moser. Simple paths on polyhedra. *Pacific J. Math.*, 13:629-631, 1963.
- [27] T. S. Motzkin. Nonmixed connecting lines. Abstract 67T 605, *Notices Amer. Math. Soc.*, 14(1967), p. 837.
- [28] W. T. Tutte. A theorem on planar graphs. *Trans. Amer. Math. Soc.*, 82:99-116, 1956.