

On probe 2-clique graphs and probe diamond-free graphs

Flavia Bonomo^{1,4*} Celina M. H. de Figueiredo^{2†} Guillermo Durán^{1,5,6‡}
 Luciano N. Grippo^{7§} Martín D. Safe^{7¶} Jayme L. Szwarcfiter^{2,3||}

¹CONICET, Argentina

²COPPE, Universidade Federal do Rio de Janeiro, Brazil

³NCE, Universidade Federal do Rio de Janeiro, Brazil

⁴Depto. de Computación, FCEN, Universidad de Buenos Aires, Buenos Aires, Argentina

⁵Depto. de Matemática and Instituto de Cálculo, FCEN, Universidad de Buenos Aires, Buenos Aires, Argentina

⁶Depto. de Ingeniería Industrial, FCFM, Universidad de Chile, Santiago, Chile

⁷Instituto de Ciencias, Universidad Nacional de General Sarmiento, Los Polvorines, Argentina

received 6th Jan. 2014, accepted 14th Mar. 2015.

Given a class \mathcal{G} of graphs, probe \mathcal{G} graphs are defined as follows. A graph G is *probe \mathcal{G}* if there exists a partition of its vertices into a set of *probe vertices* and a stable set of *nonprobe vertices* in such a way that non-edges of G , whose endpoints are nonprobe vertices, can be added so that the resulting graph belongs to \mathcal{G} . We investigate probe 2-clique graphs and probe diamond-free graphs. For probe 2-clique graphs, we present a polynomial-time recognition algorithm. Probe diamond-free graphs are characterized by minimal forbidden induced subgraphs. As a by-product, it is proved that the class of probe block graphs is the intersection between the classes of chordal graphs and probe diamond-free graphs.

Keywords: 2-clique graphs, diamond-free graphs, probe graphs

*Email: fbonomo@dc.uba.ar. Partially supported by ANPCyT PICT 2012-1324, UBACyT Grant 20020130100808BA, and CONICET PIP 112-201201-00450CO (Argentina).

†Email: celina@cos.ufrj.br. Partially supported by the Brazilian research agencies FAPERJ and CNPq

‡Email: gduran@dm.uba.ar. Partially supported by FONDECYT Grant 1140787 and Millennium Science Institute “Complex Engineering Systems” (Chile), and ANPCyT PICT 2012-1324, UBACyT Grant 20020130100808BA, and CONICET PIP 112-201201-00450CO (Argentina)

§Email: lgrippo@ungs.edu.ar. Partially supported by ANPCyT PICT 2012-1324, UBACyT Grant 20020130100808BA, and CONICET PIP 112-201201-00450CO (Argentina)

¶Email: msafe@ungs.edu.ar. Partially supported by ANPCyT PICT 2012-1324, UBACyT Grant 20020130100808BA, and CONICET PIP 112-201201-00450CO (Argentina)

||Email: jayme@nce.ufrj.br. Partially supported by the Brazilian research agencies CAPES and CNPq.

1 Introduction

Let \mathcal{G} be a class of graphs. A graph $G = (V, E)$ is *probe \mathcal{G}* if its vertices can be partitioned into *probe vertices* (P) and a stable set of *nonprobe vertices* (N) in such a way that there exists a set of non-edges F of G , whose endpoints belong to N , such that $G^* = (V, E \cup F)$ is in \mathcal{G} . The graph G^* is defined to be a *probe \mathcal{G} completion* of G , if there is no confusion we just call it a *completion of G* . By (N, P) , we denote a *probe partition* of G . A graph is *interval* if it is the intersection graph of intervals in the real line. Probe interval graphs were defined by Zhang [19]. He introduced them to deal with a problem concerning DNA mapping. A graph $G = (P \cup N, E)$ is said to be a *partitioned graph* if its vertex set is partitioned into two sets: a set P of probe vertices and a stable set N of nonprobe vertices. We say that G is a *partitioned probe \mathcal{G} graph* if there exists a completion $G^* = (P \cup N, E \cup F)$ of G belonging to \mathcal{G} , where all the edges belonging to F have both endpoints in N . We will call *nonpartitioned probe \mathcal{G} graphs* to those probe \mathcal{G} graphs without a prescribed probe partition, when it is clear from the context we will just call them probe \mathcal{G} graphs. In the case of \mathcal{G} being the class of interval graphs, many progresses have been reached in the study of the class of probe \mathcal{G} graphs, known as probe interval graphs. Given a graph a G with a prescribed partition (P, N) of its vertex set into probe (P) and nonprobe (N) vertices, the problem of decided whether there exists a completion of $G = (P \cup N, E)$ into an interval graph $G^* = (P \cup N, E \cup F)$ can be solved in $O(n^2)$ -time [13], this algorithm involves modified PQ-trees. Another algorithm, which deals with this problem, is the one presented in [16] that uses modular decomposition and whose complexity is $O(n + m \log(n))$. In the case of nonpartitioned probe interval graphs, an $O(m^2)$ recognition algorithm was found when the input graph is restricted to cocomparability graphs [3]. In [7] is presented a polynomial-time algorithm to recognize nonpartitioned probe interval graphs, but the complexity of this algorithm is not given.

Recently, several researchers have started to study probe partitioned and nonpartitioned \mathcal{G} graphs for different kind of classes \mathcal{G} of graphs (see e.g., [1], [4], [6] [9], [10]). In this work, we present a structural result on probe 2-clique graphs that leads to a polynomial-time recognition algorithm for this class, and a forbidden induced subgraph characterization of probe diamond-free graphs implying a forbidden induced subgraph characterization for probe block graphs. In [5], it is given a polynomial-time recognition algorithm for deciding if a given graph is probe block. In this article are considered both the partitioned and the nonpartitioned problem. Later, a linear-time algorithm is presented in [15]. The recognition complexity for probe chordal graphs given a partition of its vertices into probe and nonprobe is $O(|P||E|)$ [2], whereas in the case that no partition is given the complexity is $O(|E|^2)$ [2].

For concepts and notation not defined here we refer the reader to [18]. All graphs in this paper are without loops and without multiple edges. Given a graph $G = (V, E)$ and a set $A \subseteq V$, we denote by $G[A]$ the subgraph of G induced by A . A graph H is a *spanning subgraph* of G if $V(H) = V(G)$ and $E(H) \subseteq E(G)$. Spanning subgraphs are not necessarily induced. By C_n we denote a chordless cycle on n vertices. By $d_G(v)$, we denote the degree of a vertex v in G . A *clique* is a set of pairwise adjacent vertices. The *complete graph* is a graph whose vertices are pairwise adjacent. The complete graph on n vertices is denoted by K_n . Complete graphs on three vertices are called *triangles*. By $N_G(v)$, we denote the *neighborhood* of a vertex v in a graph G , if it is clear from the context we may use $N(v)$. By $N_G[v] = N_G(v) \cup \{v\}$, we denote the *closed neighborhood* of v . Given a set of graphs \mathcal{H} , we denote by \mathcal{H} -free the family of those graphs that do not contain any graph in \mathcal{H} as induced subgraph. If \mathcal{H} consists of a single element H , we use H -free for short. A vertex v is said to be *complete (anticomplete)* to a set of vertices A , if v is adjacent (nonadjacent) to all the vertices of A .

The graph $K_4 - e$ is called *diamond* (see the graph H_1 in Figure 1). A graph G is a *block graph* if every connected induced subgraph of it either is a complete graph or has a cut-vertex. In [14] it is proved that block graphs are exactly the diamond-free chordal graphs. A graph is *chordal* if it has no C_n with $n \geq 4$ as an induced subgraph. A graph is *Ptolemaic* if it is chordal and gem-free. (A *gem* is the graph F_2 in Figure 2). A graph is *split* if its vertex set can be partitioned into a clique and a stable set; more details on this class of graphs can be found in [8]. A graph is *2-clique* also referred, in the literature, as co-bipartite) if its vertices can be partitioned into two cliques. This graph class has been studied in connection with circular-arc graphs in [12, 17]

The article is organized as follows. In Section 2, we present a structural result on probe 2-clique graphs. Such a result leads to a polynomial-time algorithm to recognize the class. In Section 3, we present a characterization by forbidden structures for partitioned probe diamond-free graphs, which is used in Section 4 to characterize probe diamond-free graphs by forbidden induced subgraphs. Besides, it is shown that the class of probe block graphs is exactly the intersection between the classes of chordal graphs and probe diamond-free graphs. We conclude the article with a section of further remarks.

2 Probe 2-clique graphs

Denote by $T(G)$ the spanning subgraph of G whose edge set contains precisely the edges that are contained in some triangle of G .

Notice that if G is probe 2-clique then we can add some set of edges C (possibly empty), whose endpoints form a stable set S in G , so that the resulting graph is 2-clique. Since the fact of adding edges to a 2-clique graph gives a 2-clique graph, we could add to G all the edges whose endpoints belong to S and the resulting graph would be also 2-clique. Therefore, if G is probe 2-clique, then there exists a complete graph C' , contained in \overline{G} , having a set of edges $E(C')$ such that $\overline{G} - E(C')$ is bipartite. As every subgraph of a bipartite graph is bipartite, $\overline{G} - E(C')$ is also bipartite. Moreover, for any complete graph C containing C' , $\overline{G} - E(C)$ is bipartite.

Consequently, we have the following results.

Lemma 1 *Let G be a graph such that \overline{G} is triangle-free. Then, G is probe 2-clique if and only if \overline{G} contains an edge e such that $\overline{G} - e$ is bipartite.*

Proof: If G is a graph with no edge the result is satisfied trivially, so we will consider that the graph G contains at least an edge. Let G be a probe 2-clique graph such that \overline{G} is triangle-free. Since G is triangle-free and thus the only complete graphs in \overline{G} are its edges, by the observation above, there exists an edge e such that $\overline{G} - e$ is bipartite. Conversely, if there exists an edge e such that $\overline{G} - e$ is bipartite, then (since the edge e is a clique in \overline{G}) G is probe 2-clique. \square

For instance, the graph $\overline{C_7}$, whose complement graph is clearly triangle-free, is not 2-clique and its complement graph which is not bipartite becomes bipartite by deleting any of its edges and therefore $\overline{C_7}$ is probe 2-clique.

Lemma 2 *Let G be such that \overline{G} contains triangles. Then, G is probe 2-clique if and only if $T(\overline{G})$ has a complete subgraph C such that $\overline{G} - E(C)$ is bipartite.*

Proof: Let G be a graph such that \overline{G} contains a triangle H . Suppose that G is probe 2-clique. Then, \overline{G} has a complete subgraph C such that $\overline{G} - E(C)$ is bipartite. If $|V(C)| \geq 3$, each edge of C is contained in a

triangle of \overline{G} , and C is a complete subgraph of $T(\overline{G})$. If $|V(C)| \leq 2$, then C must be an edge of H , thus a complete subgraph of $T(\overline{G})$. Conversely, suppose $T(\overline{G})$ has a complete subgraph C such that $\overline{G} - E(C)$ is bipartite. Clearly, any complete subgraph of $T(\overline{G})$ is also a complete subgraph of \overline{G} . Consequently, \overline{G} has a complete subgraph C such that $\overline{G} - E(C)$ is bipartite, meaning that G is probe 2-clique. \square

For instance, the graph C_6 is not 2-clique and, since $T(\overline{C_6})$ is the disjoint union of two copies of K_3 , C_6 is not probe 2-clique. Notice also that, in Lemma 2 complete subgraph can be replaced by maximal complete subgraph and the lemma is still true. This observation is crucial to obtain a polynomial-time algorithm to recognize the class of probe 2-clique graphs.

Theorem 3 *Let G be a probe 2-clique graph such that \overline{G} contains triangles. Then, $T(\overline{G})$ is a split graph.*

Proof: By Lemma 2, $T(\overline{G})$ has a complete subgraph C such that $\overline{G} - E(C)$ is bipartite. Let S be the subset of vertices of $T(\overline{G})$ not contained in $V(C)$. Suppose $T(\overline{G})$ has an edge e linking two vertices of S . Then e forms a triangle in \overline{G} with some vertex v . However, such a triangle has none of its edges in $E(C)$ and thus $\overline{G} - E(C)$ cannot be bipartite, a contradiction. Therefore, $T(\overline{G})$ is a split graph. \square

Algorithmic aspects: If \overline{G} is triangle-free, then check if for some edge e of \overline{G} , $\overline{G} - e$ is bipartite. Otherwise, find all the triangles of \overline{G} and construct $T(\overline{G})$. If $T(\overline{G})$ is not a split graph, then G is not probe 2-clique. Otherwise, find each maximal complete subgraph C of $T(\overline{G})$ and verify if $\overline{G} - E(C)$ is bipartite. Since a split graph has a linear number of maximal complete subgraphs, all these steps can be clearly performed in $O(n^3)$ time.

3 Partitioned probe diamond-free graphs

Let $G = (P \cup N, E)$ and $H = (P' \cup N', E')$ be two partitioned graphs with stable sets N and N' . The graph H is defined to be a *partitioned subgraph* (a *partitioned induced subgraph*) of G , if H is a subgraph (an induced subgraph) of G , $N' \subseteq N$ and $P' \subseteq P$. When the context is clear, we just say that H is (an induced subgraph) a subgraph of G . We say that G is *isomorphic to H* if and only if there exists a one-to-one function $f : P \cup N \rightarrow P' \cup N'$ preserving adjacency and such that $f(v) \in N'$ for all $v \in N$, and $f(v) \in P'$ for all $v \in P$. We say that the partitioned graph G *does not contain H as induced subgraph* or *does not contain an induced H* if no partitioned induced subgraph of G is isomorphic to H . Given a set of partitioned graphs \mathcal{H} , G is defined to be \mathcal{H} -free if G does not contain an induced H belonging to \mathcal{H} . If \mathcal{H} is a set with a single element H , we use H -free for short. We call *tips of a diamond* the vertices of degree two of a diamond.

In this section we study the structure of probe diamond-free graphs. As a first step towards proving the main theorem of this section, partitioned probe diamond-free graphs are characterized by forbidden partitioned induced subgraphs, through the following theorem. In addition, it gives a strategy to construct a diamond-free graph from a partitioned graph $G = (P \cup N, E)$ by adding those edges whose endpoints are tips of a diamond and belong to N . Notice that the partitioned diamonds H_1, H_2, H_3 of Figure 1 are all the possible partitions of a diamond with at least one of its tips in P . This guarantees that, if $G = (P \cup N, E)$ is a partitioned $\{H_1, H_2, H_3\}$ -free graph, every diamond in G can be eliminated by adding the missing edge (because both tips are in N). The more difficult part of the proof is that of proving that no new diamond is created during this procedure.




Fig. 1: Forbidden partitioned induced subgraphs for probe diamond-free graphs. Black vertices and white vertices represent probe vertices and nonprobe vertices, respectively.

Theorem 4 *Let $G = (P \cup N, E)$ be a partitioned graph. Then, G is a partitioned probe diamond-free graph if and only if G does not contain any partitioned graph depicted in Figure 1 as induced subgraph. Moreover, a probe completion is given by adding the edge set F of non-edges of G whose endpoints belong to N and they both are tips of the same induced diamond of G .*

Proof: It is easy to see that none of the partitioned graphs in Figure 1 is a partitioned probe diamond-free graph. Conversely, let G be a partitioned graph not containing any partitioned induced graph depicted in Fig. 1. Let F be the set of non-edges of G whose endpoints belong to N and they both are tips of the same induced diamond of G . It suffices to prove that the completion $G^* = (N \cup P, E \cup F)$ of G is diamond-free. The proof follows by contradiction and it is split into three cases. Suppose, by the way of contradiction, that G^* is not diamond-free; i.e, there exists a vertex set $\{u, v, x, y\}$ of G^* inducing a diamond $H = G^*[\{u, v, x, y\}]$. Notice that, since G does not contain H_1, H_2 and H_3 as partitioned induced subgraph, G^* does not contain any induced diamond with at most one nonprobe vertex and thus F is well defined.

Case 1: Exactly two vertices of $\{u, v, x, y\}$ are nonprobe vertices. Suppose, without loss of generality, that $u, v \in N$ and $x, y \in P$. By definition of G^* , the tips of H cannot be u and v , so $uv \in F$. By symmetry, we only need to consider two cases: either u and x are the tips of H , or x and y are. First, suppose that u and x are the tips of H . Since $uv \in F$, there exist vertices $w_1, w_2 \in P$ such that u and v are the tips of a diamond induced by $\{u, v, w_1, w_2\}$. In particular, both w_1 and w_2 are different from x . Suppose without loss of generality that w_1 is different from y . If y were adjacent to w_1 , then either $\{w_1, v, x, y\}$ would induce H_3 or $\{w_1, u, x, y\}$ would induce H_2 in G , a contradiction. So, in particular, w_2 is different from y and, by symmetry, nonadjacent to it. Therefore, $\{u, v, w_1, w_2, y\}$ induces H_4 in G , leading to a contradiction. Suppose now that x and y are the tips of H . Again, consider the vertices $w_1, w_2 \in P$ such that u and v are the tips of a diamond induced by $\{u, v, w_1, w_2\}$. Since x and y are nonadjacent, we can suppose without loss of generality that w_1 is different from x and y . Suppose that w_1 is adjacent to either x or y . If w_1 is adjacent to x and nonadjacent to y , then $\{w_1, x, y, u, v\}$ induces H_4 in G , a contradiction. Analogously, if w_1 is adjacent to y and nonadjacent to x , then $\{w_1, x, y, u, v\}$ induces H_4 in G , a contradiction. Consequently, if w_1 were adjacent to either x or y , then w_1 would be adjacent to x and y . So, $\{w_1, x, y, v\}$ would induce H_3 in G , a contradiction. So, in particular, w_2 is different from x and y and, by symmetry, nonadjacent to them. Consequently, $\{w_1, w_2, u, v, x\}$ induces H_4 in G , a contradiction. In what follows we can assume that G^* does not contain any induced diamond with at most two vertices in N .

Case 2: Exactly three vertices of $\{u, v, x, y\}$ are nonprobe vertices.

Suppose first that $u, v, w \in N$ induce a triangle in G^* . We are going to prove it implies that there exists an edge $rs \in E(G)$ whose endpoints belong to P and are complete to $\{u, v, w\}$ (recall that a vertex v is

said to be complete (anticomplete) to a set of vertices A , if v is adjacent (nonadjacent) to all the vertices of A). Suppose, by the way of contradiction, that there is no edge in G whose endpoints belong to P and are complete to $\{u, v, w\}$. Since $uv \in F$, there exist two vertices x_{uv}, y_{uv} belonging to P such that $\{u, v, x_{uv}, y_{uv}\}$ induces a diamond in G . Consequently, w is neither adjacent to x_{uv} nor to y_{uv} , because otherwise, since there is no edge whose both endpoints are adjacent to u, v and w , $\{x_{uv}, y_{uv}, v, w\}$ would induce a diamond in G^* with exactly two nonprobe vertices, a contradiction. Thus, there exist two vertices x_{vw} and y_{vw} , different from x_{uv} and y_{uv} , such that they both are adjacent to v and w and nonadjacent to u . If x_{vw} (resp. y_{vw}) were adjacent to exactly one of $\{x_{uv}, y_{uv}\}$, then $\{x_{uv}, y_{uv}, x_{vw}, v\}$ (resp. $\{x_{uv}, y_{uv}, y_{vw}, v\}$) would induce H_3 in G , a contradiction. Consequently, if x_{vw} (resp. y_{vw}) is adjacent to either x_{uv} or y_{uv} , then it is adjacent to both of them and thus $\{x_{uv}, y_{uv}, x_{vw}, u\}$ (resp. $\{x_{uv}, y_{uv}, y_{vw}, v\}$) induces H_2 in G , a contradiction. So, x_{vw} and y_{vw} are nonadjacent to x_{uv} and y_{uv} . Analogously, there exist two vertices x_{uw} and y_{uw} , both of them adjacent to u and v and different from x_{uv}, x_{vw}, y_{uv} and y_{vw} . Besides, x_{uw} (resp. y_{uw}) is nonadjacent to x_{uv}, x_{vw}, y_{uv} and y_{vw} . Consequently $\{x_{uv}, x_{vw}, x_{uw}, y_{uv}, y_{vw}, y_{uw}\} \cup \{u, v, w\}$ induces H_5 in G , a contradiction. The contradiction arose from supposing that there is no edge in G whose endpoints are complete to $\{u, v, w\}$. So, there exists an edge $rs \in E(G)$ whose endpoints belong to P and are complete to $\{u, v, w\}$. We will prove now that there is no probe vertex z adjacent to u and v and nonadjacent to w . Suppose, by way of contradiction, that there exists a vertex $z \in P$ such that z is adjacent to u and v and nonadjacent to w . If z were nonadjacent to either r or s , then either $\{r, u, v, z\}$ or $\{s, u, v, z\}$ would induce a diamond in G^* with two nonprobe vertices, a contradiction. Therefore, z is adjacent to r and s . So, since z is nonadjacent to w , $\{r, s, w, z\}$ induces H_2 in G , a contradiction. We have already proved that there is no induced diamond in G^* with exactly three nonprobe vertices inducing a triangle. Now, suppose, by the way of contradiction, that $\{u, v, x, y\}$ induces a diamond in G^* with tips y and u , where u, v and y are nonprobe vertices and x is a probe vertex. Since u is adjacent to v in G^* and v is adjacent to y in G^* , then there exists a probe vertex w such that w is adjacent to either u and v , or v and y . Notice that, since G^* does not contain a diamond with at most two nonprobe vertices, w is adjacent to x . Consequently, w is adjacent to y and u , otherwise G^* would contain a diamond with exactly two nonprobe vertices. Since u is nonadjacent to y in G^* , $\{u, w, x, y\}$ induces a diamond with exactly two nonprobe vertices in G^* , a contradiction.

In what follows we can assume that there is no induced diamond in G^* with at most three nonprobe vertices.

Case 3: G^ contains a diamond with four nonprobe vertices.* Finally, we will prove that there is no diamond in G^* with all its edges belonging to F . Suppose, by way of contradiction, that there exist four vertices u, v, w and z belonging to N and inducing a diamond in G^* with tips w and z . Since $\{u, v, w\}$ and $\{u, v, z\}$ induce triangles in G^* , we know that there exist two adjacent vertices x and y belonging to P such that $\{u, v, w\}$ is complete to $\{x, y\}$ and two probe adjacent vertices r and s complete to $\{u, v, z\}$.

Let $e = xy$ and $e' = rs$. Notice that e and e' do not have a common endpoint, because otherwise there would be a vertex in G^* , say x , such that it is adjacent to every vertex in $\{u, v, w, z\}$. Consequently, $\{x, u, w, z\}$ would induce a diamond in G^* with exactly three nonprobe vertices, a contradiction. Besides, the vertices x and y (resp., r and s) are nonadjacent to z (resp., w). Therefore, $\{x, y\}$ and $\{r, s\}$ are anticomplete. Indeed, suppose, by the way of contradiction, they are not anticomplete; and also suppose, without loss of generality, that x is adjacent to r . So, since r is nonadjacent to w , $\{u, r, x, w\}$ induces a diamond in G^* with two nonprobe vertices, contradiction. Consequently, $\{u, v, x, y, r\}$ induces the H_4 in G^* , a contradiction. \square




Fig. 2: Some forbidden induced subgraphs for probe diamond-free graphs.




Fig. 3: Some forbidden induced subgraphs for probe diamond-free graphs.

4 Nonpartitioned probe diamond-free graphs

Let G and F be graphs. We will say that F is a subgraph of G with induced diamonds if F is isomorphic to a subgraph of G (not necessarily induced) in such a way that all the induced diamonds of F induce also diamonds in G , i.e., the tips of a diamond in F are also not adjacent in G .

Before presenting the characterization by forbidden induced subgraphs for probe diamond-free graphs, we will present the following technical lemma whose proof will be postponed up to §4.1.

Lemma 5 *Let G be a $\{F_1, F_2, F_3, F_4, F_6\}$ -free graph. If G contains either S or T_1 as a subgraph with induced diamonds, then G contains one of the graphs depicted in Figure 3 as induced subgraph.*

Theorem 6 *Let G be a graph. G is probe diamond-free if and only if G does not contain any graph depicted in Figures 2 and 3 as induced subgraph. Moreover, a suitable probe partition is obtained by defining N as the set of vertices of G that are tips of an induced diamond in G and P as $V \setminus N$ and a probe completion is given by adding the edge set F of non-edges of G whose endpoints are tips of the same induced diamond.*

Proof: First notice that in the graphs $F_1, F_2, F_4, F_6, S, T_1, \dots, T_{10}$ the set of tips of induced diamonds is not a stable set. In F_3 and F_5 the set of tips of induced diamonds forms a maximal stable set, and thus any probe partition of G having the tips of induced diamonds as nonprobe vertices would contain a partitioned induced subgraph isomorphic to H_4 and H_5 , respectively.

Conversely, let G be a graph not containing any graph depicted in Figures 2 and 3 as induced subgraph. Let N be the set of vertices of G that are tips of an induced diamond in G and $P = V \setminus N$. Let F be a set of non-edges of G whose endpoints are tips of the same induced diamond.

First, we are going to prove that N is a stable set of G . Suppose, by the way of contradiction, that there exist two adjacent vertices u and a belonging to N . Suppose that u belongs to a diamond D_1 induced by the vertices $\{u, v, w, x\}$ whose other tip is x , and a belongs to another diamond D_2 induced by $\{a, b, c, d\}$ whose other tip is d . If $V(D_1)$ does not intersect $V(D_2)$, then $D_1 \cup D_2$ would induce a subgraph in G that contains T_1 as subgraph with induced diamonds D_1 and D_2 . By Lemma 5, since G does not contain F_i for $i = 1, \dots, 4, 6$ as induced subgraph, G contains one of the graphs depicted in Figure 3 as induced subgraph, a contradiction. Hence, we can assume that the diamonds D_1 and D_2 have at least one vertex in common.

If they have exactly one vertex in common, it should have the same number of neighbors in both diamonds, because otherwise $D_1 \cup D_2$ would induce a subgraph in G that contains S as subgraph with induced diamonds D_1 and D_2 . So we are left with two possibilities up to symmetries and considering the fact that u and a are different and adjacent: either $d = x$ or $b = w$.

First, suppose that $d = x$ and $\{u, v, w\} \cap \{a, b, c\} = \emptyset$. Since G is F_6 -free, there exists at least one edge different from au such that one of its endpoints belongs to $\{u, v, w, x\}$ and the other one belongs to $\{a, b, c, d\}$. By symmetry, we can assume that it is either wa or wb . Suppose that w is adjacent to a . Since G is F_2 -free, v is adjacent to a . Consequently, $\{a, d, u, v, w\}$ induces F_1 , a contradiction. Therefore, we can assume that a is nonadjacent to w . By symmetry, we can also assume that v is nonadjacent to a and u is nonadjacent to b and c . Suppose now that w is adjacent to b . Since G is F_2 -free and w is nonadjacent to a , w is adjacent to c . So, $\{a, b, c, d, w\}$ induces F_1 , a contradiction. Therefore, w is nonadjacent to b . Symmetrically, w is nonadjacent to c and v is nonadjacent to b and nonadjacent to c . This leads to a contradiction because G is F_6 -free.

Suppose now that $b = w$ and $\{u, v, x\} \cap \{a, c, d\} = \emptyset$. Since G is F_2 -free, a is adjacent to either v or x . If a were adjacent to v , since G is F_1 -free, a is would be adjacent to x . On the other hand, if a were adjacent to x , since G is F_4 -free, then a would be adjacent to v . Consequently, a is adjacent to v and x . By symmetry, u is adjacent to c and d . Since the vertex set $\{a, b, c, d, x\}$ does not induce F_2 , F_4 nor F_1 , x is adjacent to d and c . Consequently, $\{a, b, u, x, d\}$ induces F_4 , a contradiction. Hence, we can assume that D_1 and D_2 have at least two vertices in common.

These two vertices cannot be nonadjacent because a and u are different and adjacent. So at most one vertex on each diamond is a tip. Suppose that none of them is a tip, and with no loss of generality $b = w$ and $c = v$. Since G is F_1 -free, a is adjacent to x and u is adjacent to d . But then $\{a, b, u, x, d\}$ induces either F_2 or F_4 , a contradiction.

Suppose now that exactly one common vertex is a tip in one of the diamonds. Up to symmetries, we have to consider the cases: $a = v, b = w$ and $d = v, b = w$. In the first case, since the vertex set $\{a, b, c, d, x\}$ does not induce F_2 , F_4 nor F_1 , x is adjacent to d and c . Analogously, u is adjacent to d and c . Consequently, $\{a, b, u, x, d\}$ induces F_4 , a contradiction. In the second case, $\{a, b, u, x, d\}$ induces either F_2 or F_4 , a contradiction.

Finally, suppose that there is one tip of each diamond in the set of common vertices. Up to symmetries, we have to consider the cases: $a = w, b = u$; $a = w, b = x$; $d = w, b = x$ and $d = x, b = w$.

Suppose first that $a = w$ and $b = u$. Since $\{a, b, c, d, v\}$ does not induce F_2 , F_4 nor F_1 , v is adjacent to d and c . But then $\{a, b, v, x, d\}$ induces either F_2 or F_4 , a contradiction. Suppose next that $a = w$ and $b = x$. Again, since $\{a, b, c, d, v\}$ does not induce F_2 , F_4 nor F_1 , v is adjacent to d and c , and now $\{a, b, u, d, v\}$ induces either F_2 or F_4 , a contradiction. Suppose now that $d = w$ and $b = x$. Once more, since $\{a, b, c, d, v\}$ does not induce F_2 , F_4 nor F_1 , v is adjacent to a and c , and so $\{a, b, u, d, v\}$ induces F_4 , a contradiction. Finally, suppose that $d = x$ and $b = w$. Since $\{a, b, c, d, v\}$ does not induce F_1 , F_2

nor F_4 , v is adjacent to a and c . By symmetry, c is adjacent to u . But then $\{a, u, v, c, d\}$ induces F_1 , a contradiction.

Hence, we can assume that D_1 and D_2 have exactly three vertices in common. These vertices cannot induce a 2-edge path, because in that case either $u = a$ or $u = d$, a contradiction with the assumption that a and u are different and adjacent. So, D_1 and D_2 share a triangle, and we have to consider two cases, up to symmetries: either $u = b$ or $u \in V(D_2) \setminus V(D_1)$. In the first case, x should be in $V(D_2) \setminus V(D_1)$, so it is nonadjacent to b and, without loss of generality, it is adjacent to c and d . Therefore, $\{a, b, c, d, x\}$ induces either F_2 or F_4 , depending on the existence of the edge ax , which contradicts our assumptions. In the second case, without loss of generality, either $x = b$ or $x = d$, so u is adjacent to c and to at most one of $\{b, d\}$. In that case, $\{a, b, c, d, u\}$ induces either F_1 , F_2 or F_4 , depending on the existence of one or none of the edges $\{ub, ud\}$, a contradiction again.

Finally, it remains to prove that the completion $G^* = (V, E \cup F)$ is diamond-free. By Theorem 4, it suffices to prove that the partitioned graph $G = (N \cup P, E)$ with probe partition (N, P) does not contain any of the partitioned graphs depicted in Figure 3. By the construction of the partition (N, P) , $G = (N \cup P, E)$ does not contain H_1 , H_2 and H_3 . Finally, since G is $\{F_3, F_5\}$ -free, the partitioned graph $G = (N \cup P, E)$ does not contain the partitioned subgraphs H_4 and H_5 . □

4.1 Proof of Lemma 5

Proof of Lemma 5: In what follows, we mean by “ S (resp. T_1) is a subgraph of G ”, S (resp. T_1) is a subgraph of G with induced diamonds. Let G be a $\{F_1, F_2, F_3, F_4, F_6\}$ -free graph. We will prove the lemma by contradiction. Suppose that G does not contain any graph depicted in Fig. 3 as induced subgraph. We are going to split the proof into two cases.

Case 1: G contains a subgraph H isomorphic to S where the diamonds of S are induced in G . By our assumption, H is not an induced subgraph. Suppose that the vertex set of the subgraph H is labeled by the set $\{a, b, c, d, f, g\}$, where the set $\{a, b, c, d\}$ induces a diamond whose tips are a and d , and $\{b, e, f, g\}$ induces a diamond whose tips are b and g . Since H is not induced, there is at least one edge in $E(G) \setminus E(H)$ whose endpoints belong to $\{a, b, c, d, e, f, g\}$. Since the diamonds are induced in G , the edge is neither ad nor bg .

By the symmetry of the graph, we may assume that at least one of ae , ag , ce and cg does exist. First, suppose that a is adjacent to e . So, a is either adjacent to f or adjacent to g because, otherwise, $\{a, b, e, f, g\}$ would induce F_2 . On the one hand, if a were adjacent to f and nonadjacent to g , $\{a, b, e, f, g\}$ would induce F_4 . On the other hand, if a were adjacent to f and nonadjacent to g , $\{a, b, e, f, g\}$ would induce F_1 . Consequently, since G is $\{F_1, F_4\}$ -free, a is adjacent to f and g . By symmetry, where $\{a, b, c, d\}$ plays the role of $\{b, e, f, g\}$ and e (resp. f) plays the role of a , we conclude that both e and f are adjacent to c and d . Now, by symmetry, where $\{b, e, f, g\}$ is the diamond and d plays the role of a , it follows that d is adjacent to g . If c were nonadjacent to g , $\{b, c, e, f, g\}$ would induce F_1 . Therefore, c is adjacent to g . Consequently, $\{a, b, c, d, g\}$ induces F_4 , a contradiction. This contradiction arose from supposing that a is adjacent to e . So, in what follows, we can assume that a is nonadjacent to e . Symmetrically, we can also assume that a is nonadjacent to f and d is nonadjacent to f and e .

Suppose now that a is adjacent to g . Then, since a is nonadjacent to e and f , $\{a, b, e, f, g\}$ induces F_3 , a contradiction. This contradiction arose from supposing that a is adjacent to g . Therefore, we can assume that a is nonadjacent to g . By symmetry, we can also assume that d is nonadjacent to g .

Now, suppose that c is adjacent to e . Since b is nonadjacent to g , c is adjacent to either f or g . On the one hand, if c were adjacent to g and nonadjacent to f , $\{c, b, e, f, g\}$ would induce F_4 . On the other hand, if c were adjacent to f and nonadjacent to g , $\{c, b, e, f, g\}$ would induce F_1 . Consequently, c is adjacent to f and g . Since b is nonadjacent to g and a is neither adjacent to g nor to f , $\{a, b, c, f, g\}$ induces F_2 . So, we can assume that c is nonadjacent to e . By symmetry, c can be also assumed not to be adjacent to f .

Finally, suppose that c is adjacent to g . Since c is neither adjacent to e nor to f , $\{b, c, e, f, g\}$ induces F_3 , a contradiction.

In what follows, we can assume that G contains no subgraph isomorphic to S with induced diamonds.

Case 2: G contains a subgraph H isomorphic to T_1 where the diamonds of T_1 are induced in G .

By our assumptions, H is not an induced subgraph. Suppose that the vertex set of the subgraph H is labeled by the set $\{a, b, c, d, r, s, t, u\}$, where $\{a, b, c, d\}$ induces a diamond whose tips are a and d , $\{r, s, t, u\}$ induces a diamond whose tips are r and u , and a is adjacent to r . Since H is not induced, there is at least one edge in $E(G) \setminus E(H)$ whose endpoints belong to $\{a, b, c, d, r, s, t, u\}$. Since the diamonds are induced in G , the edge is neither ad nor ru .

By the symmetry of the graph, we may assume that at least one of as , bs , au , bu and du does exist. First, suppose that a is adjacent to s . Since G is F_2 -free and r is nonadjacent to u , a is adjacent to either u or t . On the one hand, if a were adjacent to u and nonadjacent to t , $\{a, r, s, t, u\}$ would induce F_4 . On the other hand, if a were adjacent to t and nonadjacent to u , $\{a, r, s, t, u\}$ would induce F_1 . Therefore, a is adjacent to t and u . Consequently, G contains the graph S as a subgraph with induced diamonds, a contradiction. The contradiction arose from supposing that a is adjacent to s and thus a is nonadjacent to s and symmetrically a is nonadjacent to t . Besides, by symmetry, r is nonadjacent to b and c . Consequently, a is nonadjacent to u , because otherwise $\{a, r, s, t, u\}$ would induce F_3 . Symmetrically, r is nonadjacent to d .

Suppose now that b is adjacent to s . Notice that, if b were adjacent to t , $\{a, b, r, s, t\}$ would induce F_3 . So, b is nonadjacent to t and, by symmetry, s is nonadjacent to c . Therefore, if b were adjacent to u , since b is nonadjacent to t and r , then $\{b, r, s, t, u\}$ would induce F_2 , a contradiction. Consequently, b is nonadjacent to u and, by symmetry, s is nonadjacent to d . Analogously, if c is adjacent to t , then ct is the only edge in $E(G) \setminus E(H)$ having an endpoint in $\{c, t\}$. Thus G contains either an induced T_5 or an induced T_6 , depending on the existence of the edge du . So, we can assume that c is nonadjacent to t . Besides, c is nonadjacent to s because otherwise $\{a, b, c, r, s\}$ would induce F_3 . Suppose now that c is adjacent to u . Since G is $\{F_2, T_8\}$ -free, d is nonadjacent to t and u . So, $\{a, b, c, d, r, s, t, u\}$ induces T_4 , a contradiction. Hence, we can assume that c is nonadjacent to u . Symmetrically, t is nonadjacent to d . Finally, if d were adjacent to u , $\{a, b, c, d, r, s, t, u\}$ would induce T_{10} , a contradiction. Therefore, since G is T_2 -free, from now on, we can assume that b is nonadjacent to s . Symmetrically, we can also assume that b is nonadjacent to t , and c is nonadjacent to s and t .

Suppose that b is adjacent to u . If d were adjacent to u , then u would be adjacent to c . Because, otherwise $\{a, b, c, d, u\}$ would induce F_2 . Consequently, $\{a, b, c, d, u\}$ induces F_1 , a contradiction. So, if b is adjacent to u , then u is nonadjacent to d . Symmetrically, if c is adjacent to u , then d is nonadjacent to u . In addition, by symmetry, if s (resp. t) is adjacent to d , then d is nonadjacent to u . Suppose now that c is adjacent to u . Since G is F_6 -free, either s is adjacent to d or t is adjacent to d . If s (resp. t) were adjacent to d , then $\{b, c, d, u, s\}$ (resp. $\{b, c, d, u, t\}$) would induce F_3 , a contradiction. The contradiction arose from supposing that c is adjacent to u . Therefore, we can assume that if b is adjacent to u , then c is nonadjacent to u . Symmetrically, if s is adjacent to d , then t is nonadjacent to d . Consequently, either exactly one of $\{s, t\}$ is adjacent to d and thus $\{a, b, c, d, r, s, t, u\}$ induces T_9 , or none of them are and

thus $\{a, b, c, d, r, s, t, u\}$ induces T_7 , reaching in both cases a contradiction. Therefore, we can assume that b and c are nonadjacent to u , and s and t are nonadjacent to d .

Finally, since G is T_1 -free, d must be adjacent to u . Consequently, $\{a, b, c, d, r, s, t, u\}$ induces T_3 , a contradiction. This finishes the proof. \square

Next, the characterization for probe diamond-free graphs is used to characterize probe block graphs. Partitioned and nonpartitioned probe block graphs were characterized by forbidden induced subgraphs in [15]. Earlier and independently, probe block graphs were characterized by forbidden induced subgraphs in [11]. We will present a new characterization for this class, which establishes that probe block graphs are exactly the chordal probe diamond-free graphs. Specifically speaking, Theorem 4 and Theorem 6 are generalizations of those characterizations by forbidden induced subgraphs for probe block graphs presented in [15].

Lemma 7 *Let G be a probe block graph. Then, G is chordal.*

Proof: Let $G = (V, E)$ be a probe block graph. Suppose, by the way of contradiction, that G contains an induced cycle H of length at least four. Since the graph induced by $V(H)$ in any probe block completion G^* of G will be connected and will not have a cut-vertex, it should be complete. Since in G every vertex of H has a non-neighbor in H , all the vertices of H have to be nonprobe vertices, a contradiction because they do not form a stable set. \square

Clearly a probe block graph is probe diamond-free and we have already proved that it is also chordal. The following lemma proves that the graph obtained by adding all the edges to a chordal probe diamond-free graph whose endpoints are tips of a diamond remains chordal. Consequently, by Theorem 6, every graph which is chordal and probe diamond-free is probe block.

Lemma 8 *Let $G = (V, E)$ be a connected chordal probe diamond-free graph and F be the set of non-edges of G whose endpoints are tips of some diamond in G . Then, $G^* = (V, E \cup F)$ is chordal.*

Proof: Throughout the proof, index sums should be considered modulo k . Let F be the subset of edges of G^* defined as in the lemma. Suppose, by way of contradiction, that $G^* = (V, E \cup F)$ contains an induced cycle $H = v_1, \dots, v_k, v_1$ for $k \geq 4$ as induced subgraph. Since G is chordal, $v_i v_{i+1} \in F$ for some $i = 1, \dots, k$. Assume that the cycle H contains the minimum number of nonprobe vertices among all the induced cycles contained in G^* . By construction, there exists a vertex $w_1 \in P$ adjacent to v_i and v_{i+1} . By minimality on the number of nonprobe vertices of H and since G^* is diamond-free, w_1 is anticomplete to $V(H) - \{v_i, v_{i+1}\}$ in G^* . If $E(H) \cap F = \{v_i v_{i+1}\}$, then $V(H) \cup \{w_1\}$ would induce a cycle in G , a contradiction. Thus, we can assume that there exists an edge $v_j v_{j+1} \in F$ with $i \neq j$ such that $v_j v_{j+1} \in F$. Therefore, there exists a vertex $w_2 \neq w_1$ belonging to P and adjacent to v_j and v_{j+1} which is also anticomplete to $E(H) - \{v_j v_{j+1}\}$. In addition, by the minimality of the number of nonprobe vertices in H , it follows that $w_1 w_2 \notin E(G)$. Again, if there were no other edges belonging to H in F , G would contain an induced cycle greater than 4, a contradiction. Repeating this procedure, if were necessary, for any edge of H belonging to F , we conclude that G is not chordal, a contradiction again. \square

By combining Theorem 6, Lemma 7 and Lemma 8, it follows the characterization for probe block graphs. This characterization points out the relationship between the class of probe block graphs and

Ptolemaic graphs. Indeed, the corollary below shows that probe block graphs are a subclass of Ptolemaic graphs, and they are exactly those probe diamond-free graphs that are chordal.

Corollary 9 *Let G be a connected graph. The following statements are equivalent:*

1. G is a probe block graph.
2. G is chordal and probe diamond-free.
3. G is Ptolemaic and $\{F_1, S, T_1\}$ -free.

Proof: 1. \Rightarrow 2. Since block graphs are chordal and diamond-free, the class of probe block graphs is contained in the class of probe diamond-free graphs. On the other hand, by Lemma 7, it follows that the class of probe block graphs is contained in the class of chordal graphs.

2. \Rightarrow 1. It is a consequence of Theorem 6 and Lemma 8.

2. \Rightarrow 3. Let G be a probe diamond-free and chordal graph. By Theorem 6, G is, in particular, $\{F_1, S, T_1\}$ -free and F_2 -free. Consequently, since G is chordal, G is $\{F_1, S, T_1\}$ -free and Ptolemaic.

3. \Rightarrow 2. It is a consequence of Theorem 6, since the only chordal graphs in Figures 2 and 3 are F_1 , F_2 , S and T_1 . \square

5 Further remarks

From Theorem 4, Lemma 7 and Lemma 8 it follows that partitioned probe block graphs are exactly the class of partitioned chordal $\{H_1, H_2, H_3\}$ -free graphs (see also [15]). Let \mathcal{G} be a class of graphs. The \mathcal{G} -width of a graph G is the minimum number k of independent sets N_1, \dots, N_k in G such that there exists an embedding of a graph $H \in \mathcal{G}$ on G such that for every edge $e = (x, y)$ of H which is not an edge of G there exists an i with $x, y \in N_i$. In the case that $k = 1$, G is a probe \mathcal{G} -graph. In [5], it is presented a polynomial-time algorithm to recognize whether a given graph has \mathcal{B} -width at most k , being \mathcal{B} the class of block graphs. An interesting open problem is to try to characterize, by forbidden induced subgraphs, those graphs with \mathcal{B} -width at most k , for $k \geq 2$.

References

- [1] D. Bayer, V. B. Le, and H. N. de Ridder. Probe threshold and probe trivially perfect graphs. *Theoretical Computer Science*, 410(47-49):4812–4822, 2009.
- [2] A. Berry, M. C. Golumbic, and M. Lipshteyn. Recognizing chordal probe graphs and cycle-bicolorable graphs. *SIAM Journal on Discrete Mathematics*, 21(3):573–591, 2007.
- [3] D. E. Brown and L. J. Langley. Probe interval orders. In *Brams, Gehrlein, and Roberts, editors, The Mathematics of Preference Choice and Order, Essays in Honor of P. C. Fishburn*, pages 313–322. Springer Berlin Heidelberg, 2009.
- [4] D. B. Chandler, M.-S. Chang, T. Kloks, J. Liu, and S.-L. Peng. Recognition of probe cographs and partitioned probe distance hereditary graphs. In S.-W. Cheng and C. K. Poon, editors, *Proceedings of the Second International Conference on Algorithmic Aspects in Information and Management, AAIM 2006, Hong Kong, China, June 20-22, 2006*, volume 4041 of *Lecture Notes in Computer Science*, pages 267–278. Springer, 2006.

- [5] M.-S. Chang, L.-J. Hung, T. Kloks, and S.-L. Peng. Block-graph width. *Theoretical Computer Science*, 412(23):2496–2502, 2011.
- [6] M.-S. Chang, L.-J. Hung, and P. Rossmanith. Recognition of probe distance-hereditary graphs. *Discrete Applied Mathematics*, 161(3):336–348, 2013.
- [7] J. L. G. J. Chang, A. J. J. Kloks and S.-L. Peng. The pigs full monty - a floor show of minimal separators, in stacs 2005, proceedings. volume 3404 of *Lecture Notes in Computer Science*, pages 521–532. Springer, 2005.
- [8] M. C. Golumbic. *Algorithmic graph theory and perfect graphs*, volume 57 of *Annals of Discrete Mathematics*. Elsevier, Amsterdam, second edition, 2004.
- [9] M. C. Golumbic and M. Lipshteyn. Chordal probe graphs. *Discrete Applied Mathematics*, 143(1-3):221–237, 2004.
- [10] M. C. Golumbic, F. Maffray, and G. Morel. A characterization of chain probe graphs. *Annals of Operations Research*, 188:175–183, 2011.
- [11] L. N. Grippo. *Structural characterizations of intersection graphs*. PhD thesis, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Argentina, 2011.
- [12] P. Hell and J. Huang. Interval bigraphs and circular arc graphs. *Journal of Graph Theory*, 46(4):313–327, 2004.
- [13] J. L. Johnson and J. P. Spinrad, editors. *A polynomial time recognition algorithm for probe interval graphs*, In *Proceedings of the Twelfth Annual ACM-SIAM Symposium on Discrete Algorithms (Washington,DC, 2001)*. SIAM, 2001.
- [14] D. C. Kay and G. Chartrand. A characterization of certain Ptolemaic graphs. *Canadian Journal of Mathematics*, 17:342–346, 1965.
- [15] V. Le and S.-L. Peng. Characterizing and recognizing probe block graphs. In R.-S. Chang, L. C. Jain, and S.-L. Peng, editors, *Proceedings of the Workshop on Algorithms, Bioinformatics, and Computation Theory of the International Computer Symposium, ICS 2012, Hualien, Taiwan, December 12-14, 2012*, volume 20 of *Smart Innovation, Systems and Technologies*, pages 7–13. Springer Berlin Heidelberg, 2013.
- [16] R. M. McConnell and J. Spinrad. Construction of probe interval models. In *Proceedings of the Thirteenth Annual ACM-SIAM Symposium on Discrete Algorithms, January 6-8, 2002, San Francisco, CA, USA.*, pages 866–875, 2002.
- [17] J. Spinrad. Circular-arc graphs with clique cover number two. *J. Comb. Theory, Ser. B*, 44(3):300–306, 1988.
- [18] D. B. West. *Introduction to graph theory*. Prentice Hall Inc., Upper Saddle River, NJ, 1996.
- [19] P. Zhang, E. Schon, S. Fischer, E. C. amd J. Weiss, S. Kistler, and P. Bourne. An algorithm based on graph theory for ghe assembly of contigs in physical mapping of DNA. *Computer Applications in the Biosciences*, 10(3):309–317, 1994.

