

Sequential selection of the k best out of n rankable objects

F. Thomas Bruss, Guy Louchard

► **To cite this version:**

F. Thomas Bruss, Guy Louchard. Sequential selection of the k best out of n rankable objects. Discrete Mathematics and Theoretical Computer Science, DMTCS, 2016, Vol. 18, no 3. hal-01208327v3

HAL Id: hal-01208327

<https://hal.archives-ouvertes.fr/hal-01208327v3>

Submitted on 23 Sep 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Sequential selection of the κ best out of n rankable objects

F. Thomas Bruss¹

Guy Louchard²

¹ *Université Libre de Bruxelles, Département de Mathématique, CP 210 Boulevard du Triomphe, B-1050 Bruxelles, Belgium, tbruss@ulb.ac.be*

² *Université Libre de Bruxelles, Département d'Informatique, CP 212, Boulevard du Triomphe, B-1050 Bruxelles, Belgium, louchard@ulb.ac.be*

received 5th Oct. 2015, revised 12th July 2016, accepted 22nd Aug. 2016.

The objective of this paper is to find in a setting of n sequential observations of objects a good online policy to select the κ best of these n uniquely rankable objects. This focus is motivated by the fact that it is hard to find closed form solutions of optimal strategies for general κ and n . Selection is without recall, and the idea is to investigate threshold functions which maintain all present information, that is thresholds which are functions of all selections made so far. Our main interest lies in the asymptotic behaviour of these thresholds as $n \rightarrow \infty$ and in the corresponding asymptotic performance of the threshold algorithm.

Keywords: secretary problem, odds-theorem, multiple stopping, monotonicity, unimodality, relative ranks, record-values, optimality

1 Introduction

The so-called secretary problem is a classical problem of optimal stopping on a sequence of rankable observations. It is the problem of choosing the best of n rankable options appearing in random arrival order with probability $1/n!$ each. It was first solved by Lindley [8] in 1961. See e.g. Ferguson [7] for an interesting review. This is the special case $\kappa = 1$ of the problem we consider here, namely to find with one stop without recall the best of n uniquely rankable objects or options. Arguably the shortest solution to this problem is provided by the more general odds-theorem of optimal stopping (Bruss [2]). The optimal strategy is to wait until the threshold index $s \in N$ and to accept the first relative rank 1 from s onwards (if any). Here s is defined as the largest $m \in N$ such that $\sum_{j=m}^n r_j \geq 1$ where r_j denotes the odds of the j th event being a candidate for stopping. If no such m exists, then $s := 1$ by definition.

For the classical secretary problem $r_j = (1/j)/(1 - 1/j) = 1/(j - 1)$ because the best candidate among the first j is the j th in chronological order with probability $1/j$. The corresponding win probability is then

$$V_n = \frac{s-1}{n} \sum_{j=s-1}^{n-1} \frac{1}{j}.$$

Putting $s := s(n)$ it is well-known that $\lim_{n \rightarrow \infty} s_n/n = 1/e$. V_n has the same limit as the function s_n/n , namely $1/e$. Interestingly, (Bruss [3]), the value $1/e$ is also the precise lower bound in much more general problems of stopping with maximum probability on *any* last specific event. For further generalizations for the case $\kappa = 1$ see the review by Dendievel [6].

κ best choices

The problem of making sequentially the κ best choices, or in short the κ -best problem, is to find an efficient stopping rule to select (all) the κ best objects online from a sequence of n objects. The study of this problem is the main objective of the present paper. Indeed, if $\kappa > 1$ then the optimal solution as well as the limiting behaviour of the thresholds seem difficult. Actually, one would not expect such a difficulty because one might expect a tractable recursive structure. Some recursive structure is there, of course, but a little reflection shows that once one has selected $1 \leq \kappa' < \kappa$ objects, the optimal continuation does not only depend $\kappa - \kappa'$ but also on the values of the chosen ones. This difficulty motivated several authors before, and the present paper complements earlier results by Vanderbei [13] and Tamaki [12] by an independent approach rather than really solving it. However, as we shall argue, this complement has interest on its own.

The problem is mathematically interesting. We agree that there are probably not many situations in everyday life where our model would directly fit (except for $\kappa = 1$ or $\kappa = 2$). However, we know of several people who have tried to get answers and found the problem untractable. We felt that one should not give up and try to give at least an (arguably) good answer.

Related work

Platen [9] studies the problem of choosing a fixed number of objects (secretaries) from a sequence of n but his objective function is different. He thinks of the rank k as generating the cost k and considers the problem of minimizing the sum of the cost of all selections. Notice that this is different from the problem of minimizing the expected rank as studied by Chow et al [5] and also again quite different from our problem of maximizing the expected payoff of one if and only if all the κ best objects are chosen, and zero otherwise. Platen also considers more general payoff (cost) functions, but, motivated by thinking of secretary problems, limits his interest to additive functions, which do not apply to our problem.

A more weakly related problem has been studied by Rose [10] for the case $\kappa = 2$. Rose introduces in the same model two decision makers, one of which is defined as dominant. The objective is to select the two best objects, the best one being assigned to the dominant decision maker. Rose considers a strategy as successful if and only if both decision makers reach their goal under the online assignment constraint. Hence here the optimal success probability should be (and is, as we shall see) smaller than in our problem without assignments. Note also that the concurrent assignment constraint gives Roses' problem quite a different character.

Another κ -choice problem was studied by Ano et al. [1] who maximise the probability of finding the *best* (only) candidate with κ available sequential choices. Szajowski and Lebek [11] solve an interesting related investment problem for $\kappa = 2$.

Wilson [15] generalizes the model of Rose for general κ , also with simultaneous online assignment of ranks. The paper displays important optimality conditions for general κ , but closed-form solutions seem now to become hard.

Specifically related work of Tamaki and Vanderbei

Tamaki [12] looks at the same problem for $\kappa = 3$ and shows that, interestingly, in contrast to the case $\kappa = 1$ (secretary problem) and the case $\kappa = 2$ the optimal strategy is now no longer simple. Here *simple* means that, for the κ selections, the strategy waits until a threshold index s_k and selects then consecutively the next k (if possible) record candidates. Tamaki's observation of the non-simplicity of the optimal rule for $\kappa = 3$ (and $\kappa \geq 3$) is important in this context because it gives additional motivation to the work of the approach presented in this paper as well as to earlier work of Vanderbei [13] described below.

Vanderbei [13] studies the corresponding problem without online assignments with an original approach by formulating the problem as a control problem defined on a Markov chain. An explicit solution is found for n even and $\kappa = n/2$. The corresponding optimal win probability equals $\frac{1}{n/2+1}$. For general κ , this article confines its interest to studying a simple sub-optimal solution. Thus Vanderbei studies the same problem as ours but the approach is different. As far as we understand, Vanderbei's sufficient condition (see the last line of [13] p.481) is

not a sufficient condition in the set of strategies which we consider. Indeed, we need in general also information about 2-records, 3-records, up to $(\kappa - 1)$ -records, where a s -record is a candidate which is at the time of its arrival of relative rank s . Having said this we see by Theorem 5.1 of [13] that the sufficient condition we require is implicit by the definition of “marginal”, and the definition of the sets T and S' , so that our strategies may coincide. A closer comparison between approaches is difficult, however, since Vanderbei’s approach leads to a difference-differential equation (see (5.3) in [13]) whereas our approach is purely algorithmic. Without being able to say more in definite terms, we think of our contribution as of being an attractive alternative to Vanderbei’s result and as a way of access to a *good* solution for *general* κ .

Let us finally mention another interesting result by Vanderbei [14]. In this paper, he considers a minor variant of this classical secretary problem. He wishes to pick not the best but the second best postdoc (the best is going to Harvard). In this case, an explicit solution can be given both for the optimal strategy and the associated optimal success probability. The probability of success is $k_0(n - k_0)/(n(n - 1))$, where $k_0 = \lfloor n/2 \rfloor$. Clearly, as n goes to infinity, the probability of success tends to $1/4$. Hence, it is easier to pick the best than the second best, a phenomenon which is not only true in several other selection problems but, often enough, felt to be true in real life.

2 Algorithm

The approach to the algorithm we will propose consists of two parts: firstly, defining a suitable strategy, secondly, computing sequentially the necessary parameters we have to plug in. We are interested in understanding the asymptotic form of an optimal or, at least, efficient, strategy, and its corresponding value as $n \rightarrow \infty$.

The κ choices threshold strategy:

Our strategy, based on thresholds, is defined as follows:

we use κ thresholds: $1 \leq j_1 < j_2 < \dots < j_\kappa \leq n$. The way we compute these thresholds will be characterized later on. It will turn out that there are no general closed forms for the optimum values for these thresholds.

Following Vanderbei’s terminology, we define first two kinds of candidates:

1. *compulsory candidates* i.e. candidates whom we *must* retain, given the relative ranks of the candidates we have already chosen. For instance, if we have chosen so far ℓ candidates, with relative ranks $1 < 2 < \dots < \ell$ (relative rank 1 corresponds to the best candidate seen so far, rank 2 to the second best seen so far, etc.), we *must* retain any candidate with relative rank $\leq \ell$.
2. *marginal candidates* i.e. candidates whose relative rank is equal to $\ell + 1$.

According to our strategy, we first choose the first record (if any) arriving at position $u_1 \geq j_1$. This is done for all possible values of j_1 where j_1 is some index between 1 and n . We then compute for each possible position u_1 an optimal threshold $j_2(u_1) \geq u_1$. The reason behind this is that we have two possibilities, namely:

1. We observe a *compulsory candidate* at position $u_2 < j_2$. We take this candidate and we start again the strategy at position u_2 , with a new threshold $j_3(u_2)$. The strategy iterates then this procedure accordingly over the position values. At this stage it is difficult to say already more, and the details are postponed to the beginning of Sec.4.
2. We do not observe any *compulsory candidate* before j_2 . Then, from j_2 on, we retain a *compulsory candidate* or a *marginal candidate* (if any) at position u_2 . We now start again the strategy at position u_2 , with a new threshold $j_3(u_2)$ and, as before, we iterate the procedure.

Examples

To give an example, we first define the 2-choice strategy. We define a (j_1, j_2) -strategy as the policy to act as follows:

1. wait until index j_1 without any action;

2. select, from j_1 onwards, the first record (if any). Thereafter instructions will split according to two possibilities:
3. (a) if we select only one record value up to a certain index $j_2 \in (u_1, n)$ then we accept a record value or 2-record value after j_2 ,
or else,
(b) we select two consecutive records up to j_2 (if any.)

For $\kappa = 3$, the definition of a (j_1, j_2, j_3) -strategy if $\kappa = 3$ is in principle analogous but, again, it is better to postpone the details to the beginning of Sec.4. The corresponding optimal thresholds are denoted by j_1^*, j_2^*, j_3^* . The order of computation is backwards.

The asymptotic results for $\kappa = 2$ and $\kappa = 3$ are summarized in the following theorem:

Theorem 2.1 • With $j_1^* = j_1^*(n)$, $j_2^* = j_2^*(n)$, the optimal thresholds j_1^* and j_2^* for the case $\kappa = 2$ satisfy the asymptotic relationship

$$\begin{aligned} j_1^*/n \rightarrow \chi_1^* &= -e^{-1/2}W(-\exp(-3 + e^{1/2})) = 0.2291147286\dots \\ &\text{where } W(x) \text{ is the Lambert's function,} \\ j_2^*/n \rightarrow \chi_2^* &= e^{-1/2} = 0.6065306596\dots, \end{aligned}$$

The asymptotic success probability of the (j_1^*, j_2^*) -strategy equals 0.2254366561\dots

- In the case $\kappa = 3$, the corresponding asymptotic relationships are

$$\begin{aligned} j_1^*/n \rightarrow \chi_1^* &= .1666171752\dots, \\ j_2^*/n \rightarrow \chi_2^* &= -e^{-1/3}W(-\exp(-5/2 + e^{1/3})) = .4369818602\dots, \\ j_3^*/n \rightarrow \chi_3^* &= e^{-1/3} = .7165313106\dots, \end{aligned}$$

and the asymptotic success probability of the (j_1^*, j_2^*, j_3^*) -strategy equals .1625200069\dots

Our computational technique can be extended to any κ . We will illustrate the optimal thresholds computation and analyze the performance, i.e. the success probability of our algorithm for the cases $\kappa = 2, \kappa = 3$.

3 The case $\kappa = 2$

Let us first compute $j_2(u_1) \geq u_1$, given u_1 .

As explained in Sec.1, we have two possibilities: $u_1 < j_2 \leq u_2$, or, alternatively, $u_1 < u_2 < j_2$. The probability of success of the first case is given by

$$\begin{aligned} &\sum_{u_2=j_2}^n \prod_{k=u_1+1}^{j_2-1} \left(1 - \frac{1}{k}\right) \prod_{\ell=j_2}^{u_2-1} \left(1 - \frac{2}{\ell}\right) \frac{2}{u_2} \prod_{s=u_2+1}^n \left(1 - \frac{2}{s}\right) \\ &= \sum_{u_2=j_2}^n \frac{u_1}{n} \frac{2}{u_2-2} \frac{j_2-2}{n-1} =: u_1 F_1(j_2) \text{ say.} \end{aligned}$$

To see this note that after index j_2 , we have two possible types of *candidates*, a *compulsory* or a *marginal* candidate at position u_2 . After u_2 , we must exclude any *compulsory candidate*.

We note that $u_1 F_1(j_2)$ is unimodal in j_2 because $u_1 F_1(j_2) - u_1 F_1(j_2 + 1)$ yields after a few steps of simplification the form

$$\frac{1}{n(n-1)} 2u_1 \left(1 - \sum_{u_2=j_2+1}^n \frac{1}{u_2-2} \right)$$

which changes sign at most once according to the value j_2 .

The probability of success of the second case is given by

$$\begin{aligned} & \sum_{u_2=u_1+1}^{j_2-1} \prod_{k=u_1+1}^{u_2-1} \left(1 - \frac{1}{k} \right) \frac{1}{u_2} \prod_{\ell=u_2+1}^n \left(1 - \frac{2}{\ell} \right) \\ &= \sum_{u_2=u_1+1}^{j_2-1} \frac{u_1}{n(n-1)} =: u_1 F_2(u_1, j_2), \\ & \text{with } F_2(u_1, j_2) := \frac{j_2 - u_1 - 1}{n(n-1)}. \end{aligned}$$

Here we note that after u_1 and before j_2 , we have one possible *compulsory candidate* at position u_2 . After u_2 , we must exclude any *compulsory candidates*. $u_1 F_2(u_1, j_2)$ is clearly monotone in j_2 because it is, as seen in the last sum term, affine linear in j_2 . Consequently $u_1 F_1(j_2) + u_1 F_2(u_1, j_2)$ is unimodal in j_2 because the sum of a unimodal function and an affine linear function is unimodal.

Of course, $F_2(u_1, j_2)$ is quite simple here, but, viewing general κ , we prefer to keep the general notation.

As $n \rightarrow \infty$, we use the Euler Maclaurin formula to replace sums by integrals (see also the continuous-time model Tamaki [12]). Moreover, we use continuous variables $u'_1 := u_1/n, u'_2 := u_2/n, \dots$. By a slight abuse of notations, in order to simplify our expressions, we will continue to use u_1, u_2, \dots as continuous time variables. Also, to avoid any confusion, we will use the notation χ_k for the (continuous) asymptotic of j_k/n .

Since both parts contributing to the win probability have the same factor u_1 , it now suffices to maximize the sum $F_1(j_2) + F_2(u_1, j_2)$, conditioning on u_1 , that is we must maximize the asymptotic expression

$$V_1(u_1, \chi_2) := \int_{\chi_2}^1 2 \frac{\chi_2}{u_2} du_2 + \int_{u_1}^{\chi_2} 1 du_2 = -u_1 + \chi_2 - 2\chi_2 \ln(\chi_2). \quad (1)$$

This gives $\chi_2^* = e^{-1/2} = 0.6065306596 \dots$

We note that $V_1(u_1, \chi_2)$ can be seen as the sum of two functions $\varphi_1(u_1)$ and $\varphi_2(\chi_2)$ namely $\varphi_1(u_1) + \varphi_2(\chi_2)$. Since $\frac{d\varphi_2(\chi_2)}{d\chi_2} = 1 - 2 \ln(\chi_2) - 2$ is monotone in χ_2 , the function φ_2 is unimodal. From now on, the unimodality will sometimes be derived in the continuous model. A plot of $\varphi_2(\chi_2)$ is given in Figure 1.

We must now compute j_1 . But we note that the first record can occur *after* j_2^* . It is clear from the unimodality of $\varphi(\chi_2)$, that, asymptotically, $j_2(u_1)$ is then exactly given by u_1 . Actually

- $j_2(u_1) = j_2^*$ if $u_1 < j_2^*$,
- $j_2(u_1) = u_1$ if $u_1 \geq j_2^*$.

So we obtain the following success probability $P(j_1) = P_1(j_1) + P_2(j_1)$, with, on the one hand

$$\begin{aligned} u_1 < j_2^* : P_1(j_1) &= \sum_{u_1=j_1}^{j_2^*-1} \prod_{i=j_1}^{u_1-1} \left(1 - \frac{1}{i} \right) \frac{1}{u_1} [F_1(j_2^*) + F_2(u_1, j_2^*)] \\ &= \sum_{u_1=j_1}^{j_2^*-1} \frac{j_1 - 1}{u_1 - 1} [F_1(j_2^*) + F_2(u_1, j_2^*)], \end{aligned} \quad (2)$$




Fig. 1: $\varphi_2(\chi_2)$

and on the the other hand

$$\begin{aligned}
 j_2^* \leq u_1 < u_2 : P_2(j_1) &= \sum_{u_1=j_2^*}^{n-1} \sum_{u_2=u_1+1}^n \prod_{i=j_1}^{u_1-1} \left(1 - \frac{1}{i}\right) \frac{1}{u_1} \prod_{\ell=u_1+1}^{u_2-1} \left(1 - \frac{2}{\ell}\right) \frac{2}{u_2} \prod_{s=u_2+1}^n \left(1 - \frac{2}{s}\right) \\
 &= \sum_{u_1=j_2^*}^{n-1} \sum_{u_2=u_1+1}^n \frac{2}{u_2-2} \frac{j_1-1}{n} \frac{1}{n-1}. \tag{3}
 \end{aligned}$$

Note that (3) is again affine linear in j_1 . Therefore, if $P_1(j_1)$ defined in (2) is unimodal in j_j then $P(j_1) = P_1(j_1) + P_2(j_1)$ is unimodal in j_1 because the sum of an affine linear function and a unimodal function is unimodal. However, since j_1 figures in (2) both in the numerator and as the starting point of summation we cannot decide upon unimodality of $P_1(j_1)$ without being able to quantify the summands $[F_1(j_2^*) + F_2(u_1, j_2^*)]$.

However, we cannot do this, that is, here the problem becomes circular. Motivated by staying in line with our determination to keep things tractable (see also below) and to focus our interest on the asymptotic case as $n \rightarrow \infty$, we disregard in this place the problem of unimodality for all n . (Here it is good to know that unimodality, which is a sufficient condition for our approach to be solid, need not be necessary!)

The asymptotic total success probability is, independently of the problem of unimodality, always of interest and readily accessible, namely

$$\begin{aligned}
 V_2(\chi_1) &= \chi_1 \int_{\chi_1}^{\chi_2^*} \frac{1}{u_1} V_1(u_1, \chi_2^*) du_1 + \chi_1 \int_{u_1=\chi_2^*}^1 \int_{u_2=u_1}^1 \frac{2}{u_2} du_2 du_1 \\
 &= \chi_1 \left[\chi_1 - 2 \ln(\chi_1) e^{-1/2} - 5e^{-1/2} + 2 \right].
 \end{aligned}$$

Maximizing $V_2(\chi_1)$ then yields the asymptotically optimal threshold

$$\chi_1^* = -e^{-1/2} W(-\exp(-3 + e^{1/2})) = 0.2291147286 \dots$$

where $W(x)$ is the Lambert function, that is the solution of

$$W(x)e^{W(x)} = x.$$

We use its principal branch which is analytic at 0. This leads from the above to $V_2(\chi_1^*) = 0.2254366561\dots$ and the value χ_1^* is unique since $V_2(\chi_1)$ is unimodal in χ_1 . A plot of $V_2(\chi_1)$ is given in Figure 2.




Fig. 2: $V_2(\chi_1)$

We have compared, for $n = 500$, $V_2(\chi_1)$ with $P(j_1)$. This is shown in Figure 3 (line= $V_2(\chi_1)$, box= $P(j_1)$). This is quite satisfactory. The relative error at j_1^* is given by $\frac{V_2(\chi_1^*)}{P(j_1^*)} - 1 = -0.0033478319\dots$

Remark on the circular part of the problem

How good is the approximation for finite n ? Here the problem is the same as above; the mentioned circular part problem persists. We do not know, and we cannot do better because we would have to know the optimal strategy and its precise value.

However, there are good reasons to be optimistic. The few comparisons we could do with simplified problems (comparisons with upper bounds or lower bound stemming from problems for which we know the optimal strategy and its value) suggest that the resulting values should not be far off the optimum. Moreover, the asymptotic behaviour (see Fig. 3) of the values seems very nice.

Similarly, we cannot guarantee that the use of the asymptotic j_2^* is mathematically rigorous. What we can say is that we do not see what may go wrong in doing so, and, in the worst case, our results are at least lower bounds for the optimal success probability.

4 The case $\kappa = 3$

The strategy for general κ proceeds as follows. Assume that we are at time $u \in [j_k^*, j_{k+1}^*)$,

- if we have already chosen $k - 1$ candidates or less, we select a *compulsory* or *marginal candidate* (if any) in $[u, j_{k+1}^*)$;
- if we have already chosen more than $k - 1$ candidates, we select a *compulsory candidate* (if any) in $[u, j_{k+1}^*)$.




Fig. 3: Comparison, for $n = 500$ of $V_2(\chi_1)$ (line) with $P(j_1)$ (box).

The initial condition is the following: for $u \in [j_1^*, j_2^*)$, the *marginal candidate* is the first record, by definition.

Let us now consider the case $\kappa = 3$. We must first compute $j_3(u_2) \geq u_2$, given u_2 (at this time, we have selected the second candidate). Again, we have two possibilities: $u_2 < j_3 \leq u_3$, or, alternatively, $u_2 < u_3 < j_3$.

The probability of success of the first case is given by

$$\begin{aligned} & \sum_{u_3=j_3}^n \prod_{s=u_2+1}^{j_3-1} \left(1 - \frac{2}{s}\right) \prod_{t=j_3}^{u_3-1} \left(1 - \frac{3}{t}\right) \frac{3}{u_3} \prod_{v=u_3+1}^n \left(1 - \frac{3}{v}\right) \\ &= \sum_{u_3=j_3}^n \frac{3}{u_3-3} \prod_{s=u_2+1}^n \left(1 - \frac{2}{s}\right) \frac{j_3-3}{n-2} =: \prod_{s=u_2+1}^n \left(1 - \frac{2}{s}\right) F_3(j_3) \text{ say,} \end{aligned}$$

Note that this expression is unimodal in j_3 if and only if

$$g(j_3) := \frac{3}{n-2} \sum_{u_3=j_3}^n \frac{1}{u_3-3} (j_3-3)$$

is unimodal in j_3 . This is true if

$$h(j) := \left(\sum_{u=j}^n \frac{j-3}{u-3} \right) - \left(\sum_{u=j+1}^n \frac{(j-3)+1}{u-3} \right)$$

changes, as a function of j , at most once its sign. An easy calculation shows that

$$h(j) = 1 - \sum_{u=j+1}^n \frac{1}{u-3}$$

so that $h(j)$ is monotone increasing in j .

The probability of success of the second case is given by

$$\begin{aligned} & \sum_{u_3=u_2+1}^{j_3-1} \prod_{s=u_2+1}^{u_3-1} \left(1 - \frac{2}{s}\right) \frac{2}{u_3} \prod_{t=u_3+1}^n \left(1 - \frac{3}{t}\right) \\ &= \sum_{u_3=u_2+1}^{j_3-1} \frac{2}{n-2} \prod_{s=u_2+1}^n \left(1 - \frac{2}{s}\right) =: \prod_{s=u_2+1}^n \left(1 - \frac{2}{s}\right) F_4(u_2, j_3) \text{ say,} \end{aligned}$$

which is clearly monotone in j_3 . Asymptotically, from $F_3(j_3) + F_4(u_2, j_3)$, conditioning on u_2 , we must maximize, with respect to the index χ_3 , the function

$$V_3(u_2, \chi_3) := \int_{\chi_3}^1 3 \frac{\chi_3}{u_3} du_3 + \int_{u_2}^{\chi_3} 2 du_3 = -2u_2 + 2\chi_3 - 3\chi_3 \ln(\chi_3).$$

Note the similarity with V_1 given in equation (1). This gives $\chi_3^* = e^{-1/3} = .7165313106\dots$. Again we note that $V_3(u_2, \chi_3)$ has the form $\varphi_3(u_2) + \varphi_4(\chi_3)$, and that $\varphi_4(\chi_3)$ is unimodal in χ_3 .

We must now compute $j_2(u_1) \geq u_1$ given u_1 . But we note that the relevant position u_2 can occur *after* j_3^* . It is then clear, from the unimodality of $\varphi_4(\chi_3)$, that, asymptotically, $j_3(u_2)$ is then exactly given by u_2 . Actually,

- $j_3(u_2) = j_3^*, u_2 < j_3^*$,
- $j_3(u_2) = u_2, u_2 \geq j_3^*$.

We have now to consider the following cases:

- $u_1 < j_2 \leq u_2 < j_3^*$,
- $u_1 < j_2 < j_3^* < u_2 < u_3$,
- $u_1 < u_2 \leq j_2 < j_3^*$.

This leads respectively to

$$\begin{aligned} P_3(u_1, j_2) &= \sum_{u_2=j_2}^{j_3^*-1} \prod_{k=u_1+1}^{j_2-1} \left(1 - \frac{1}{k}\right) \prod_{\ell=j_2}^{u_2-1} \left(1 - \frac{2}{\ell}\right) \frac{2}{u_2} \prod_{s=u_2+1}^n \left(1 - \frac{2}{s}\right) [F_3(j_3^*) + F_4(u_2, j_3^*)] \\ &= \sum_{u_2=j_2}^{j_3^*-1} \frac{2}{u_2-2} \frac{u_1}{n} \frac{j_2-2}{n-1} [F_3(j_3^*) + F_4(u_2, j_3^*)], \end{aligned}$$

Note the similarity with equation (2),

$$\begin{aligned} P_4(u_1, j_2) &= \sum_{u_2=j_3^*}^{n-1} \sum_{u_3=u_2+1}^n \prod_{k=u_1+1}^{j_2-1} \left(1 - \frac{1}{k}\right) \prod_{\ell=j_2}^{u_2-1} \left(1 - \frac{2}{\ell}\right) \frac{2}{u_2} \prod_{s=u_2+1}^{u_3-1} \left(1 - \frac{3}{s}\right) \frac{3}{u_3} \prod_{t=u_3+1}^n \left(1 - \frac{3}{t}\right) \\ &= \sum_{u_2=j_3^*}^{n-1} \sum_{u_3=u_2+1}^n \frac{6}{u_3-3} \frac{u_1}{n} \frac{j_2-2}{(n-1)(n-2)}, \end{aligned}$$

Note the similarity with equation (3),

$$P_5(u_1, j_2) = \sum_{u_2=u_1+1}^{j_2-1} \prod_{k=u_1+1}^{u_2-1} \left(1 - \frac{1}{k}\right) \frac{1}{u_2} \prod_{s=u_2+1}^n \left(1 - \frac{2}{s}\right) [F_3(j_3^*) + F_4(u_2, j_3^*)]$$

$$= \sum_{u_2=u_1+1}^{j_2^*-1} \frac{u_1}{n(n-1)} [F_3(j_3^*) + F_4(u_2, j_3^*)].$$

Asymptotically, conditioning on u_1 , we must then maximize

$$\begin{aligned} V_4(u_1, \chi_2) &= \int_{\chi_2}^{\chi_3^*} 2 \frac{\chi_2^2}{u_2} V_3(u_2, \chi_3^*) du_2 + \int_{u_2=\chi_3^*}^1 \int_{u_3=u_2}^1 6 \frac{\chi_2^2}{u_3} du_3 du_2 + \int_{u_1}^{\chi_2} V_3(u_2, \chi_3^*) du_2 \\ &= -3e^{-1/3} u_1 + u_1^2 + 3\chi_2^2 + 6\chi_2 - 6e^{-1/3} \chi_2 \ln(\chi_2) - 11\chi_2 e^{-1/3}. \end{aligned}$$

This gives $\chi_2^* = -e^{-1/3} W(-\exp(-5/2 + e^{1/3})) = .4369818602\dots$ Again we note that $V_4(u_1, \chi_2)$ has the form $\varphi_5(u_1) + \varphi_6(\chi_2)$ and $\varphi_6(\chi_2)$ is unimodal in χ_2 .

We now compute j_1 . We note that u_1 can occur *after* j_2^* . It is then clear, from the unimodality of $\varphi_6(\chi_2)$, that, asymptotically, $j_2(u_1)$ is then exactly given by u_1 . Actually now the split in the index j_2^* is given by

- $j_2(u_1) = j_2^*, u_1 < j_2^*$,
- $j_2(u_1) = u_1, u_1 \geq j_2^*$.

We obviously need a concise notation for all cases we will consider (this will of course be applied to cases $\kappa \geq 4$). The idea is to create a bijection between the different cases and an urn model. Indeed, the adequate model here is the Bose-Einstein urn model, that is balls correspond to chosen candidates u_1, u_2, \dots and urns to intervals $[j_k, j_{k+1} - 1]$. We throw κ indistinguishable balls into κ ordered urns labelled $1, 2, \dots, \kappa$. The number of possible cases is given by $\binom{2\kappa-1}{\kappa}$ for example 10 cases for $\kappa = 3$. We next label the balls in increasing order. Denote then by $(\alpha_1, \alpha_2, \alpha_3)$ the urn labels of balls $(1, 2, 3)$. For instance, the case $(1, 3, 3)$ corresponds to $j_1 \leq u_1 < j_2^* < j_3^* \leq u_2 < u_3$ because it means that the arrival time u_1 falls in the first urn defined by the borders j_1 and $j_2^* - 1$ and the second and third candidate fall in the third urn with bounds j_3^* and n . The corresponding probability is given by

$$\begin{aligned} & \sum_{u_1=j_1}^{j_2^*-1} \sum_{u_2=j_3^*}^{n-1} \sum_{u_3=u_2+1}^n \prod_{i=j_1}^{u_1-1} \left(1 - \frac{1}{i}\right) \frac{1}{u_1} \prod_{k=u_1+1}^{j_2^*-1} \left(1 - \frac{1}{k}\right) \prod_{\ell=j_3^*}^{u_2-1} \left(1 - \frac{2}{\ell}\right) \frac{2}{u_2} \times \\ & \times \prod_{s=u_2+1}^{u_3-1} \left(1 - \frac{3}{s}\right) \frac{3}{u_3} \prod_{t=u_3+1}^n \left(1 - \frac{3}{t}\right). \end{aligned}$$

We will not detail the tedious, but routine computations corresponding to our 10 cases. We provide only the final asymptotic integrals.

For this we must maximize the success probability $V_5(\chi_1) = \chi_1[K_1 + K_2 + K_3 + K_4 + K_5]$ with

cases $(1, 1, 1), (1, 1, 2), (1, 1, 3), (1, 2, 2), (1, 2, 3), (1, 3, 3)$:

$$K_1 = \int_{\chi_1}^{\chi_2^*} \frac{1}{u_1} V_4(u_1, \chi_2^*) du_1,$$

case $(2, 2, 3)$:

$$K_2 = \int_{u_1=\chi_2^*}^{\chi_3^*} \int_{u_2=u_1}^{\chi_3^*} \int_{u_3=\chi_3^*}^1 6 \frac{\chi_3^*}{u_2 u_3} du_3 du_2 du_1,$$

case $(2, 3, 3)$:

$$K_3 = \int_{u_1=\chi_2^*}^{\chi_3^*} \int_{u_2=\chi_3^*}^1 \int_{u_3=u_2}^1 6 \frac{1}{u_3} du_3 du_2 du_1,$$

$$\begin{aligned} & \text{case } (2, 2, 2) : \\ K_4 &= \int_{u_1=\chi_2^*}^{\chi_3^*} \int_{u_2=u_1}^{\chi_3^*} \int_{u_3=u_2}^{\chi_3^*} 4 \frac{1}{u_2} du_3 du_2 du_1, \\ & \text{case } (3, 3, 3) : \\ K_5 &= \int_{u_1=\chi_3^*}^1 \int_{u_2=u_1}^1 \int_{u_3=u_2}^1 6 \frac{1}{u_3} du_3 du_2 du_1. \end{aligned}$$

Maximizing $V_5(\chi_1)$ leads to an equation of the type

$$C_1 \chi_1^2 + C_2 \chi_1 + C_3 \ln(\chi_1) + C_4 = 0,$$

where C_i depends on χ_2^*, χ_3^* . This equation does not have an explicit solution.

However, if we introduce a “generalized” Lambert function $W_1(x, y)$ as being defined by

$$W_1 e^{y W_1 + W_1^2} = x,$$

then we can express χ_1^* as

$$\begin{aligned} \chi_1^* &= C_5 W_1(x, y), \\ x &= e^{-C_4/C_3} / C_5, \\ y &= C_2 C_5 / C_3, \\ C_5 &= \sqrt{C_3 / C_1}. \end{aligned}$$

Numerically, this gives $\chi_1^* = .1666171752\dots$ and the success probability $V_5(\chi_1^*) = .1625200069\dots$. Again, $V_5(\chi_1)$ is unimodal in χ_1 .

We note that the much easier problem of predicting an index from which onwards we have exactly three record values has an optimal value of $0.22404\dots$. See Bruss and Paindaveine [4], Sec. 6, equation (13):

$$\begin{aligned} V(\ell) &= \frac{\ell^\ell}{\ell! e^\ell}, \\ \ell = 3 : V(3) &= \frac{9}{2} e^{-3} = 0.22404\dots \end{aligned}$$

Note that the three very best coincide with the last three record values only if the absolute ranks 3, 2, and 1 appear exactly in that order. There is a clear difference between $0.22404\dots$ and the *lower bound* of $.1625200\dots$ we obtained for getting the three *very best* under optimal play. Still, to know now that is actually not that much harder to get the three very best is at least not evident at all.

5 Conclusion

It is in the nature of the studied problem that even asymptotic answers are already computationally involved. Still, this provides a feasible access. Using similar simple probabilistic arguments, we can compute a set of asymptotic thresholds and the final success probability for any κ . Further work would be to build a computer algebra system which could mechanically compute the different cases and the corresponding asymptotic probabilities. This computer algebra should bring the relevant involved functions in a sufficient tractable form to prove the unimodality for all κ in the asymptotic continuous case. (Remember that unimodality is only a sufficient condition and perhaps not needed.) This would be all we need to prove asymptotic optimality also for the case $\kappa \geq 4$.

Hence there remain two interesting open problems:

6 Open problems

PROBLEM 1: Is our procedure for finding the κ best secretaries out of n candidates optimal for all κ ?

PROBLEM 2: Can this procedure be extended to the case where κ depends on n ?

Acknowledgement.

We would like to thank the referee for careful reading and many useful suggestions that improved the paper.

References

- [1] K. Ano, H. Kakinuma, and N. Miyoshi. Odds theorem with multiple selection chances. *Journal of Applied Probability*, 47(4):1093–1104, 2010.
- [2] F.T. Bruss. Sum the odds to one and stop. *Annals of Probability*, 28(3):1384–1391, 2000.
- [3] F.T. Bruss. A note on the odds-theorem of optimal stopping. *Annals of Probability*, 31(4):1859–1861, 2003.
- [4] F.T. Bruss and D. Paindaveine. Selecting a sequence of last successes in independent trials. *Journal of Applied Probability*, 37(2):389–399, 2000.
- [5] Y.S. Chow, S. Moriguti, H. Robbins, and S.M. Samuels. Optimal selection based on relative rank. *Israel Journal of mathematics*, 2(2):81–90, 1964.
- [6] R. Dendievel. New developments of the odds-theorem. *Mathematical Scientist*, 38:111–123, 2013.
- [7] T.S. Ferguson. Who solved the secretary problem? *Statistical Science*, 4(3):282–289, 1989.
- [8] D.V. Lindley. Dynamic programming and decision theory. *Applied Statistics*, 10:35–51, 1961.
- [9] H. Platen. About secretary problems. *Mathematical Statistics, Banach Center Publications*, 6:257–266, 1980.
- [10] J.S. Rose. A problem of optimal choice and assignment. *Operations Research*, 30:172–181, 1982.
- [11] K. Szajowski and D. Lebek. Optimal strategies in high risk investments. *Bulletin of the Belgian Mathematical Society*, 14(1):143–155, 2007.
- [12] M. Tamaki. The secretary problem with optimal assignments. *Operations Research*, 32:847–858, 1984.
- [13] R. J. Vanderbei. The optimal choice of a subset of a population. *Mathematics of Operations Research*, 5(4):481–487, 1980.
- [14] R.J. Vanderbei. The postdoc variant of the secretary problem. Technical report, Princeton University.
- [15] J.G. Wilson. Optimal choice and assignment of the best m of n randomly arriving items. *Stochastic Processes and Their Applications*, 39:325–343, 1991.