

New Hopf Structures on Binary Trees

Stefan Forcey,¹ Aaron Lauve² and Frank Sottile^{2 †}

¹*Department of Mathematics
Tennessee State University
3500 John A Merritt Blvd
Nashville, Tennessee 37209*

²*Department of Mathematics
Texas A&M University
MS 3368
College Station, Texas 77843*

Abstract. The multiplihedra $\mathcal{M}_\bullet = (\mathcal{M}_n)_{n \geq 1}$ form a family of polytopes originating in the study of higher categories and homotopy theory. While the multiplihedra may be unfamiliar to the algebraic combinatorics community, it is nestled between two families of polytopes that certainly are not: the permutahedra \mathfrak{S}_\bullet , and associahedra \mathcal{Y}_\bullet . The maps $\mathfrak{S}_\bullet \rightarrow \mathcal{M}_\bullet \rightarrow \mathcal{Y}_\bullet$ reveal several new Hopf structures on tree-like objects nestled between the Hopf algebras $\mathfrak{S}Sym$ and $\mathcal{Y}Sym$. We begin their study here, showing that $\mathcal{M}Sym$ is a module over $\mathfrak{S}Sym$ and a Hopf module over $\mathcal{Y}Sym$. An elegant description of the coinvariants for $\mathcal{M}Sym$ over $\mathcal{Y}Sym$ is uncovered via a change of basis—using Möbius inversion in posets built on the 1-skeleta of \mathcal{M}_\bullet . Our analysis uses the notion of an *interval retract* that should be of independent interest in poset combinatorics. It also reveals new families of polytopes, and even a new factorization of a known projection from the associahedra to hypercubes.

Résumé. Les multiplihèdres $\mathcal{M}_\bullet = (\mathcal{M}_n)_{n \geq 1}$ forment une famille de polytopes en provenant de l'étude des catégories supérieures et de la théorie de l'homotopie. Tandis que les multiplihèdres sont peu connus dans la communauté de la combinatoire algébrique, ils sont nichés entre deux familles des polytopes qui sont bien connus: les permutahèdres \mathfrak{S}_\bullet , et les associahèdres \mathcal{Y}_\bullet . Les morphismes $\mathfrak{S}_\bullet \rightarrow \mathcal{M}_\bullet \rightarrow \mathcal{Y}_\bullet$ dévoilent plusieurs nouvelles structures de Hopf sur les arbres binaires entre les algèbres de Hopf $\mathfrak{S}Sym$ et $\mathcal{Y}Sym$. Nous commençons son étude ici, en démontrant que $\mathcal{M}Sym$ est un module sur $\mathfrak{S}Sym$ et un module de Hopf sur $\mathcal{Y}Sym$. Une description élégante des coinvariants de $\mathcal{M}Sym$ sur $\mathcal{Y}Sym$ est trouvée par moyen d'un change de base—en utilisant une inversion de Möbius dans certains posets construits sur le 1-squelette de \mathcal{M}_\bullet . Notre analyse utilise la notion d'*interval retract*, ce devrait être intéressante par soi-même dans la théorie des ensembles partiellement ordonnés. Notre analyse donne lieu également à des nouvelles familles des polytopes, et même une nouvelle factorisation d'une projection connue des associahèdres aux hypercubes.

Keywords: multiplihedron, permutations, permutahedron, associahedron, binary trees, Hopf algebras

[†] Sottile supported by NSF grant DMS-0701050

Introduction

In the past 30 years, there has been an explosion of interest in combinatorial Hopf algebras related to the classical ring of symmetric functions. This is due in part to their applications in combinatorics and representation theory, but also in part to a viewpoint expressed in the elegant commuting diagram

$$\begin{array}{ccc} NSym & \hookrightarrow & \mathfrak{S}Sym \\ \downarrow & & \downarrow \\ Sym & \hookrightarrow & QSym \end{array} .$$

Namely, much information about an object may be gained by studying how it interacts with its surroundings. From this picture, we focus on the right edge, $\mathfrak{S}Sym \twoheadrightarrow QSym$. We factor this map through finer and finer structures (some well-known and some new) until this edge is replaced by a veritable zoo of Hopf structures. A surprising feature of our results is that each of these factorizations may be given geometric meaning—they correspond to successive polytope quotients from permutahedra to hypercubes.

The (known) cast of characters

Let us reacquaint ourselves with some of the characters who have already appeared on stage.

$\mathfrak{S}Sym$ – the Hopf algebra introduced by Malvenuto and Reutenauer [13] to explain the isomorphism $QSym \simeq (NSym)^*$. A graded, noncommutative, noncocommutative, self-dual Hopf algebra, with basis indexed by permutations, it offers a natural setting to practice noncommutative character theory [4].

$\mathcal{Y}Sym$ – the (dual of the) Hopf algebra of trees introduced by Loday and Ronco [11]. A graded, noncommutative, noncocommutative Hopf algebra with basis indexed by planar binary trees, it is important for its connections to the Connes-Kreimer renormalization procedure.

$QSym$ – The Hopf algebra of quasisymmetric functions introduced by Gessel [9] in his study of P -partitions. A graded, commutative, noncocommutative Hopf algebra with basis indexed by compositions, it holds a special place in the world of combinatorial Hopf algebras [1].

The new players

In this extended abstract, we study in detail a family of planar binary trees that we call *bi-leveled trees*, which possess two types of internal nodes (circled or not, subject to certain rules). These objects are the vertices of Stasheff’s multiplihedra [18], originating from his study of A_∞ categories. The multiplihedra were given the structure of CW-complexes by Iwase and Mimura [10] and realized as polytopes later [8]. They persist as important objects of study, among other reasons, because they catalog all possible ways to multiply objects in the domain and range of a function f , when both have nonassociative multiplication rules. More recently, they have appeared as moduli spaces of “stable quilted discs” [14].

In Section 2, we define a vector space $\mathcal{M}Sym$ with basis indexed by these bi-leveled trees. We give $\mathcal{M}Sym$ a module structure for $\mathfrak{S}Sym$ by virtue of the factorization

$$\mathfrak{S}Sym \xrightarrow{\beta} \mathcal{M}Sym \xrightarrow{\phi} \mathcal{Y}Sym$$

(evident on the level of planar binary trees) and a splitting $\mathcal{M}Sym \hookrightarrow \mathfrak{S}Sym$. We also show that $\mathcal{M}Sym$ is a Hopf module for $\mathcal{Y}Sym$ and we give an explicit realization of the fundamental theorem of Hopf modules. That is, we find the coinvariants for this action. Our proof, sketched in Section 3, rests on a result about poset maps of independent interest.

We conclude in Section 4 with a massive commuting diagram—containing several new families of planar binary trees—that further factors the map from $\mathfrak{S}Sym$ to $QSym$. The remarkable feature of this diagram is that it comes from polytopes (some of them even new) and successive polytope quotients. Careful study of the interplay between the algebra and geometry will be carried out in future work.

1 Basic combinatorial data

1.1 Ordered and planar binary trees

We recall a map τ from permutations $\mathfrak{S}_n = \bigcup_n \mathfrak{S}_n$ to planar binary trees $\mathcal{Y}_n = \bigcup_n \mathcal{Y}_n$ that has proven useful in many contexts [19, 12]. Its behavior is best described in the reverse direction as follows. Fix a tree $t \in \mathcal{Y}_n$. The n internal nodes of t are equipped with a partial order, viewing the root node as maximal. An *ordered tree* is a planar binary tree, together with a linear extension of the poset of its nodes. These are in bijection with permutations, as the nodes are naturally indexed left-to-right by the numbers $1, \dots, n$. The map τ takes an ordered tree (permutation) to the unique tree whose partial order it extends.

Example 1 The permutations 1423, 2413, and 3412 share a common image under τ :

$$\tau^{-1} \left(\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \right) = \left\{ \begin{array}{c} \color{red}1 \quad \color{red}2 \quad \color{red}3 \\ \color{red}4 \\ \color{red}1 \quad \color{red}3 \\ \color{red}2 \quad \color{red}4 \end{array}, \begin{array}{c} \color{red}2 \quad \color{red}1 \quad \color{red}3 \\ \color{red}4 \\ \color{red}2 \quad \color{red}3 \\ \color{red}1 \quad \color{red}4 \end{array}, \begin{array}{c} \color{red}3 \quad \color{red}1 \quad \color{red}2 \\ \color{red}4 \\ \color{red}3 \quad \color{red}2 \\ \color{red}1 \quad \color{red}4 \end{array} \right\}$$

There are two right inverses to τ that will be useful later. Let $\min(t)$ (respectively, $\max(t)$) denote the unique 231-avoiding (132-avoiding) permutation mapping to t under τ . Loday and Ronco show that $\tau^{-1}(t)$ is the interval $[\min(t), \max(t)]$ in the weak Bruhat order on the symmetric group [12, Thm. 2.5], and that \min and \max are both order-preserving with respect to the Tamari order on \mathcal{Y}_n .

1.2 Bi-leveled trees and the multiplihedra

We next describe a family of *bi-leveled trees* intermediate between the ordered and unordered ones. These trees arrange themselves as vertices of the *multiplihedra* $\mathcal{M}_n = \bigcup_n \mathcal{M}_n$, a family of polytopes introduced by Stasheff in 1970 [18] (though only proven to be polytopes much later [8]). Stasheff introduced this family to represent the fundamental structure of a weak map f between weak structures, such as weak n -categories or A_n spaces. The vertices of \mathcal{M}_n correspond to associations of n objects, pre- and post-application of f , e.g., $(f(a)f(b))f(c)$ and $f(a)f(bc)$. This leads to a natural description of \mathcal{M}_n in terms of “painted binary trees” [5], but we use here the description of Saneblidze and Umble [16].

A *bi-leveled tree* is a pair (t, C) with $t \in \mathcal{Y}_n$ and $C \subseteq [n]$ designating some nodes of t as lower than the others (indexing the nodes from left-to-right by $1, \dots, n$). Viewing t as a poset with root node maximal, C is an increasing order ideal in t where the leftmost node is a minimal element. Graphically, C indexes a collection of nodes of t circled according to the rules: (i) the leftmost node is circled and has no circled children; (ii) if a node is circled, then its parent node is circled.

Define a map β from \mathfrak{S}_n to \mathcal{M}_n as follows. Given a permutation $\sigma = \sigma_1\sigma_2 \cdots \sigma_n$, first represent σ as an ordered tree. Next, forget the ordering on the nodes, save for circling all nodes σ_i with $\sigma_i \geq \sigma_1$.

Example 2 Consider again the permutations 1423, 2413, and 3412 of Example 1. Viewed as ordered trees, their images under β are distinct:

$$\left(\begin{array}{c} \color{red}1 \quad \color{red}2 \quad \color{red}3 \\ \color{red}4 \\ \color{red}1 \quad \color{red}3 \\ \color{red}2 \quad \color{red}4 \end{array}, \begin{array}{c} \color{red}2 \quad \color{red}1 \quad \color{red}3 \\ \color{red}4 \\ \color{red}2 \quad \color{red}3 \\ \color{red}1 \quad \color{red}4 \end{array}, \begin{array}{c} \color{red}3 \quad \color{red}1 \quad \color{red}2 \\ \color{red}4 \\ \color{red}3 \quad \color{red}2 \\ \color{red}1 \quad \color{red}4 \end{array} \right) \xrightarrow{\beta} \left(\begin{array}{c} \circ \quad \circ \quad \circ \\ \circ \\ \circ \quad \circ \\ \circ \quad \circ \end{array}, \begin{array}{c} \circ \quad \circ \quad \circ \\ \circ \\ \circ \quad \circ \\ \circ \quad \circ \end{array}, \begin{array}{c} \circ \quad \circ \quad \circ \\ \circ \\ \circ \quad \circ \\ \circ \quad \circ \end{array} \right).$$

Denote by ϕ the map from bi-leveled trees to trees that forgets which nodes are circled. The map ϕ helps define a partial order on bi-leveled trees that extends the Tamari lattice on planar binary trees: say that the bi-leveled tree s precedes the bi-leveled tree t in the partial order if $\phi(s) \leq \phi(t)$ and the circled nodes satisfy $C_t \subseteq C_s$. We call this the weak order on bi-leveled trees. See Figure 1 below for an example.

The equality $\phi \circ \beta = \tau$ is evident. Remarkably, this factorization

$$\mathfrak{S}_n \xrightarrow{\beta} \mathcal{M}_n \xrightarrow{\phi} \mathcal{Y}_n$$




FIG. 1: The weak order on \mathcal{M}_4 , the bi-leveled trees on 4 nodes.

extends to the level of face maps between polytopes (see Figure 2). Our point of departure was the observation that it is also a factorization as poset maps.




FIG. 2: β and ϕ extend to face (and poset) maps from the permutahedra to the associahedra. The distinguished vertices 1234 , $\beta(1234)$, and $\phi(\beta(1234))$ are indicated.

1.3 Dimension enumeration

Fix a field k of characteristic zero and let $\mathfrak{S}Sym = \bigoplus_{n \geq 0} \mathfrak{S}Sym_n$ denote the graded vector space whose n^{th} graded piece has the “fundamental” basis $\{F_\sigma \mid \sigma \text{ an ordered tree in } \mathfrak{S}_n\}$. Define $\mathcal{M}Sym$ and $\mathcal{Y}Sym$ similarly, replacing \mathfrak{S}_n by \mathcal{M}_n and \mathcal{Y}_n , respectively. We follow convention and say that $\mathfrak{S}Sym_0$ and $\mathcal{Y}Sym_0$ are 1-dimensional. By contrast, we agree that $\mathcal{M}Sym_0 = \{0\}$. (See [7] for categorical rationale; briefly, Stasheff’s \mathcal{M}_1 is already 0-dimensional, so \mathcal{M}_0 has no clear significance.)

In Section 2, we give these three vector spaces a variety of algebraic structures. Here we record some

information about the dimensions of the graded pieces for later reference.

$$\text{Hilb}_q(\mathfrak{S}Sym) = \sum_{n \geq 0} n! q^n = 1 + q + 2q^2 + 6q^3 + 24q^4 + 120q^5 \dots \tag{1}$$

$$\text{Hilb}_q(\mathcal{M}Sym) = \sum_{n \geq 1} A_n q^n = q + 2q^2 + 6q^3 + 21q^4 + 80q^5 + \dots \tag{2}$$

$$\text{Hilb}_q(\mathcal{Y}Sym) = \sum_{n \geq 0} C_n q^n = 1 + q + 2q^2 + 5q^3 + 14q^4 + 42q^5 + \dots \tag{3}$$

Of course, C_n is the n^{th} Catalan number. The enumeration of bi-leveled trees is less familiar: the n^{th} term satisfies $A_n = C_{n-1} + \sum_{k=1}^{n-1} A_k A_{n-k}$ [17, A121988]. A little generating function arithmetic can show that the quotient of (2) by (3) expands as a power series with nonnegative coefficients,

$$\frac{\text{Hilb}_q(\mathcal{M}Sym)}{\text{Hilb}_q(\mathcal{Y}Sym)} = q + q^2 + 3q^3 + 11q^4 + 44q^5 + \dots \tag{4}$$

We will recover this with a little algebra in Section 2.3. The positivity of the quotient of (1) by (3) is established by [3, Theorem 7.2].

2 The Hopf module $\mathcal{M}Sym$

Let τ , β , and ϕ be the maps between the vector spaces $\mathfrak{S}Sym$, $\mathcal{M}Sym$, and $\mathcal{Y}Sym$ induced by τ , β , and ϕ on the fundamental bases. That is, for permutations σ and bi-leveled trees t , we take

$$\tau(F_\sigma) = F_{\tau(\sigma)} \qquad \beta(F_\sigma) = F_{\beta(\sigma)} \qquad \phi(F_t) = F_{\phi(t)}$$


Below, we recall the product and coproduct structures on the Hopf algebras $\mathfrak{S}Sym$ and $\mathcal{Y}Sym$. In [13] and [11], these were defined in terms of the fundamental bases. Departing from these definitions, rich structural information was deduced about $\mathfrak{S}Sym$, $\mathcal{Y}Sym$, and the Hopf algebra map τ between them in [2, 3]. This information was revealed via a change of basis—from fundamental to “monomial”—using Möbius inversion. We take the same tack below with $\mathcal{M}Sym$ and meet with similar success.

2.1 The Hopf algebras $\mathfrak{S}Sym$ and $\mathcal{Y}Sym$

Following [3], we define the product and coproduct structures on $\mathfrak{S}Sym$ and $\mathcal{Y}Sym$ in terms of p -splittings and graftings of trees. A p -splitting of a tree t with n nodes is a forest (sequence) of $p + 1$ trees with n nodes in total. This sequence is obtained by choosing p leaves of t and splitting them (and all parent branchings) right down to the root. By way of example, consider the 3-splitting below (where the third leaf is chosen twice and the fifth leaf is chosen once).



Denote a p -splitting of t by $t \xrightarrow{\gamma} (t_0, \dots, t_p)$. The grafting of a forest (t_0, t_1, \dots, t_p) onto a tree with p nodes is also best described in pictures; for the forest above and $s = \tau(213)$, the tree



is the grafting of (t_0, t_1, \dots, t_p) onto s , denoted $(t_0, t_1, t_2, t_3)/s$. Splittings and graftings of ordered trees are similarly defined. One remembers the labels originally assigned to the nodes of t in a p -splitting, and if t has q nodes, then one increments the labels of s by q in a grafting $(t_0, t_1, t_2, t_3)/s$. See [3] for details.

Definition 3 Fix two ordered or ordinary trees s and t with p and q internal nodes, respectively. We define the product and coproduct by

$$F_t \cdot F_s = \sum_{t \xrightarrow{Y} (t_0, t_1, \dots, t_p)} F_{(t_0, t_1, \dots, t_p)/s} \quad \text{and} \quad \Delta(F_t) = \sum_{t \xrightarrow{Y} (t_0, t_1)} F_{t_0} \otimes F_{t_1}. \quad (5)$$

(In the coproduct for ordered trees, the labels in t_0 and t_1 are reduced to be permutations of $|t_0|$ and $|t_1|$.)

2.2 Module and comodule structures

We next modify the structure maps in (5) to give \mathcal{MSym} the structure of (left) \mathcal{GSym} -module and (right) \mathcal{YSym} -Hopf module. Given a bi-leveled tree b , let $b \xrightarrow{Y} (b_0, \dots, b_p)$ represent any p -splitting of the underlying tree, together with a circling of all nodes in each b_i that were originally circled in b .

Definition 4 (action of \mathcal{GSym} on \mathcal{MSym}) For $w \in \mathcal{G}$ and $s \in \mathcal{M}_p$, write $b = \beta(w)$ and set

$$F_w \cdot F_s = \sum_{b \xrightarrow{Y} (b_0, b_1, \dots, b_p)} F_{(b_0, b_1, \dots, b_p)/s} \quad (6)$$

where the circling rules in $(b_0, b_1, \dots, b_p)/s$ are as follows: every node originating in s is circled whenever $|b_0| > 0$, otherwise, every node originating in $b = \beta(w)$ is uncircled.

This action may be combined with any section of β to define a product on \mathcal{MSym} . For example,

Theorem 5 The action $\mathcal{GSym} \otimes \mathcal{MSym} \rightarrow \mathcal{MSym}$ and the product $\mathcal{MSym} \otimes \mathcal{MSym} \rightarrow \mathcal{MSym}$ are associative. Moreover, putting $\mathcal{MSym}_0 := k$, they make β into an algebra map that factors τ .

Unfortunately, no natural coalgebra structure exists on \mathcal{MSym} that makes β into a Hopf algebra map.

Definition 6 (action and coaction of \mathcal{YSym} on \mathcal{MSym}) Given $b \in \mathcal{M}$, let $b \xrightarrow{Y_+} (b_0, \dots, b_p)$ denote a p -splitting satisfying $|b_0| > 0$. For $s \in \mathcal{Y}_p$, set

$$F_b \cdot F_s = \sum_{b \xrightarrow{Y_+} (b_0, b_1, \dots, b_p)} F_{(b_0, b_1, \dots, b_p)/s} \quad \text{and} \quad \rho(F_b) = \sum_{b \xrightarrow{Y_+} (b_0, b_1)} F_{b_0} \otimes F_{\phi(b_1)}, \quad (7)$$

where in $(b_0, b_1, \dots, b_p)/s$ every node originating in s is circled, and in $\phi(b_1)$ all circles are forgotten.

Example 7 In the fundamental bases of \mathcal{MSym} and \mathcal{YSym} , the action looks like

while the coaction looks like

The significance of our definition of ρ will be seen in Corollary 11. Our next result requires only slight modifications to the original proof that \mathcal{YSym} is a Hopf algebra (due to the restricted p -splittings).

Theorem 8 *The maps $\cdot : \mathcal{MSym} \otimes \mathcal{YSym} \rightarrow \mathcal{MSym}$ and $\rho : \mathcal{MSym} \rightarrow \mathcal{MSym} \otimes \mathcal{YSym}$ are associative and coassociative, respectively. They give \mathcal{MSym} the structure of \mathcal{YSym} -Hopf module. That is, $\rho(F_b \cdot F_s) = \rho(F_b) \cdot \Delta(F_s)$.*

2.3 Main results

We next introduce “monomial bases” for \mathcal{GSym} , \mathcal{MSym} , and \mathcal{YSym} . Given $t \in \mathcal{M}_n$, define

$$M_t = \sum_{t \leq t'} \mu(t, t') F_{t'},$$

where $\mu(\cdot, \cdot)$ is the Möbius function on the poset \mathcal{M}_n . Define the monomial bases of \mathcal{GSym} and \mathcal{YSym} similarly (see (13) and (17) in [3]). The coaction ρ in this basis is particularly nice, but we need a bit more notation to describe it. Given $t \in \mathcal{M}_p$ and $s \in \mathcal{Y}_q$, let $t \setminus s$ denote the bi-leveled tree on $p + q$ internal nodes formed by grafting the root of s onto the rightmost leaf of t .

Theorem 9 *Given a bi-leveled tree t , the coaction ρ on M_t is given by $\rho(M_t) = \sum_{t=t' \setminus s} M_{t'} \otimes M_s$.*

Example 10 Revisiting the trees in the previous example, the coaction in the monomial bases looks like

$$\begin{aligned} \rho(M_{\text{tree}_1}) &= M_{\text{tree}_1} \otimes 1 + M_{\text{tree}_2} \otimes M_{\text{tree}_3}, \\ \rho(M_{\text{tree}_4}) &= M_{\text{tree}_4} \otimes 1 + M_{\text{tree}_5} \otimes M_{\text{tree}_6} + M_{\text{tree}_7} \otimes M_{\text{tree}_8}. \end{aligned}$$

Recall that the *coinvariants* of a Hopf module M over a Hopf algebra H are defined by $M^{\text{co}} = \{m \in M \mid \rho(m) = m \otimes 1\}$. The fundamental theorem of Hopf modules provides that $M \simeq M^{\text{co}} \otimes H$. The monomial basis of \mathcal{MSym} demonstrates this isomorphism explicitly.

Corollary 11 *A basis for the coinvariants in the Hopf module \mathcal{MSym} is given by $\{M_t\}_{t \in \mathcal{T}}$, where \mathcal{T} comprises the bi-leveled trees with no uncircled nodes on their right branches.*

This result explains the phenomenon observed in (4). It also parallels Corollary 5.3 of [3] to an astonishing degree. There, the right-grafting idea above is defined for pairs of planar binary trees and used to describe the coproduct structure of \mathcal{YSym} in its monomial basis.

3 Towards a proof of the main result

We follow the proof of [3, Theorem 5.1], which uses properties of the monomial basis of \mathcal{GSym} developed in [2] to do the heavy lifting. In [3], the section $\mathcal{Y} \xrightarrow{\max} \mathcal{G}$ of τ is shown to satisfy $\tau(M_{\max(t)}) = M_t$ and $\tau(M_\sigma) = 0$ if σ is not 132-avoiding. This was proven using the following result about Galois connections.

Theorem 12 ([15, Thm. 1]) *Suppose P and Q are two posets related by a Galois connection, i.e., a pair of order-preserving maps $\varphi : P \rightarrow Q$ and $\gamma : Q \rightarrow P$ such that for any $v \in P$ and $t \in Q$, $\varphi(v) \leq t \iff v \leq \gamma(t)$. Then the Möbius functions μ_P and μ_Q are related by*

$$\forall v \in P \text{ and } t \in Q, \quad \sum_{\substack{w \in \varphi^{-1}(t), \\ v \leq w}} \mu_P(v, w) = \sum_{\substack{s \in \gamma^{-1}(v), \\ s \leq t}} \mu_Q(s, t).$$


There is a twist in our present situation. Specifically, no Galois connection exists between \mathfrak{S}_n and \mathcal{M}_n . On the other hand, we find that no order-preserving map $\iota : \mathcal{M}_n \hookrightarrow \mathfrak{S}_n$ satisfies $\beta(M_{\iota(t)}) = M_t$. Rather,

$$\beta\left(\sum_{\sigma \in \beta^{-1}(t)} M_\sigma\right) = M_t. \tag{8}$$


This fact is the key ingredient in our proof of Theorem 9. Its verification required modification of the notion of Galois connection—a relationship between posets that we call an *interval retract* (Section 3.2).

3.1 Sections of the map $\beta : \mathfrak{S}_n \rightarrow \mathcal{M}_n$.


Bi-leveled trees t are in bijection with pairs $\{s, \mathbf{s}\}$, where s is a planar binary tree, with p nodes say, and $\mathbf{s} = (s_1, \dots, s_p)$ is a forest (sequence) of planar binary trees. In the bijection, s comprises the circled nodes of t and s_i is the binary tree (of uncircled nodes) sitting above the i^{th} leaf of s . For example,



A natural choice for a section $\iota : \mathcal{M}_n \rightarrow \mathfrak{S}_n$ would be to, say, build $\min(s)$ and $\min(s_i)$ for each i and splice these permutations together in some way to build a word on the letters $\{1, 2, \dots, n\}$. Let $\text{mm}(t)$ denote the choice giving s_1 smaller letters than s_2 , s_2 smaller letters than s_3 , \dots , s_{p-1} smaller letters than s_p , and s_p smaller letters than s :



This choice does not induce a poset map. The similarly defined MM also fails (choosing maximal permutations representing s and \mathbf{s}), but Mm has the properties we need:



We define this map carefully. Given $t \in \mathcal{Y}_n$ and any subset $S \subseteq \mathbb{N}$ of cardinality n , write $\min_S(t)$ for the image of $\min(t)$ under the unique order-preserving map from $[n]$ to S ; define $\max_S(t)$ similarly.

Definition 13 [The section Mm] Let $t \leftrightarrow \{s, \mathbf{s}\}$ be a bi-leveled tree on n nodes with p circled nodes. Write $u = u_1 \cdots u_p = \min_{[a,b]}(s)$ for $[a, b] = \{n - p + 1, \dots, n\}$ and write $v^i = \max_{[a_i, b_i]}(s_i)$ ($1 \leq i \leq p$), where the intervals $[a_i, b_i]$ are defined recursively as follows:

$$a_p = 1 \quad \text{and} \quad b_p = a_p + |s_p| - 1,$$

$$a_i = 1 + \max_{j>i} \bigcup S_j \quad \text{and} \quad b_i = a_i + |s_i| - 1.$$

Finally, define $\text{Mm}(t)$ by the concatenation $\text{Mm}(t) = u_1 v^1 u_2 v^2 \cdots u_p v^p$.

Remark 14 Alternatively, $\text{Mm}(t)$ is the unique $w \in \beta^{-1}(t)$ avoiding the pinned patterns $\underline{0}231$, $\underline{3}021$, and $\underline{2}031$, where the underlined letter is the first letter in w . The first two patterns fix the embeddings of s_i ($0 \leq i \leq p$), the last one makes the letters in s_i larger than those in s_{i+1} ($1 \leq i < p$).

The important properties of $Mm(t)$ are as follows.

Proposition 15 *The section $\iota : \mathcal{M}_n \rightarrow \mathfrak{S}_n$ given by $\iota(t) = Mm(t)$ is an embedding of posets. The map $\beta : \mathfrak{S}_n \rightarrow \mathcal{M}_n$ satisfies $\beta(\iota(t)) = t$ for all $t \in \mathcal{M}_n$ and $\beta^{-1}(t) \subseteq \mathfrak{S}_n$ is the interval $[mm(t), MM(t)]$.*

3.2 Interval retracts

Let $\varphi : P \rightarrow Q$ and $\gamma : Q \rightarrow P$ be two order-preserving maps between a lattice P and a poset Q . If

$$\forall t \in Q \quad \varphi(\gamma(t)) = t \quad \text{and} \quad \varphi^{-1}(t) \text{ is an interval,}$$

then we say that φ and γ demonstrate P as an *interval retract* of Q .

Theorem 16 *If P and Q are two posets related by an interval retract (φ, γ) , then the Möbius functions μ_P and μ_Q are related by*

$$\forall s < t \in Q \quad \sum_{\substack{v \in \varphi^{-1}(s) \\ w \in \varphi^{-1}(t)}} \mu_P(v, w) = \mu_Q(s, t).$$


The proof of Theorem 16 exploits Hall’s formula for Möbius functions. An immediate consequence is a version of (8) for any P and Q related by an interval retract. Verifying that (β, Mm) is an interval retract between \mathfrak{S}_n and \mathcal{M}_n (Proposition 15) amounts to basic combinatorics of the weak order on \mathfrak{S}_n .

4 More families of binary trees and their polytopes

We have so far ignored the algebra $QSym$ of quasisymmetric functions advertised in the introduction. A basis for its n^{th} graded piece is naturally indexed by compositions of n , but may also be indexed by trees as follows. To a composition (a_1, a_2, \dots) , say $(3, 2, 1, 4)$, we associate a sequence of *left-combs*


$$(\begin{array}{c} \diagup \diagdown \\ \diagup \diagdown \\ \diagup \diagdown \end{array}, \begin{array}{c} \diagup \diagdown \\ \diagup \diagdown \end{array}, \begin{array}{c} \diagup \diagdown \\ \diagup \diagdown \end{array}, \begin{array}{c} \diagup \diagdown \\ \diagup \diagdown \\ \diagup \diagdown \\ \diagup \diagdown \end{array}),$$

i.e., trees with a_i leaves and all internal leaves rooted to the rightmost branch and left-pointing. These may be hung on another tree, a right-comb with right-pointing leaves, to establish a bijection between compositions of n and “combs of combs” with n internal nodes:



To see how $QSym$ and the hypercubes fit into the picture, we briefly revisit the map β of Section 1.2.

We identified bi-leveled trees with pairs $\{s, s\}$, where s is the tree of circled nodes and $s = (s_0, \dots, s_p)$ is a forest of trees (the uncircled nodes). Under this identification, β may be viewed as a pair of maps (τ, τ) —with the first factor τ making a (planar binary) trees out of the nodes greater than or equal to σ_1 , and the second factor τ making trees out of the smaller nodes:



See also Figure 3. Two more fundamental maps are γ_r and γ_l , taking trees to (right- or left-) combs, e.g.,




Figure 3 displays several combinations of the maps τ , γ_r , and γ_l . The algebra \mathcal{QSym} corresponds to the terminal object there—the set denoted $\left\{ \frac{\dots \text{combs} \dots}{\text{comb}} \right\}$.

The new binary tree-like structures appearing in the factorization of $\mathcal{S}Sym \rightarrow \mathcal{QSym}$ (i.e., those trees not appearing on the central, vertical axis of Figure 3) will be studied in upcoming papers. It is no surprise that $\mathcal{S}Sym \rightarrow \mathcal{QSym}$ factors through so many intermediate structures. What is remarkable, and what our binary tree point-of-view reveals, is that each family of trees in Figure 3 can be arranged into a family of polytopes. See Figure 4. The (Hopf) algebraic and geometric implications of this phenomenon will also be addressed in future work.

References

- [1] Marcelo Aguiar, Nantel Bergeron, and Frank Sottile, *Combinatorial Hopf algebras and generalized Dehn-Sommerville relations*, *Compos. Math.* **142** (2006), no. 1, 1–30.
- [2] Marcelo Aguiar and Frank Sottile, *Structure of the Malvenuto-Reutenauer Hopf algebra of permutations*, *Adv. Math.* **191** (2005), no. 2, 225–275.
- [3] ———, *Structure of the Loday-Ronco Hopf algebra of trees*, *J. Algebra* **295** (2006), no. 2, 473–511.
- [4] Dieter Blessenohl and Manfred Schocker, *Noncommutative character theory of the symmetric group*, Imperial College Press, London, 2005.
- [5] J. M. Boardman and R. M. Vogt, *Homotopy invariant algebraic structures on topological spaces*, *Lecture Notes in Mathematics*, Vol. 347, Springer-Verlag, Berlin, 1973.
- [6] Satyan Devadoss and Stefan Forcey, *Marked tubes and the graph multiplihedron*, preprint, arXiv:0807.4159.
- [7] Stefan Forcey, *Quotients of the multiplihedron as categorified associahedra*, preprint, arXiv:0803.2694.
- [8] ———, *Convex hull realizations of the multiplihedra*, *Topology Appl.* **156** (2008), 326–347.
- [9] Ira M. Gessel, *Multipartite P -partitions and inner products of skew Schur functions*, *Combinatorics and algebra* (Boulder, Colo., 1983), *Contemp. Math.*, vol. 34, Amer. Math. Soc., Providence, RI, 1984, pp. 289–317.
- [10] Norio Iwase and Mamoru Mimura, *Higher homotopy associativity*, *Algebraic topology* (Arcata, CA, 1986), *Lecture Notes in Math.*, vol. 1370, Springer, Berlin, 1989, pp. 193–220.
- [11] Jean-Louis Loday and María O. Ronco, *Hopf algebra of the planar binary trees*, *Adv. Math.* **139** (1998), no. 2, 293–309.
- [12] ———, *Order structure on the algebra of permutations and of planar binary trees*, *J. Algebraic Combin.* **15** (2002), no. 3, 253–270.
- [13] Claudia Malvenuto and Christophe Reutenauer, *Duality between quasi-symmetric functions and the Solomon descent algebra*, *J. Algebra* **177** (1995), no. 3, 967–982.
- [14] Sikimeti Mau and Chris Woodward, *Geometric realizations of the multiplihedron and its complexification*, preprint, arXiv:0802.2120v4.
- [15] Gian-Carlo Rota, *On the foundations of combinatorial theory. I. Theory of Möbius functions*, *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **2** (1964), 340–368 (1964), Reprinted in: J.P.S. Kunt (Ed.), *Gian-Carlo Rota on Combinatorics: Introductory Papers and Commentaries*, Birkhäuser, Boston, 1995.
- [16] Samson Sanedidze and Ronald Umble, *Diagonals on the permutahedra, multiplihedra and associahedra*, *Homology Homotopy Appl.* **6** (2004), no. 1, 363–411 (electronic).
- [17] N. J. A. Sloane, *The on-line encyclopedia of integer sequences*, published electronically at www.research.att.com/~njas/sequences/.
- [18] James Stasheff, *H-spaces from a homotopy point of view*, *Lecture Notes in Mathematics*, Vol. 161, Springer-Verlag, Berlin, 1970.
- [19] Andy Tonks, *Relating the associahedron and the permutohedron*, *Operads: Proceedings of Renaissance Conferences* (Hartford, CT/Luminy, 1995), *Contemp. Math.*, vol. 202, Amer. Math. Soc., Providence, RI, 1997, pp. 33–36.




FIG. 3: A commuting diagram of tree-like objects. The spaces $\mathfrak{S}Sym$, \mathcal{MSym} , \mathcal{YSym} and \mathcal{QSym} appear, top to bottom, along the center. The unlabeled dashed line represents the usual map from \mathcal{YSym} to \mathcal{QSym} (see [11], Section 4.4). It is incompatible with the given map $(\gamma_r, \gamma_l) : \mathcal{MSym} \rightarrow \mathcal{QSym}$.

We explore the Hopf module structures of objects mapping to \mathcal{YSym} and \mathcal{QSym} in future work. At least some of these will be full-fledged Hopf algebras (e.g., note that there is a bijection of sets between $\{\frac{\dots \text{trees} \dots}{\text{comb}}\}$ and $\{\text{trees}\}$, the latter indexing the Hopf algebra \mathcal{YSym}).




FIG. 4: A commuting diagram of polytopes based on tree-like objects with 4 nodes, corresponding position-wise to Figure 3 (image of ϕ is suppressed). Notation is taken from [6]: $\mathcal{P}(4)$ is the permutohedron, $\mathcal{J}(4)$ is the multiplihedron, $\mathcal{K}(5)$ is the associahedron, and $\mathcal{CK}(4)$ is the composihedron. \mathcal{JG}_d is the domain quotient of the permutohedron and \mathcal{JG}_r is its range quotient.

The cellular projections shown include neither the Tonks projection nor the Loday Ronco projection from the associahedron to the hypercube. However, the map from the multiplihedron to the cube passing through the associahedron appears in [5].