

# Generalized triangulations, pipe dreams, and simplicial spheres

Luis Serrano and Christian Stump

*LaCIM, Université du Québec à Montréal, Montréal, Canada*

---

**Abstract.** We exhibit a canonical connection between maximal  $(0, 1)$ -fillings of a moon polyomino avoiding north-east chains of a given length and reduced pipe dreams of a certain permutation. Following this approach we show that the simplicial complex of such maximal fillings is a vertex-decomposable and thus a shellable sphere. In particular, this implies a positivity result for Schubert polynomials. For Ferrers shapes, we moreover construct a bijection to maximal fillings avoiding south-east chains of the same length which specializes to a bijection between  $k$ -triangulations of the  $n$ -gon and  $k$ -fans of Dyck paths. Using this, we translate a conjectured cyclic sieving phenomenon for  $k$ -triangulations with rotation to  $k$ -flagged tableaux with promotion. **Résumé.** Nous décrivons un lien canonique entre les  $(0, 1)$ -remplissages maximaux d'un polyomino-lune évitant les chaînes Nord-Est d'une longueur donnée, et les "pipe dreams" réduits d'une certaine permutation. En suivant cette approche nous montrons que le complexe simplicial de tels remplissages maximaux est une sphère "vertex-decomposable" et donc "shellable". En particulier, cela entraîne un résultat de positivité sur les polynômes de Schubert. De plus, nous construisons, dans le cas des diagrammes de Ferrers, une bijection vers les remplissages maximaux évitant les chaînes Sud-Est de même longueur, qui se spécialise en une bijection entre les  $k$ -triangulations d'un  $n$ -gone et les  $k$ -faisceaux de chemins de Dyck. À l'aide de celle-ci, nous traduisons une instance conjecturale du phénomène de tamis cyclique pour les  $k$ -triangulations avec rotation dans le cadre des tableaux  $k$ -marqués avec promotion.

**Keywords:**  $k$ -triangulation, enumerative combinatorics, pipe dream, fans of Dyck paths, flagged Schur function, Schubert polynomial, Edelman-Greene insertion

---

## 1 Introduction

Fix positive integers  $n$  and  $k$  such that  $2k < n$ . A  **$k$ -triangulation** of a convex  $n$ -gon is a maximal collection of diagonals in the  $n$ -gon such that no  $k + 1$  diagonals mutually cross. A  **$k$ -fan of Dyck paths** of length  $2\ell$  is a collection of  $k$  Dyck paths from  $(0, 0)$  to  $(\ell, \ell)$  which do not cross (although they may share edges).

The following theorem is the first main result in this article. It extends results by S. Elizalde [Eli07] and C. Nicolás [Nic09].

**Theorem 1.1** *There is an explicit bijection between  $k$ -triangulations of a convex  $n$ -gon and  $k$ -fans of Dyck paths of length  $2(n - 2k)$ .*

A **north-east chain** of length  $\ell$  in a Ferrers shape  $\lambda$  is a sequence of  $\ell$  boxes in  $\lambda$  such that every box in the sequence is strictly north and strictly east of the preceding one, and for which the smallest rectangle containing all boxes in the sequence is also contained in  $\lambda$ . A  **$k$ -north-east filling** of  $\lambda$  is a  $(0, 1)$ -filling which does not contain any north-east chain of 1's of length  $k + 1$ , and in which the number of 1's is maximal. As usual, we identify a  $(0, 1)$ -filling with its set of boxes filled with 1's and draw them by marking its set of boxes by +'s. See Figure 1(a) for an example. The set of all  $k$ -north-east fillings of  $\lambda$  is denoted by  $\mathcal{F}_{NE}(\lambda, k)$ . **South-east chains**,  **$k$ -south-east fillings** and  $\mathcal{F}_{SE}(\lambda, k)$  are defined similarly.

It is well known that  $k$ -triangulations of the  $n$ -gon can be seen as  $k$ -north-east fillings of the staircase shape  $(n - 1, \dots, 2, 1)$ , and furthermore,  $k$ -fans of Dyck paths of length  $2(n - 2k)$  can be seen as  $k$ -south-east fillings of the same staircase (see e.g. [Kra06, Rub06]). Thus, the second main theorem is a clear extension of the first. It answers a questions raised by C. Krattenthaler in [Kra06].

**Theorem 1.2** *Let  $\lambda$  be a Ferrers shape and let  $k$  be a positive integer. There is an explicit bijection between  $k$ -north-east and  $k$ -south-east fillings of  $\lambda$ .*

The constructed bijection goes through two intermediate objects, namely through pipe dreams and flagged tableaux, both arising in the theory of Schubert polynomials. The third main theorem is a central step in the proof of Theorem 1.2 and it concerns the connection between north-east chains and reduced pipe dreams.

**Theorem 1.3** *Let  $\lambda$  be a Ferrers shape and let  $k$  be a positive integer. There exists a canonical bijection between  $k$ -north-east fillings of  $\lambda$  and reduced pipe dreams of a permutation depending on  $\lambda$  and  $k$ .*

This bijection will be described in Section 2. A variation of the argument gives the following generalization to moon polyominoes as defined in Section 2.2.

**Theorem 1.4** *Let  $M$  be a moon polyomino and let  $k$  be a positive integer. Then there exists a canonical bijection between  $k$ -north-east fillings of  $M$  and reduced pipe dreams (of a given permutation) living inside  $M$ .*

We will use the construction to obtain new properties and simple proofs for known properties of  $k$ -north-east fillings and of  $k$ -triangulations. In particular, we obtain the following corollaries.

**Corollary 1.5** *The simplicial complex with facets being  $k$ -north-east fillings of a moon polyomino  $M$  is the join of a vertex-decomposable, triangulated sphere with a full simplex. In particular, it is shellable and Cohen-Macaulay.*

**Corollary 1.6** *Let  $S$  be a stack polyomino and  $\lambda$  the Ferrers shape obtained from  $S$  by properly rearranging its columns. Let  $\sigma$  and  $\tau$  be the associated permutations. Then the difference*


$$\mathfrak{S}_\sigma(x_1, x_2, \dots) - \mathfrak{S}_\tau(x_1, x_2, \dots)$$

*of Schubert polynomials is monomial positive.*

The bijection for  $k$ -triangulations has the additional property that the cyclic action given by rotation of the  $n$ -gon corresponds to a promotion-like operation on flagged tableaux and thus transforms a conjectured cyclic sieving phenomenon (CSP) into the context of  $k$ -flagged tableaux.

**Conjecture 1.7** *Let  $\mathcal{FT}(\lambda, k)$  be the set of  $k$ -flagged tableaux and let  $\rho$  be the promotion-like cyclic action on  $\mathcal{FT}(\lambda, k)$ . The triple*

$$\left( \mathcal{FT}(\lambda, k), \langle \rho \rangle, F(q) \right),$$



**Fig. 1:** A 2-north-east filling of  $\lambda = (8, 6, 6, 5, 4, 4, 1)$  and its associated reduced pipe dream.

exhibits the CSP, where

$$F(q) := \prod_{1 \leq i \leq j < n-2k} \frac{[i + j + 2k]_q}{[i + j]_q}$$

is a natural  $q$ -analogue of the cardinality of  $\mathcal{F}_{NE}(\lambda, k)$ .


## 2 From north-east fillings to pipe dreams

In this section we exhibit a connection between  $k$ -north-east fillings of Ferrers shapes as well as of stack and moon polyominoes on the one hand and reduced pipe dreams on the other. This generalizes a construction by the second author for  $k$ -triangulations [Stu10].

Reduced pipe dreams (or  $rc$ -graphs) were introduced by S. Fomin and A. Kirillov in [FK96] (see also [BB93] and [KM05, Section 1.4]). They play a central role in the combinatorics of Schubert polynomials of A. Lascoux and M.-P. Schützenberger. A **pipe dream** of size  $n$  is a filling of the staircase shape  $(n-1, \dots, 2, 1)$  where each box contains two crossing pipes  $\times$  or two turning pipes  $\curvearrowright$ . See Figure 1(b) for an example. A pipe dream is identified with its set of boxes containing two crossing pipes  $\times$ . The permutation  $\pi(D)$  of a pipe dream  $D$  is obtained by following the pipes starting from the top and going all the way to the left, and then reading  $\pi(D)$  on the left from top to bottom in one line notation. For example, the permutation of the pipe dream in Figure 1(b) is  $[1, 2, 7, 6, 5, 8, 3, 4, 9, 10]$ . A pipe dream is **reduced** if two pipes cross at most once. We say that a pipe dream lives inside a set  $M$  of boxes in the staircase shape if all its crossings are contained in  $M$ . For a given permutation  $\pi$  and a set  $M$  of boxes, denote the set of reduced pipe dreams for  $\pi$  by  $\mathcal{RP}(\pi)$  and the set of reduced pipe dreams for  $\pi$  which live inside  $M$  by  $\mathcal{RP}(\pi, M)$ .

### 2.1 A bijection between north-east fillings and reduced pipe dreams

Starting with a  $k$ -north-east filling of  $\lambda$ , one obtains a pipe dream by replacing every 1 by two turning pipes and every 0 by two crossing pipes. Afterwards,  $\lambda$  is embedded into the smallest staircase containing it, and all boxes in the staircase outside of  $\lambda$  are replaced by turning pipes. In other words, a  $k$ -north-east



**Fig. 2:**  $\sigma_2(\lambda) = [1, 2, 7, 6, 5, 8, 3, 4, 9, 10]$  for  $\lambda = (8, 6, 6, 5, 4, 4, 1)$ .

filling of  $\lambda$  and its associated pipe dream are complementary  $(0, 1)$ -fillings of  $\lambda$  when both are identified with their sets of boxes. For example, the  $\perp$ 's in the pipe dream in Figure 1(b) and the marked boxes in (a) are complementary  $(0, 1)$ -fillings of  $\lambda$ . The pieces in boxes outside of  $\lambda$  are drawn in the pipe dream in red whereas pieces within  $\lambda$  are drawn in green. We call this identification between  $k$ -north-east fillings of  $\lambda$  and reduced pipe dreams **complementary map**.

For a permutation  $\sigma \in \mathcal{S}_n$ , define its **(Rothe) diagram** (see [Man01, Section 2.1]) to be the set of boxes in the staircase shape given by

$$D(\sigma) := \{(i, \sigma_j) : i < j, \sigma_i > \sigma_j\}.$$

For example, the diagram of  $\sigma_2(\lambda)$  in Figure 2 is given by the shaded area. Clearly, the number of boxes in  $D(\sigma)$  equals the **length** of  $\sigma$ , i.e., the minimal number of simple transpositions needed to write  $\sigma$ . A permutation is called **dominant** if its diagram is a Ferrers shape containing the box  $(1, 1)$ . By construction, different permutations in  $\mathcal{S}_n$  have different shapes and one can obtain every Ferrers shape in this way for some  $n$ . Thus, starting with a Ferrers shape  $\lambda$ , let  $\sigma(\lambda)$  be the unique dominant permutation  $\sigma \in \mathcal{S}_n$  for which  $D(\sigma) = \lambda$ , where  $n$  is given by the smallest staircase shape containing  $\lambda$ . Moreover, define  $\sigma_k(\lambda)$  to be

$$\mathbf{1}_k \times \tau := [1, 2, \dots, k, \tau_1 + k, \dots, \tau_n + k] \in \mathcal{S}_{n+k}$$

where  $\tau = \sigma(\mu)$  and  $\mu$  is obtained from  $\lambda$  by removing its first  $k$  rows and columns. Graphically, this means that  $\sigma_k(\lambda)$  is obtained by removing the first  $k$  columns and rows from  $\sigma(\lambda)$ . Note that the north-west corner of  $\sigma_k(\lambda)$  remains in box  $(k + 1, k + 1)$ . See Figure 2 for  $\sigma_2(\lambda)$  with  $\lambda$  as in Figure 1. The following theorem is a more precise reformulation of Theorem 1.3.

**Theorem 2.1** *Let  $\lambda$  be a Ferrers shape and let  $\sigma = \sigma_k(\lambda)$ . The complementary map from  $k$ -north-east fillings to pipe dreams is a bijection between  $\mathcal{F}_{NE}(\lambda, k)$  and  $\mathcal{RP}(\sigma)$ .*

### 2.2 Generalizations to moon polyominoes

The results in the previous section can be partially generalized to moon polyominoes which were studied by J. Jonsson in [Jon05]. A polyomino  $M$  (i.e., a set of boxes in the positive integer quadrant) is called




Fig. 3: A 1-north-east filling of a moon polyomino and its associated pipe dream.


**convex** if for any two boxes in  $M$  lying in the same row or column, all boxes in between are also contained in  $M$ . Moreover,  $M$  is called **intersection-free** if for any two columns (or equivalently, rows) of  $M$ , one is contained in the other. A polyomino is called a **moon polyomino** if it is convex and intersection-free.

Without loss of generality we consider always moon polyominoes which are north-west justified, namely, they contain boxes both in the first row and in the first column. Observe that Ferrers shapes are special types of moon polyominoes. A  $k$ -filling of a moon polyomino is defined exactly in the same way as for a Ferrers shape. See Figure 3 for an example.

To connect  $k$ -north-east fillings of a moon polyomino  $M$  and pipe dreams of a certain permutation  $\sigma = \sigma_k(M)$ , we must relate maximal fillings of  $M$  which do not contain a  $(k + 1)$ -north-east chain in one of its maximal rectangles and maximal fillings of the staircase  $(n - 1, \dots, 2, 1)$  which do not contain a north-east chain of length  $r_\sigma(p, q) + 1$  in any rectangle  $[p] \times [q]$ . Define  $\sigma_k(M)$  as follows: for a maximal rectangle  $R$  in  $M$  of width and height both strictly larger than  $k$ , let  $(a + 1, b + 1)$  and  $(i, j)$  be its north-west and south-east corner boxes. Mark the box  $(i + b, j + a)$  with  $a + b + k$ .  $\sigma_k(M)$  is the permutation with this collection as its essential set. This means that the diagram  $D(\sigma)$  of  $\sigma$  has  $(i + b, j + a)$  as a south-east corner with labels  $r_\sigma(i + b, j + a) = a + b + k$ . Using [Man01, 2.2.8], it is easy to see that this construction is well defined. Note that maximal rectangles of width or height less than or equal to  $k$  cannot contain any north-east chain of length larger than  $k$  and thus do not contribute to the essential set of the corresponding permutation. For example, the moon polyomino  $M$  in Figure 3(a) has maximal rectangles  $(a + 1, b + 1) - (i, j)$  given by

$$\begin{aligned} &(1, 3) - (5, 5), \quad (1, 3) - (6, 4), \\ &(3, 1) - (4, 7), \quad (2, 2) - (5, 5), \\ &(2, 1) - (4, 6), \end{aligned}$$

where the first maximal rectangle is highlighted. Thus, for  $k = 1$ , the resulting essential set and the associated diagram can be seen in Figure 4 and the associated permutation is  $\sigma_1(M) = [1, 2, 8, 10, 3, 7, 6, 5, 4, 9]$ . As all maximal rectangles in a Ferrers shape are of the form  $[p] \times [q]$ , the definition of  $\sigma_k(\lambda)$  reduces in



**Fig. 4:** The essential set and the diagram  $D(\sigma)$  for  $\sigma = [1, 2, 8, 10, 3, 7, 6, 5, 4, 9]$ .

this case to the definition given in the previous section.

The following theorem is a more precise reformulation of Theorem 1.4.

**Theorem 2.2** *The complementary map from  $k$ -north-east fillings of a moon polyomino  $M$  to pipe dreams of  $\sigma = \sigma_k(M)$  is a bijection between  $\mathcal{F}_{NE}(M, k)$  and  $\mathcal{RP}(\sigma, M)$ .*


We now use this theorem together with the main theorem in [Jon05] to get new insights on pipe dreams. A **stack polyomino** is a moon polyomino where every column starts in the first row. Let  $S$  be a stack polyomino and let  $\lambda$  be the Ferrers shape obtained from  $S$  by properly rearranging the columns. J. Jonsson proved in [Jon05, Theorem 14] that the number of  $k$ -north-east fillings of  $S$  with a given number of  $+$ 's in every row equals the number of  $k$ -north-east fillings in  $\lambda$  with the same number of  $+$ 's in every row. Moreover, he conjectured that this property still holds if the stack polyomino  $S$  is replaced by a moon polyomino. Therefore, we obtain the following corollary and the conjecture for the analogous statement for moon polyominoes.

**Corollary 2.3** *Let  $S$  be a stack polyomino and let  $\lambda$  be the associated Ferrers shape. The number of pipe dreams in  $\mathcal{RP}(\sigma_k(S), S)$  with a given number of crossings in every row is equal to the number of pipe dreams in  $\mathcal{RP}(\sigma_k(\lambda))$  with the same number of crossings in every row.*

### 2.3 The simplicial complex of north-east-fillings

We are now in position to obtain Corollary 1.5. The canonical connection between  $k$ -north-east fillings and reduced pipe dreams can be used in the same way as described in the proof of [Stu10, Corollary 1.3] for  $k$ -triangulations in this more general setting. For the necessary background on simplicial complexes and in particular on subword complexes, we refer to [KM04]. A box in a moon polyomino  $M$  is called **passive** if it is not contained in any north-east chain in  $M$  of length  $k + 1$ . Let  $\Delta(M, k)$  be the simplicial complex with vertices being the collection of boxes in  $M$ , and with facets being  $k$ -north-east fillings of  $M$ .

**Corollary 2.4**  *$\Delta(M, k)$  is the join of a vertex-decomposable, triangulated sphere and a full simplex of dimension  $i - 1$ , where  $i$  equals the number of passive boxes in  $M$ . In particular, it is shellable and Cohen-Macaulay.*



**Fig. 5:** Labelling of the crossing boxes in the pipe dream in Figure 1(b), the corresponding compatible sequence, and its insertion and recording tableau.

### 2.4 A mutation-like operation on pipe dreams

Generalizing the notion in the previous section, one can define a pure simplicial complex  $\Delta(\sigma)$  for any  $\sigma \in \mathcal{S}_n$  by defining the facets as the complements in the staircase of reduced pipe dreams in  $\mathcal{RP}(\sigma)$  (see [KM04]). Using the property that two pipes in a reduced pipe dream  $D$  cross at most once, one can define a mutation-like operation on facets of  $\Delta(\sigma)$  as follows. One can mutate the facet  $F(D)$  of  $\Delta(\sigma)$  associated to  $D$  at a vertex  $b$  if the two pipes in  $D$  which touch in  $b$  cross somewhere else. In other words, one can mutate  $F(D)$  at a vertex  $b$  if the starting points  $i < j$  of the two pipes in  $D$  which touch in  $b$  form an inversion of  $\sigma$ . The mutation of  $F(D)$  at such a vertex  $b$  is then defined to be the facet  $F(D')$  for the reduced pipe dream  $D'$  such that

- (i) the two turning pipes in  $b$  are replaced in  $D'$  by two crossing pipes,
- (ii) the unique crossing  $b'$  of those two pipes is replaced in  $D'$  by two turning pipes.

By construction, the pipe dream  $D' = (D \cup b) \setminus b'$  is again in  $\mathcal{RP}(\sigma)$  and thus its complement  $F(D') = (F(D) \setminus b) \cup b'$  forms another facet of  $\Delta(\sigma)$ .

## 3 From pipe dreams to south-east fillings

In this section we describe a bijection between pipe dreams for  $\sigma_k(\lambda)$  and  $k$ -south east fillings of  $\lambda$ , for a Ferrers shape  $\lambda$ . For the sake of readability, we do this construction in several steps.

### 3.1 From pipe dreams to flagged tableaux

Define a  **$k$ -flagged tableau** as a semistandard tableau in which the entries in the  $i$ -th row are smaller than or equal to  $i + k$ , and denote the set of  $k$ -flagged tableaux of shape  $\lambda$  by  $\mathcal{FT}(\lambda, k)$ . These were introduced by M. Wachs [Wac85] in the study of **flagged Schur functions**, thus the choice of terminology. We now present a bijection between the set  $\mathcal{RP}(\sigma)$  of reduced pipe dreams of  $\sigma = \sigma_k(\lambda)$  and the set  $\mathcal{FT}(\mu, k)$  of  $k$ -flagged tableaux of shape  $\mu = D(\sigma)$ .

For a reduced pipe dream  $D \in \mathcal{RP}(\sigma)$  with  $\sigma$  being of length  $\ell$ , define the **reading biword** to be the  $2 \times \ell$  array by reading  $\binom{i}{i+j-1}$  for every crossing box  $(i, j)$  in  $D$  row by row from east to west and from north to south. See Figure 5 for an example. It is known (and easy to check) that this gives a bijection between  $\mathcal{RP}(\sigma)$  and the set of **compatible sequences**  $\mathcal{CS}(\sigma)$ , defined by S. Billey, W. Jockush

and R. Stanley in [BJS93] as the set of all  $2 \times \ell$  arrays of the form  $\begin{pmatrix} a_1, \dots, a_\ell \\ b_1, \dots, b_\ell \end{pmatrix}$  satisfying the following properties:

1.  $a_1 \leq a_2 \leq \dots \leq a_\ell$ ,
2. if  $a_i = a_{i+1}$ , then  $b_i > b_{i+1}$ ,
3.  $b_1 b_2 \dots b_\ell$  is a reduced word for  $\sigma$ , where  $i$  denotes the simple transposition  $s_i = (i, i + 1)$ , and
4.  $a_i \leq b_i$ .

One can see from the definition that a compatible sequence  $t$  for  $\sigma$  can be written as the concatenation  $t = t_1 \dots t_m$ , where  $t_i = \begin{pmatrix} i^{w_i} \\ w_i \end{pmatrix}$ , and  $w_i$  is decreasing. Observe that  $\sigma$  fixes all  $j \leq k$  and thus, every letter in  $w_i$  is larger than or equal to  $\max(i, k)$ .

Define a map  $\mathcal{CS}(\sigma) \rightarrow \mathcal{FT}(\mu, k)$  as follows. Let  $t \in \mathcal{CS}(\sigma)$  be a compatible sequence for  $\sigma$ . Insert the letters of the word formed by the bottom row of  $t$  using column Edelman–Greene insertion [EG87] into a tableau, while recording the corresponding letters from the first row. This produces an insertion tableau  $P(t)$  and a recording tableau  $Q(t)$ . The image in  $\mathcal{FT}(\mu, k)$  is now defined to be  $Q(t)$ .

**Theorem 3.1** *Let  $\sigma = \sigma_k(\lambda)$  for a Ferrers shape  $\lambda$ , and let  $\mu = D(\sigma)$ . The map sending  $D$  in  $\mathcal{RP}(\sigma)$  to the recording tableau of the reading biword of  $D$  is a bijection between  $\mathcal{RP}(\sigma)$  and  $\mathcal{FT}(\mu, k)$ .*

An example of the bijection can be seen in Figure 5.

### 3.2 A cyclic action on flagged tableaux

In this subsection we define a cyclic action on  $k$ -flagged tableaux. The **flagged promotion**  $\rho(Q)$  of a  $k$ -flagged tableau  $Q$  is defined as follows.


- (i) Delete all the instances of the letter 1,
- (ii) apply jeu de taquin to the remaining entries,
- (iii) subtract 1 from all the entries,
- (iv) label each empty box on row  $i$  with  $i + k$ .

One can easily see that  $\rho(Q)$  is indeed a  $k$ -flagged tableau, since the empty boxes after step (iii) must form a horizontal strip, which means there is at most one empty box per column. Furthermore, as every box gets moved at most up by one row, and at the end one subtracts 1 from all the entries, the tableau obtained after step (iii) is  $k$ -flagged as well. The argument is finalized with the observation that if one adds a horizontal strip in which every box gets added its maximum possible value, the tableau is still  $k$ -flagged, since the row-weakness is assured by the maximality of the value of the entries on each row, and the column-strictness is assured by the fact that the entries in row  $i - 1$  are all strictly less than the maximal value on row  $i$ .

### 3.3 From flagged tableaux to fans of paths and south-east fillings

We proceed as in [FK97] to obtain a reverse plane partition of height  $k$  from a  $k$ -flagged tableau. Let  $\lambda$  be a Ferrers shape and let  $\mu = D(\sigma_k(\lambda))$ . Since every entry in row  $i$  of a  $k$ -flagged tableau of shape  $\mu$  is less than or equal to  $i + k$  and greater than or equal to  $i$  (as the tableau is semistandard), one can subtract  $i$  from all the entries in row  $i$ , for all rows, and obtain a reverse plane partition of height  $k$  and shape  $\mu$ , or equivalently, a  $k$ -fan of noncrossing north-east paths inside  $\mu$ . To obtain a bijection between  $k$ -flagged





**Fig. 6:** Reverse plane partition corresponding to Figure 5 and its corresponding 2-fan of paths.

tableaux of shape  $\mu$  and the set  $\mathcal{F}_{SE}(\lambda, k)$  of  $k$ -south-east fillings of the shape  $\lambda$ , one lifts the  $i$ -th path from the bottom by  $i - 1$  and turns it into a path of  $+$ 's inside  $\lambda$ . See Figure 6 for an example; the red marks come from the red path, the blue from the blue path, and the additional black marks are contained in any 2-south-east filling.

Putting the described bijections together, we obtain Theorem 1.2.

**Theorem 3.2** *Let  $\lambda$  be a Ferrers shape. The composition of the described maps is a bijection between  $\mathcal{F}_{NE}(\lambda)$  and  $\mathcal{F}_{SE}(\lambda)$ .*

As mentioned in the introduction,  $k$ -triangulations of the  $n$ -gon can be seen as  $k$ -north-east fillings of the staircase shape  $(n - 1, \dots, 2, 1)$ , and  $k$ -fans of Dyck paths of length  $2(n - 2k)$  can be seen as  $k$ -south-east fillings of the same staircase (see e.g. [Kra06, Rub06]). Thus, we obtain Theorem 1.1. See Figure 7 for an example.

**Corollary 3.3** *In the case where  $\lambda$  is the staircase shape  $(n - 1, \dots, 2, 1)$ , the described map is a bijection between  $k$ -triangulations of the  $n$ -gon and  $k$ -fans of noncrossing Dyck paths of length  $2(n - 2k)$ .*

### 4 Properties of north-east fillings and $k$ -triangulations

Using Theorem 2.2, we obtain several properties of  $k$ -north-east fillings of moon polyominoes and of Ferrers shapes and  $k$ -triangulations in particular. Some of them were already known while others were only conjectured.

The first property was proved in the case of stack polyominoes by J. Jonsson in [Jon05, Theorem 10]. It follows immediately from Theorem 2.1.


**Corollary 4.1** *Every  $k$ -north-east filling of a moon polyomino  $M$  contains  $i$  many boxes where  $i$  equals the total number of boxes in  $M$  minus the length of  $\sigma_k(M)$ . In particular  $i$  equals the number of boxes in the first  $k$  rows and columns in the case of Ferrers shapes.*

The second property is part of the main theorem in [PS09, Theorem 1.4(i)] and concerns the star property as described as well in [Stu10]; for the notion used here, we refer as well to the latter.

**Corollary 4.2** *Every  $k$ -triangulation of the  $n$ -gon consists of exactly  $n - 2k$   $k$ -stars.*

Using the description of mutations for  $k$ -triangulations in Section 2.4, one can also describe the mutation of a facet in the simplicial complex

$$\Delta_{n,k} := \Delta(\mathbf{1}_k \times [n - 2k, \dots, 1]).$$



**Fig. 7:** An example of all the steps in the bijection: a 2-triangulation of the 8-gon, the 2-north-east filling of  $(7, \dots, 1)$ , the pipe dream for  $[1, 2, 6, 5, 4, 3]$ , the 2-flagged tableau of shape  $(3, 2, 1)$ , the 2-fan of Dyck paths of length 8, and finally the 2-south-east filling of  $(7, \dots, 1)$ .

This mutation corresponds to removing a diagonal in a  $k$ -triangulation and replacing it by the unique other diagonal which gives a  $k$ -triangulation. This operation is called *flip* in [PS09, Theorem 1.4(iii)].

**Corollary 4.3** *A facet  $F$  in the simplicial complex  $\Delta_{n,k}$  can be mutated at any vertex  $d = (i, j) \in F$  for which  $k < |i - j| < n - k$ .*

The next property of the constructed bijection is a refined counting of  $k$ -triangulations, as conjectured by C. Nicolas [Nic09].

**Theorem 4.4** *The number of  $k$ -triangulations of a convex  $n$ -gon having degree  $d$  in a given vertex is given by the determinantal expression*

$$\det \begin{pmatrix} \text{Cat}_{n-2k} & \cdots & \text{Cat}_{n-k-2} & B_{n-k-1}^k(d) \\ \vdots & \ddots & \vdots & \vdots \\ \text{Cat}_{n-k-1} & \cdots & \text{Cat}_{n-3} & B_{n-2}^k(d) \end{pmatrix},$$

where  $\text{Cat}_\ell$  is the usual Catalan number, and where  $B_\ell^k(d) = \frac{2k+d-3}{\ell} \binom{2\ell-2k-d+2}{\ell-1}$ .

### 4.1 Rotation of the $n$ -gon and a CSP for flagged tableaux

There is a natural cyclic action  $\rho$  on  $k$ -triangulations given by rotating the vertex labels in the  $n$ -gon counterclockwise. The following conjecture is due to V. Reiner [Rei09].

**Conjecture 4.5 (V. Reiner)** *Let  $\lambda$  be the staircase shape  $(n - 1, \dots, 2, 1)$  and let  $k$  be a positive integer. The triple*

$$\left( \mathcal{F}_{NE}(\lambda, k), \langle \rho \rangle, F(q) \right)$$

*exhibits the cyclic sieving phenomenon (CSP) as described in [RSW04].*

We can describe the cyclic action on  $k$ -triangulations induced by rotation in terms of the cyclic action on flagged tableaux as defined in Section 3.2.

**Theorem 4.6** *The constructed bijection maps the cyclic action on  $k$ -triangulations to the cyclic action given by flagged promotion on flagged tableaux.*

Using this connection, we obtain the following corollary.

**Corollary 4.7** *Conjecture 1.7 is equivalent to Conjecture 4.5.*

## 5 Schubert polynomials and geometry of Schubert varieties

In this section we use the results about moon polyominoes in Section 2.2 to obtain new properties of Schubert polynomials. It was shown in [FK96] that Schubert polynomials are a generating series for pipe dreams, more precisely, for a permutation  $\sigma$ ,

$$\mathfrak{S}_\sigma(x_1, \dots, x_n) = \sum_{D \in \mathcal{RP}(\sigma)} \prod_{(i,j) \in D} x_i.$$

We obtain the following theorem and thus Corollary 1.6.

**Theorem 5.1** *Let  $S$  be a stack polyomino, let  $\lambda$  be the associated Ferrers shape and let  $k$  be a positive integer. Then*

$$\mathfrak{S}_{\sigma_k(S)}(x_1, x_2, \dots) - \mathfrak{S}_{\sigma_k(\lambda)}(x_1, x_2, \dots)$$

*is monomial positive. In particular,  $\mathfrak{S}_{\sigma_k(S)}(1, 1, \dots)$  is greater than or equal to the number of  $k$ -flagged tableaux of shape  $\lambda$ .*

Let  $B$  be the subgroup of  $GL(n)$  consisting of upper triangular matrices. For each  $\sigma \in \mathcal{S}_n$  there is a subvariety  $X_\sigma$  of  $GL(n)/B$  known as the Schubert variety (see, e.g., [Man01, Chapter 3]). A. Knutson and E. Miller showed in [KM05] that the multiplicity of the point  $X_e$  at the Schubert variety  $X_\sigma$  is given by the specialization  $\mathfrak{S}_\sigma(1, 1, \dots)$ . Thus, from the previous theorem, one obtains the following corollary.

**Corollary 5.2** *The multiplicity of the point  $X_e$  at the Schubert variety  $X_{\sigma_k(S)}$  is greater than or equal to the multiplicity of the point  $X_e$  at the Schubert variety  $S_{\sigma_k(\lambda)}$ .*

## References

- [BB93] N. Bergeron and S. Billey, *RC-graphs and Schubert polynomials*, Experiment. Math. **2** (1993), no. 4, 257–269.
- [BJS93] S. Billey, W. Jockush, and R. Stanley, *Some combinatorial properties of Schubert polynomials*, J. Algebraic Combin. **2** (1993), 344–374.
- [EG87] P. Edelman and C. Greene, *Balanced tableaux*, Adv. Math. **63** (1987), no. 1, 42–99.

- [Eli07] S. Elizalde, *A bijection between 2-triangulations and pairs of non-crossing Dyck paths*, J. Combin. Theory Ser. A **114** (2007), no. 8, 1481–1503.
- [FK96] S. Fomin and A. Kirillov, *The Yang-Baxter equation, symmetric functions, and Schubert polynomials*, Discrete Math. **153** (1996), 123–143.
- [FK97] ———, *Reduced words and plane partitions*, J. Algebraic Combin. **6** (1997), no. 4, 311–319.
- [Jon05] J. Jonsson, *Generalized triangulations and diagonal-free subsets of stack polyominoes*, J. Comb. Theory, Ser. A **112** (2005), 117–142.
- [KM04] A. Knutson and E. Miller, *Subword complexes in Coxeter groups*, Adv. Math. **184** (2004), no. 1, 161–176.
- [KM05] ———, *Gröbner geometry of Schubert polynomials*, Ann. of Math. **161** (2005), no. 3, 1245–1318.
- [Kra06] C. Krattenthaler, *Growth diagrams, and increasing and decreasing chains in fillings of Ferrers shapes*, Adv. in Appl. Math. **37** (2006), 404–431.
- [Man01] L. Manivel, *Symmetric functions, Schubert polynomials and degeneracy loci*, SMF/AMS Texts and Monographs **6** (2001).
- [Nic09] C. Nicolás, *Another bijection between 2-triangulations and pairs of non-crossing Dyck paths*, in DMTCS as part of the FPSAC 2009 conference proceedings (2009).
- [PS09] V. Pilaud and F. Santos, *Multitriangulations as complexes of star polygons*, Discrete Comput. Geom. **41** (2009), no. 2, 284–317.
- [Rei09] V. Reiner, *personal communication*.
- [RSW04] V. Reiner, D. Stanton, and D. White, *The cyclic sieving phenomenon*, J. Combin. Theory Ser. A **108** (2004), 17–50.
- [Rub06] M. Rubey, *Increasing and decreasing sequences in fillings of moon polyominoes*, to appear in Adv. in Appl. Math., available at [arXiv:math/0604140](https://arxiv.org/abs/math/0604140) (2006).
- [Stu10] C. Stump, *A new perspective on  $k$ -triangulations*, J. Combin. Theory Ser. A **118** (2010), no. 6, 1794–1800.
- [Wac85] M. Wachs, *Flagged Schur functions, Schubert polynomials, and symmetrizing operators*, J. Combin. Theory Ser. A **40** (1985), no. 2, 276–289.