

On the quotients between the (revised) Szeged index and Wiener index of graphs

Huihui Zhang^{1,3*}Jing Chen^{2†}Shuchao Li^{3‡}¹ Department of Mathematics, Luoyang Normal Univeristy, Luoyang 471002, P.R. China² Department of Mathematics, Hunan First Normal University, Changsha, Hunan 410205, P.R. China³ Faculty of Mathematics and Statistics, Central China Normal University, Wuhan 430079, P.R. Chinareceived 26th June 2016, revised 6th Mar. 2017, accepted 14th Apr. 2017.

Let $Sz(G)$, $Sz^*(G)$ and $W(G)$ be the Szeged index, revised Szeged index and Wiener index of a graph G . In this paper, the graphs with the fourth, fifth, sixth and seventh largest Wiener indices among all unicyclic graphs of order $n \geq 10$ are characterized; and the graphs with the first, second, third, and fourth largest Wiener indices among all bicyclic graphs are identified. Based on these results, further relation on the quotients between the (revised) Szeged index and the Wiener index are studied. Sharp lower bound on $Sz(G)/W(G)$ is determined for all connected graphs each of which contains at least one non-complete block. In addition the connected graph with the second smallest value on $Sz^*(G)/W(G)$ is identified for G containing at least one cycle.

Keywords: Szeged index; Revised Szeged index; Wiener index

1 Introduction

We consider that all graphs in this paper are finite, undirected and simple. We follow the notations and terminologies in Bondy and Murty (2008) except otherwise stated. Let $G = (V_G, E_G)$ be a connected graph with vertex set V_G and edge set E_G . A connected graph is *cyclic* if it contains at least one cycle. In particular, a connected graph G is *unicyclic* (resp. *bicyclic*) if $|E_G| = |V_G|$ (resp. $|E_G| = |V_G| + 1$). For convenience, let $|G| := |V_G|$.

In the subsequent sections, we use $G - v$, or $G - uv$ to denote the graph obtained from G by deleting vertex $v \in V_G$, or edge $uv \in E_G$, respectively (it is naturally extended if at least two vertices or edges are deleted). Let $G + uv$ be the graph obtained from G by adding an edge $uv \notin E_G$. For a subset S of V_G , let $G[S]$ be the subgraph induced by S . For $v \in V_G$, we denote by $N_G(v)$ (or $N(v)$ for short) the set of all neighbors of v in G and let $d_G(v) = |N_G(v)|$ be the degree of v in G . Call u a *pendant vertex* or *leaf* in G , if $d_G(u) = 1$. We denote by P_n, C_n, S_n and K_n the path, cycle, star and complete graph of order n ,

*She is fully supported by the excellent doctoral dissertation cultivation grant from Central China Normal University (Grant No. 2016YBZZ084).

†She is fully supported by the National Natural Science Foundation of China (Grant No.11501188).

‡Corresponding author and I am fully supported by the National Natural Science Foundation of China (Grant No. 11671164).

respectively. We call $L_{n,r}$ a *lollipop* if it is obtained by identifying some vertex of C_r with an end-vertex of P_{n-r+1} .

For $u, v \in V_G$, the *distance* between u and v in G , denoted by $d_G(u, v)$, is the length of a shortest path connecting u and v . The *diameter*, $d(G)$, of G is equal to $\max_{u, v \in V_G} d_G(u, v)$. For all $v \in V_G$, let $n_G^i(v) = |\{u \in V_G : d(u, v) = i\}|$. The symbol \cong denotes that two graphs in question are isomorphic.

The *Wiener index* of G is defined as the sum of all distances between pairs of unordered vertices in G , i.e.,

$$W(G) = \sum_{\{u, v\} \subseteq V_G} d_G(u, v) = \frac{1}{2} \sum_{u \in V_G} D_G(u), \quad (1)$$

where $D_G(u) = \sum_{x \in V_G} d_G(x, u)$. This distance-based graph invariant was in chemistry introduced back in Wiener (1947) and in mathematics about 30 years later; Entringer et al. (1976). Nowadays, the Wiener index is an extensively studied graph invariant; see the reviews Dobrynin et al. (2001, 2002). A collection of recent papers dedicated to the investigations of the Wiener index; see Knor et al. (2016); Li and Song (2014); Liu et al. (2016). The problem of finding an upper bound on the Wiener index of a graph is quite challenging; see Mukwembi and Vetrík (2014). Only a few papers considered the upper bounds on the Wiener index of graphs; see Deng (2007); Zhang et al. (2010); Tang and Deng (2008); Dong and Zhou (2012).

Given an edge $e = uv$ in G , define three sets with respect to e as follows:

$$\begin{aligned} N_u(e) &= \{w \in V_G : d_G(u, w) < d_G(v, w)\}, & N_v(e) &= \{w \in V_G : d_G(v, w) < d_G(u, w)\}, \\ N_0(e) &= \{w \in V_G : d_G(u, w) = d_G(v, w)\}. \end{aligned}$$

Clearly, $V_G = N_u(e) \cup N_v(e) \cup N_0(e)$. For convenience, let $n_u(e) = |N_u(e)|$, $n_v(e) = |N_v(e)|$ and $n_0(e) = |N_0(e)|$. It is easy to see $n_u(e) + n_v(e) + n_0(e) = |V_G|$. If G is bipartite, then $N_0(e) = \emptyset$ holds for all $e \in E_G$. Consequently, for any bipartite graph G with $e \in E_G$, $n_u(e) + n_v(e) = |V_G|$.

From Dobrynin et al. (2001); Gutman and Polansky (1986); Wiener (1947), we know that, for a tree T , its Wiener index can be defined alternatively as

$$W(T) = \sum_{e=uv \in E_T} n_u(e)n_v(e). \quad (2)$$

Motivated by (2), Gutman (1994) introduced the *Szeged index* of graph G , which is defined by

$$Sz(G) = \sum_{e=uv \in E_G} n_u(e)n_v(e).$$

Randić (2002) observed that the vertices at equal distances from the end-vertices of an edge do not contribute to the Szeged index, and so he proposed the *revised Szeged index* $Sz^*(G)$ of a graph G as follows:

$$Sz^*(G) = \sum_{e=uv \in E_G} \left(n_u(e) + \frac{n_0(e)}{2} \right) \left(n_v(e) + \frac{n_0(e)}{2} \right).$$

For more recent results on (revised) Szeged index, one may be referred to these in Aouchiche and Hansen (2010); Ilić (2010); Klavžar and Nadjafi-Arani (2014); Li and Liu (2013); Pisanski and Randić (2010); Simić et al. (2000); Xing and Zhou (2011).

By (2) and (1), we know that $Sz(T) = W(T)$ holds for any tree T . Then, many researchers focused on the difference between the Szeged index and the Wiener index on general graphs. Given a graph G , the difference $Sz(G) - W(G) \geq 0$ holds, which was conjectured in Gutman (1994) and proved in Klavžar et al. (1996). Moreover, $Sz(G) = W(G)$ holds if and only if every block of G is a complete graph, which was obtained by Dobrynin and Gutman (1994), and see Khodashenas et al. (2011) for another proof. Nadjafi-Arani et al. (2011) studied the structure of graphs G with $Sz(G) - W(G) = k$, here k is a positive integer. In particular, Nadjafi-Arani et al. (2012) identified the graphs for which the difference is 4 and 5. The difference between $Sz(G)$ and $W(G)$ in networks was investigated in Klavžar and Nadjafi-Arani (2013). Pisanski and Randić (2010) showed that, if G is connected, then $Sz^*(G) \geq Sz(G)$ with equality if and only if G is bipartite. Some further results on the difference between the Wiener index and the (revised) Szeged index were established in Zhang et al. (2016).

The computer program AutoGraphiX was used to study the relationship involving graph invariants; see Aouchiche et al. (2005); Caporossi and Hansen (2000); Du and Ilić (2013) for more detailed information. Hansen et al. (2010) used the computer program AutoGraphiX to generate eight conjectures on the difference (resp. quotient) between the (revised) Szeged index and Wiener index. Chen et al. (2012, 2014) confirmed three conjectures on the difference between the (revised) Szeged index and Wiener index, which can be summarized as following three theorems.

Theorem 1.1 (Chen et al. (2012, 2014)). *Let G be a connected bipartite graph with $n \geq 4$ vertices and $|E_G| \geq n$ edges. Then $Sz(G) - W(G) \geq 4n - 8$. The equality holds if and only if G is composed of a cycle C_4 on 4 vertices and a tree T on $n - 3$ vertices sharing a single vertex.*

Theorem 1.2 (Chen et al. (2014)). *Let G be a connected graph with $n \geq 5$ vertices with an odd cycle and girth $g \geq 5$. Then $Sz(G) - W(G) \geq 2n - 5$. The equality holds if and only if G is composed of a cycle C_5 on 5 vertices, and one tree rooted at a vertex of the C_5 or two trees, respectively, rooted at two adjacent vertices of the C_5 .*

Theorem 1.3 (Chen et al. (2014)). *Let G be a connected graph with $n \geq 4$ vertices and $|E_G| \geq n$ edges and with an odd cycle. Then*

$$Sz^*(G) - W(G) \geq \frac{n^2 + 4n - 6}{4}.$$

The equality holds if and only if G is composed of a cycle C_3 on 3 vertices and a tree T on $n - 2$ vertices sharing a single vertex.

Recently, Li and Zhang (2017) confirmed three additional above conjectures, which are described as the following three theorems.

Theorem 1.4 (Li and Zhang (2017)). *Let G be a cyclic graph of order $n \geq 4$.*

(i) *If G is a bipartite graph, then*

$$\frac{Sz^*(G)}{W(G)} \geq 1 + \frac{24(n-2)}{n^3 - 13n + 36}$$

with equality if and only if G is the lollipop $L_{n,4}$.

(ii) *If G is a non-bipartite graph, then*

$$\frac{Sz^*(G)}{W(G)} \geq 1 + \frac{3(n^2 + 4n - 6)}{2(n^3 - 7n + 12)}$$

with equality if and only if G is the lollipop $L_{n,3}$.

Theorem 1.5 (Li and Zhang (2017)). *Let G be a unicyclic graph on $n \geq 4$ vertices. Then*

$$\frac{Sz(G)}{W(G)} \leq \begin{cases} 2 - \frac{8}{n^2+7}, & \text{if } n \text{ is odd,} \\ 2, & \text{if } n \text{ is even} \end{cases}$$

with equality if and only if G is the lollipop $L_{n,n-1}$ if n is odd and the cycle C_n if n is even.

Theorem 1.6 (Li and Zhang (2017)). *Let G be a unicyclic graph on $n \geq 4$ vertices. Then*

$$\frac{Sz^*(G)}{W(G)} \leq \begin{cases} 2 + \frac{2}{n^2-1}, & \text{if } n \text{ is odd,} \\ 2, & \text{if } n \text{ is even} \end{cases}$$

with equality if and only if G is the cycle C_n .

Li and Zhang (2017) determined sharp lower bounds on $Sz(G)/W(G)$ for cyclic graph with girth at least 4.

Theorem 1.7 (Li and Zhang (2017)). *Let G be a cyclic graph of order $n \geq 5$ with girth at least 4.*

(i) *If G is a bipartite graph, then*

$$\frac{Sz(G)}{W(G)} \geq 1 + \frac{24(n-2)}{n^3 - 13n + 36}$$

with equality if and only if G is the lollipop $L_{n,4}$.

(ii) *If G is a non-bipartite graph, then*

$$\frac{Sz(G)}{W(G)} \geq 1 + \frac{6(2n-5)}{n^3 - 25n + 90}$$

with equality if and only if G is the lollipop $L_{n,5}$.

Dobrynin and Gutman (1995) showed that $Sz(G)/W(G) \geq 1$ with equality if and only if every block of G is complete, whereas the case that G contains at least one block being non-complete is open. Based on Theorem 1.4, we may induce that $L_{n,4}$ is the unique graph with the smallest value on $Sz^*(G)/W(G)$ among n -vertex cyclic graphs. How can we determine the graph with second smallest value on $Sz^*(G)/W(G)$ among n -vertex cyclic graphs? Theorem 1.7 only considers the case for the graph with girth at least 4, whereas the case for cyclic graph of girth 3 is still open. In order to give solutions for the above open problems, it is natural and interesting for us to study some further relation on the quotients between the (revised) Szeged index and Wiener index of connected graphs.

Our paper is organized as follows: In Section 2, we give necessary definitions and state the main results of the paper. The first result determines the graphs with the fourth, fifth, sixth and seventh largest values on $W(G)$ among all unicyclic graphs of order $n \geq 10$; while the second result characterizes the graphs with the first four greatest values on $W(G)$ in the class of all bicyclic graphs. These two results will be proved in Sections 4 and 5, respectively. The third result obtains the sharp lower bound on $Sz(G)/W(G)$ for all connected graphs each of which contains at least one non-complete block. It, respectively, extends the results obtained by Dobrynin and Gutman (1995) and Li and Zhang (2017). The last result determines the cyclic graph G with the second smallest value on $Sz^*(G)/W(G)$, which extends the result obtained by Li and Zhang (2017). These two theorems are then proved in Section 6. In Section 3, we give some preliminary results which are used to prove our main results.

2 Main results

Consider a cycle C_r whose vertices are labeled consecutively by v_1, v_2, \dots, v_r . Then let $C_r(T_1, T_2, \dots, T_r)$ be an n -vertex graph obtained from C_r and rooted trees T_i 's by identifying the root, say r_i , of T_i with v_i on C_r , $i = 1, 2, \dots, r$. Assume that $|V_{T_i} \setminus \{r_i\}| = n_i$. Clearly, for $i = 1, 2, \dots, r$, $n_i \geq 0$. Thus, $|C_r(T_1, T_2, \dots, T_r)| = \sum_{i=1}^r n_i + r$. In particular, if every rooted tree is a path whose root is just one of its pendant vertices, then we denote $C_r(T_1, T_2, \dots, T_r)$ by $C_r(P_{n_1+1}, P_{n_2+1}, \dots, P_{n_r+1})$. For sim-




Fig. 1: Graphs H_n^0, H_n^1, H_n^2 and H_n^3 .

licity, let H_n^0, H_n^1, H_n^2 and H_n^3 be four unicyclic graphs of order n as depicted in Fig. 1. Then, let $\mathcal{A}_n = \{L_{n,3}, L_{n,4}, H_n^0, H_n^1, H_n^2, H_n^3, C_3(P_{n-3}, P_2, P_1), C_3(P_{n-4}, P_3, P_1), C_4(P_{n-4}, P_1, P_2, P_1)\}$.

Tang and Deng (2008) showed that, among the unicyclic graphs of order $n \geq 6$, the graph $L_{n,3}$ is the graph with the largest Wiener index, both $C_3(P_{n-3}, P_2, P_1)$ and $L_{n,4}$ are the graphs with the second largest Wiener index and H_n^0 is the graph with the third largest Wiener index. Our first main result in this paper characterizes the unicyclic graphs of order $n \geq 10$ with the fourth, fifth, sixth and seventh largest Wiener indices.

Theorem 2.1. *Let G be a unicyclic graph of order $n \geq 10$ and G is not in \mathcal{A}_n .*

- (i) *If $n = 10$ and $G \not\cong C_3(P_5, P_4, P_1)$, then $W(G) < W(H_{10}^3) = W(H_{10}^2) < W(H_{10}^1) = W(C_3(P_5, P_4, P_1)) < W(C_4(P_6, P_1, P_2, P_1)) < W(C_3(P_6, P_3, P_1))$.*
- (ii) *If $n = 11$ and $G \not\cong C_3(P_6, P_4, P_1)$, then $W(G) < W(H_{11}^3) = W(H_{11}^2) = W(C_3(P_6, P_4, P_1)) < W(H_{11}^1) < W(C_4(P_7, P_1, P_2, P_1)) < W(C_3(P_7, P_3, P_1))$.*
- (iii) *If $n \geq 12$, then $W(G) < W(H_n^3) = W(H_n^2) < W(H_n^1) < W(C_4(P_{n-4}, P_1, P_2, P_1)) < W(C_3(P_{n-4}, P_3, P_1))$.*

Let K_4^- be the graph obtained from K_4 by deleting one of its edges. Then, let $B_n^{(1)}$ be the n -vertex graph obtained by identifying an end-vertex of P_{n-3} with a 3-degree vertex in K_4^- . For $0 \leq s \leq \lfloor \frac{n-4}{2} \rfloor$, let $B(n, s)$ be the n -vertex graph obtained by attaching paths P_{n-s-4} and P_s , respectively, to two 2-degree vertices of K_4^- . Given two cycles C_p, C_q , let $B_n^{p,q}$ be the n -vertex graph obtained by joining C_p and C_q with a path of length $n - p - q + 1$. Our next main result characterizes all the n -vertex bicyclic graphs with the first four largest Wiener indices.

Theorem 2.2. *Let G be a bicyclic graph of order $n \geq 6$ and G is not in $\{B(n, 0), B(n, 1), B_n^{(1)}, B_n^{3,3}\}$.*

- (i) *If $n = 6$, then $W(G) < W(B_6^{(1)}) < W(B(6, 1)) = W(B_8^{3,3}) < W(B(6, 0))$.*
- (ii) *If $n = 8$ and $G \not\cong B(8, 2)$, then $W(G) < W(B(8, 2)) = W(B_8^{(1)}) < W(B(8, 1)) < W(B_8^{3,3}) < W(B(8, 0))$.*
- (iii) *If $n \geq 7$ and $n \neq 8$, then $W(G) < W(B_n^{(1)}) < W(B(n, 1)) < W(B_n^{3,3}) < W(B(n, 0))$.*




Fig. 2: Graphs G' and G'' .

Based on Theorem 2.2, our next main result determines the sharp lower bound on $Sz(G)/W(G)$ for all connected graphs each of which contains at least one non-complete block.

Theorem 2.3. *Let G be an n -vertex connected graph containing at least one non-complete block, $n \geq 5$. Then*

$$\frac{Sz(G)}{W(G)} \geq 1 + \frac{12}{n^3 - 19n + 54}$$

with equality if and only if $G \cong B_n^{(1)}$.

Based on Theorem 2.1, our last main result characterizes the connected cyclic graph G with the second smallest value on $Sz^*(G)/W(G)$.

Theorem 2.4. *Let $G (\not\cong L_{n,4})$ be a cyclic graph on $n \geq 10$ vertices. Then*

$$\frac{Sz^*(G)}{W(G)} \geq 1 + \frac{24(n-2)}{n^3 - 19n + 54}$$

with equality if and only if $G \cong H_n^2$, where H_n^2 is depicted in Fig. 1.

3 Preliminaries

In this section, we give some preliminary results and definitions which are used to prove our main results in the subsequent sections.

Lemma 3.1 (Xu and Das (2013)). *Let G be an n -vertex bicyclic graph of diameter $n - 2$. Then $G \in \{B(n, s) : 0 \leq s \leq \lfloor \frac{n-4}{2} \rfloor\}$.*

Lemma 3.2 (Zhang et al. (2016)). *Let G be an n -vertex connected graph of girth $r = 3$ and at least one block is non-complete. Then $Sz(G) - W(G) \geq 2$, the equality holds if and only if $G \cong G'$, where G' is depicted in Fig. 2 satisfying each block of H_1 (resp. H_2) being a complete graph.*

Tang and Deng (2008) showed that if G is an n -vertex unicyclic graph with $n \geq 6$, then $W(G) \leq W(L_{n,3})$. In fact, if $n = 4, 5$, it is routine to check that this result also holds. Hence,

Lemma 3.3. *Let G be a unicyclic graph on $n \geq 4$ vertices, then $W(G) \leq W(L_{n,3})$.*

For the unicyclic graph $C_r(T_1, T_2, \dots, T_r)$, if the non-trivial rooted trees are just T_i, T_j, \dots, T_k with $i, j, \dots, k \in \{1, 2, \dots, r\}$, then we denote it by $C_r^{i,j,\dots,k}(T_i, T_j, \dots, T_k)$. In particular, if every non-trivial tree $T_l, l = i, j, \dots, k$, is a star whose root is just its maximal degree vertex, then we denote it by $C_r^{i,j,\dots,k}(S_{n_i+1}, S_{n_j+1}, \dots, S_{n_k+1})$; if every non-trivial tree $T_l, l = i, j, \dots, k$, is a path whose root is just an end-vertex, then we denote it by $C_r^{i,j,\dots,k}(P_{n_i+1}, P_{n_j+1}, \dots, P_{n_k+1})$.

Lemma 3.4 (Tang and Deng (2008)). Let $\hat{G} = C_r^{i,j,\dots,k}(S_{n_i+1}, S_{n_j+1}, \dots, S_{n_k+1})$ and $\tilde{G} = C_r^{i,j,\dots,k}(P_{n_i+1}, P_{n_j+1}, \dots, P_{n_k+1})$. Then

$$W(\hat{G}) \leq W(G) \leq W(\tilde{G})$$

for any graph $G = C_r^{i,j,\dots,k}(T_i, T_j, \dots, T_k)$ with $|T_t| = n_t + 1, t = i, j, \dots, k$. The equality on the left (resp. right) holds if and only if $G \cong \hat{G}$ (resp. $G \cong \tilde{G}$).

Lemma 3.5 (Li and Zhang (2017)). Let G be an n -vertex unicyclic graph with girth $r \geq 5$. If r is odd, then $W(G) \leq (n^3 - 25n + 90)/6$ with equality if and only if $G \cong L_{n,5}$.

Let $T_n(n_1^{l_1}, n_2^{l_2}, \dots, n_k^{l_k})$ be an n -vertex tree obtained by identifying one end-vertex for each of l_1 paths of length n_1, l_2 paths of length n_2, \dots, l_k paths of length n_k . Clearly, $\sum_{i=1}^k l_i n_i = n - 1$. For simplicity, let $A_{n,k} := T_n(n - k - 1, 1^k)$. A leaf in $A_{n,k}$ is called a *unit leaf* if it is adjacent to the unique maximum degree vertex. Let $A_{n,k}^i$ be a graph obtained from $A_{n,k}$ by adding i edges to connect its unit leaves.

Lemma 3.6 (Deng (2007)). Let T be a tree of order $n \geq 9$. If $T \notin \{P_n, T_n(n - 3, 1^2), T_n(n - 4, 2, 1), T_n(n - 5, 3, 1), G''\}$, where G'' is depicted in Fig. 2. Then

$$W(T) < W(T_n(n - 5, 3, 1)) < W(G'') < W(T_n(n - 4, 2, 1)) < W(T_n(n - 3, 1^2)) < W(P_n).$$

By a direct calculation, we may obtain the Wiener index of each n -vertex trees for $5 \leq n \leq 8$ (based on Table 2 of the Appendix in Cvetković et al. (1979)). Together with Lemma 3.6, one obtains

Lemma 3.7. Let T be a tree of order $n \geq 5$. If $T \notin \{P_n, T_n(n - 3, 1^2)\}$, then $W(T) < W(T_n(n - 3, 1^2)) < W(P_n)$. Furthermore, if $n = 8$ and $T \notin \{P_8, T_8(5, 1^2), T_8(4, 2, 1), T_8(3^2, 1), G''\}$, then $W(T) < W(G'') < W(T_8(3^2, 1)) < W(T_8(4, 2, 1)) < W(T_8(5, 1^2)) < W(P_n)$, where G'' is the tree on 8 vertices as depicted in Fig. 2.

Lemma 3.8. Let G be a graph obtained from vertex-disjoint connected graphs G_1 and G_2 by identifying a vertex of G_1 with a vertex of G_2 and denote the common vertex by v . Then

$$W(G) = W(G_1) + W(G_2) + (|G_2| - 1)D_{G_1}(v) + (|G_1| - 1)D_{G_2}(v).$$

Proof: Note that $d_G(x, y) = d_{G_i}(x, y)$ for $\{x, y\} \subseteq V_{G_i}, i = 1, 2$. By (1), we have

$$\begin{aligned} W(G) &= \sum_{x,y \in V_{G_1}} d_G(x, y) + \sum_{x,y \in V_{G_2}} d_G(x, y) + \sum_{\substack{x \in V_{G_1} \setminus \{v\}, \\ y \in V_{G_2} \setminus \{v\}}} d_G(x, y) \\ &= \sum_{x,y \in V_{G_1}} d_{G_1}(x, y) + \sum_{x,y \in V_{G_2}} d_{G_2}(x, y) + \sum_{\substack{x \in V_{G_1} \setminus \{v\}, \\ y \in V_{G_2} \setminus \{v\}}} (d_{G_1}(x, v) + d_{G_2}(v, y)) \\ &= W(G_1) + W(G_2) + \sum_{\substack{x \in V_{G_1} \setminus \{v\}, \\ y \in V_{G_2} \setminus \{v\}}} d_{G_1}(x, v) + \sum_{\substack{x \in V_{G_1} \setminus \{v\}, \\ y \in V_{G_2} \setminus \{v\}}} d_{G_2}(y, v) \\ &= W(G_1) + W(G_2) + (|G_2| - 1) \sum_{x \in V_{G_1} \setminus \{v\}} d_{G_1}(x, v) + (|G_1| - 1) \sum_{y \in V_{G_2} \setminus \{v\}} d_{G_2}(y, v) \\ &= W(G_1) + W(G_2) + (|G_2| - 1)D_{G_1}(v) + (|G_1| - 1)D_{G_2}(v), \end{aligned}$$

as desired. \square

The following result is a direct consequence of the above lemma.

Corollary 3.9. *Let u be a pendant vertex of an n -vertex connected graph G and v be the unique neighbor of u . Then $W(G) = W(G - u) + D_{G-u}(v) + n - 1$.*

Lemma 3.10. *Given a connected graph H containing at least one edge, let G_1 be a graph obtained by identifying a vertex, say v , of H with a vertex in C_k , G_2 be a graph obtained by identifying the vertex v of H with a minimal degree vertex in $L_{k,3}$. Then one has $W(G_1) \leq W(G_2)$ with equality if and only if $k = 3$.*

Proof: Clearly, if $k = 3$, then $G_1 \cong G_2$. So we consider $k \geq 4$ in what follows. By Lemma 3.8, we have

$$\begin{aligned} W(G_1) &= W(H) + W(C_k) + (k-1)D_H(v) + (|H|-1)D_{C_k}(v), \\ W(G_2) &= W(H) + W(L_{k,3}) + (k-1)D_H(v) + (|H|-1)D_{L_{k,3}}(v). \end{aligned}$$

Thus,

$$W(G_1) - W(G_2) = W(C_k) - W(L_{k,3}) + (|H|-1)(D_{C_k}(v) - D_{L_{k,3}}(v)). \quad (3)$$

It is easy to see that $|H| \geq 2$. On the one hand, by Lemma 3.3, we have $W(C_k) \leq W(L_{k,3})$. On the other hand, $D_{C_k}(v) - D_{L_{k,3}}(v) \leq \frac{k^2}{4} - \frac{k^2-k-2}{2} = \frac{-k^2+2k+4}{4} < 0$ for $k \geq 4$. In view of (3), we have $W(G_1) - W(G_2) < 0$, i.e., $W(G_1) < W(G_2)$, as desired. \square

Lemma 3.11. *Let G be an n -vertex connected graph of diameter at most $n-3$. For all $v \in V_G$, one has $D_G(v) \leq (n^2 - n - 6)/2$ with equality if and only if $G \in \{A_{n,4}^i : i = 0, 1, 2, 3\}$ and v is a non-unit leaf in $A_{n,4}^i$, $0 \leq i \leq 3$.*

Proof: Let d be the diameter of G , then by the definition of $D_G(v)$ for all $v \in V_G$, we have

$$\begin{aligned} D_G(v) &= \sum_{x \in V_G} d_G(x, v) \\ &\leq (1 + 2 + \cdots + d) + d(n-d-1) \end{aligned} \quad (4)$$

$$\begin{aligned} &= -\frac{1}{2} \left[\left(d - \frac{2n-1}{2} \right)^2 - \frac{(2n-1)^2}{4} \right] \\ &\leq \frac{n^2 - n - 6}{2}. \quad (\text{Since } d \leq n-3) \end{aligned} \quad (5)$$

The equality in (4) holds if and only if there are exactly $n-d$ vertices each of which is of distance d from v , whereas the equality in (5) holds if and only if $d = n-3$. Hence, $D_G(v) = \frac{n^2-n-6}{2}$ if and only if $d = n-3$, v is a non-unit leaf in G and G contains at most 3 unit leaves, i.e., $G \in \{A_{n,4}^i : i = 0, 1, 2, 3\}$. \square

Lemma 3.12. *For the lollipop $L_{n,r}$ with $r < n$, if r is even then*

$$W(L_{n,r}) = \frac{1}{6} \left[n^3 + \left(-\frac{3}{2}r^2 + 3r - 1 \right) n + \left(\frac{5}{4}r^3 - 3r^2 + r \right) \right].$$

Proof: For convenience, let u be the unique vertex of degree 3 in $L_{n,r}$. By a direct calculation, we obtain that $W(C_r) = \frac{r^3}{8}$, $W(P_{n-r+1}) = \frac{(n-r+1)(n-r)(n-r+2)}{6}$, $D_{C_r}(u) = \frac{2W(C_r)}{r} = \frac{r^2}{4}$ and $D_{P_{n-r+1}}(u) = 1 + 2 + \dots + (n-r) = \frac{(n-r)(n-r+1)}{2}$. Note that u is a cut vertex of $L_{n,r}$. Hence, we obtain (based on Lemma 3.8)

$$\begin{aligned} W(L_{n,r}) &= W(C_r) + W(P_{n-r+1}) + (r-1)D_{P_{n-r+1}}(u) + (n-r)D_{C_r}(u) \\ &= \frac{r^3}{8} + \frac{(n-r+1)(n-r)(n-r+2)}{6} + \frac{(r-1)(n-r)(n-r+1)}{2} + \frac{r^2(n-r)}{4} \\ &= \frac{1}{6} \left[n^3 + \left(-\frac{3}{2}r^2 + 3r - 1 \right) n + \left(\frac{5}{4}r^3 - 3r^2 + r \right) \right], \end{aligned}$$

as desired. \square

Recall that $C_r(P_{n_1+1}, P_{n_2+1}, \dots, P_{n_r+1})$ is an n -vertex unicyclic graph obtained from $C_r = v_1v_2 \dots v_rv_1$ and paths P_{n_i+1} 's by identifying an end-vertex of P_{n_i+1} with v_i on the cycle C_r , $i = 1, 2, \dots, r$.

Lemma 3.13. *Let $G = C_r(P_{n_1+1}, \dots, P_{n_k+1}, \dots, P_{n_t+1}, \dots, P_{n_r+1})$ be an n -vertex unicyclic graph containing at least two non-trivial rooted paths, say P_{n_k+1} and P_{n_t+1} .*

(i) *If $\sum_{\substack{i=1 \\ i \neq k,t}}^r (n_i + 1)d_G(v_i, v_k) \leq \sum_{\substack{i=1 \\ i \neq k,t}}^r (n_i + 1)d_G(v_i, v_t)$, then*

$$W(G) < W(C_r(P_{n_1+1}, \dots, P_{n_{k-1}+1}, P_1, P_{n_{k+1}+1}, \dots, P_{n_{t-1}+1}, P_{n_k+n_t+1}, P_{n_{t+1}+1}, \dots, P_{n_r+1})).$$

(ii) *If $\sum_{\substack{i=1 \\ i \neq k,t}}^r (n_i + 1)d_G(v_i, v_k) > \sum_{\substack{i=1 \\ i \neq k,t}}^r (n_i + 1)d_G(v_i, v_t)$, then*

$$W(G) < W(C_r(P_{n_1+1}, \dots, P_{n_{k-1}+1}, P_{n_k+n_t+1}, P_{n_{k+1}+1}, \dots, P_{n_{t-1}+1}, P_1, P_{n_{t+1}+1}, \dots, P_{n_r+1})).$$

Proof: For convenience, denote by $P_{n_k+1} = u_1u_2 \dots u_{n_k+1}$ and $P_{n_t+1} = w_1w_2 \dots w_{n_t+1}$, where $u_1 = v_k, w_1 = v_t$.

(i) Let $G_1 = G - u_1u_2 + u_2w_{n_t+1}$, i.e.,

$$G_1 = C_r(P_{n_1+1}, \dots, P_{n_{k-1}+1}, P_1, P_{n_{k+1}+1}, \dots, P_{n_{t-1}+1}, P_{n_k+n_t+1}, P_{n_{t+1}+1}, \dots, P_{n_r+1}).$$

In what follows, we show that $W(G) < W(G_1)$.

Let $A = (V_G \setminus V_{P_{n_k+1}}) \cup \{u_1\}$. Clearly, $A = (V_{G_1} \setminus V_{P_{n_k+1}}) \cup \{u_1\}$. By the definition of $W(G)$, we have

$$\begin{aligned} W(G) &= W(P_{n_k+1}) + \sum_{x,y \in A} d_G(x,y) + \sum_{\substack{x \in V_{P_{n_k+1}} \setminus \{u_1\}, \\ y \in A \setminus \{u_1\}}} d_G(x,y) \\ &= W(P_{n_k+1}) + \sum_{x,y \in A} d_G(x,y) + \sum_{\substack{x \in V_{P_{n_k+1}} \setminus \{u_1\}, \\ y \in A \setminus \{u_1\}}} (d_G(x, v_k) + d_G(v_k, y)) \\ &= W(P_{n_k+1}) + \sum_{x,y \in A} d_G(x,y) + (n - n_k - 1) \sum_{x \in V_{P_{n_k+1}} \setminus \{u_1\}} d_G(x, v_k) + n_k \sum_{y \in A} d_G(v_k, y). \end{aligned}$$

Similarly, we have

$$\begin{aligned} W(G_1) &= W(P_{n_k+1}) + \sum_{x,y \in A} d_{G_1}(x,y) + (n - n_k - 1) \sum_{x \in V_{P_{n_k+1}} \setminus \{u_1\}} d_{G_1}(x, w_{n_t+1}) \\ &\quad + n_k \sum_{y \in A} d_{G_1}(w_{n_t+1}, y). \end{aligned}$$

Note that $d_G(x, y) = d_{G_1}(x, y)$, $d_G(w_{n_t+1}, y) = d_{G_1}(w_{n_t+1}, y)$ and $d_G(z, v_k) = d_{G_1}(z, w_{n_t+1})$ for all $x, y \in A, z \in V_{P_{n_k+1}} \setminus \{u_1\}$. Hence,

$$W(G) - W(G_1) = n_k \sum_{y \in A} (d_G(v_k, y) - d_G(w_{n_t+1}, y)). \quad (6)$$

In what follows, we show that the right of (6) is negative. By a direct calculation, we have

$$\begin{aligned} \sum_{y \in A} d_G(v_k, y) &= \sum_{\substack{i=1 \\ i \neq k}}^r \sum_{y \in V_{P_{n_i+1}}} (d_G(v_k, v_i) + d_G(v_i, y)) \\ &= (n_t + 1)d_G(v_k, v_t) + \sum_{\substack{i=1 \\ i \neq k, t}}^r (n_i + 1)d_G(v_k, v_i) + \sum_{\substack{i=1 \\ i \neq k}}^r D_{P_{n_i+1}}(v_i) \end{aligned}$$

and

$$\begin{aligned} \sum_{y \in A} d_G(w_{n_t+1}, y) &= n_t + d_G(v_k, v_t) + \sum_{\substack{i=1 \\ i \neq k, t}}^r \sum_{y \in V_{P_{n_i+1}}} (n_t + d_G(v_t, v_i) + d_G(v_i, y)) + D_{P_{n_t+1}}(v_t) \\ &= (n - n_k - n_t - 1)n_t + d_G(v_k, v_t) + \sum_{\substack{i=1 \\ i \neq k, t}}^r (n_i + 1)d_G(v_t, v_i) + \sum_{\substack{i=1 \\ i \neq k}}^r D_{P_{n_i+1}}(v_i). \end{aligned}$$

Bearing in mind the condition in (i) we have

$$\sum_{y \in A} d_G(v_k, y) - \sum_{y \in A} d_G(w_{n_t+1}, y) \leq n_t(d_G(v_k, v_t) - n + n_k + n_t + 1). \quad (7)$$

If $d_G(v_k, v_t) = 1$, then $d_G(v_k, v_t) - n + n_k + n_t + 1 = n_k + n_t + 2 - n = -\sum_{i \neq k, t} (n_i + 1) \leq -1$. Together with (6) and (7), we have $W(G) < W(G_1)$.

If $d_G(v_k, v_t) \geq 2$, then one may observe that $n - n_k - n_t - 2 = \sum_{i \neq k, t} (n_i + 1) \geq 2d_G(v_k, v_t) - 2$. Thus, $d_G(v_k, v_t) - n + n_k + n_t + 1 \leq 1 - d_G(v_k, v_t) \leq -1$. Thus, in view of (7) we have $\sum_{y \in A} d_G(v_k, y) < \sum_{y \in A} d_G(w_{n_t+1}, y)$. Together with (6), we obtain $W(G) < W(G_1)$, as desired.

(ii) By the same discussion as in the proof of (i), we may show that (ii) holds, which is omitted here. \square

Lemma 3.14. Let $C_r(T_1, T_2, \dots, T_r)$ be an n -vertex unicyclic graph with girth $r \geq 6$. If r is even, then

$$W(C_r(T_1, T_2, \dots, T_r)) \leq \frac{n^3 - 37n + 168}{6}$$

with equality if and only if $C_r(T_1, T_2, \dots, T_r) \cong L_{n,6}$ if $n \neq 8$ and $C_r(T_1, T_2, \dots, T_r) \cong L_{8,6}$ or $L_{8,8}$, otherwise.

Proof: Repeated using Lemmas 3.4 and 3.13 yields

$$\begin{aligned} W(C_r(T_1, T_2, \dots, T_r)) &\leq W(L_{n,r}) & (8) \\ &= \frac{1}{6} \left[n^3 + \left(-\frac{3}{2}r^2 + 3r - 1 \right) n + \left(\frac{5}{4}r^3 - 3r^2 + r \right) \right]. \quad (\text{By Lemma 3.12}) \end{aligned}$$

The equality in (8) holds if and only if $G \cong L_{n,r}$.

In order to complete the proof, it suffices to compare the Wiener index of $L_{n,6}$ with that of $L_{n,r}$ for $r \geq 8$. In fact, if $r \geq 8$ then by a direct calculation we have

$$\begin{aligned} W(L_{n,6}) - W(L_{n,r}) &= \frac{1}{6} \left[\left(\frac{3}{2}r^2 - 3r - 36 \right) n - \left(\frac{5}{4}r^3 - 3r^2 + r - 168 \right) \right] \\ &\geq \frac{1}{6} \left[\left(\frac{3}{2}r^2 - 3r - 36 \right) r - \left(\frac{5}{4}r^3 - 3r^2 + r - 168 \right) \right] & (9) \end{aligned}$$

$$\begin{aligned} &= \frac{1}{6} \left(\frac{1}{4}r^3 - 37r + 168 \right) \\ &\geq 0, & (10) \end{aligned}$$

where (9) follows by the fact that $n \geq r$ and $\frac{3}{2}r^2 - 3r - 36 > 0$ for $r \geq 8$; the equality in (9) holds if and only if $n = r$; whereas (10) follows by $r \geq 8$; the equality in (10) holds if and only if $r = 8$. Hence, if $r \geq 8$, then $W(L_{n,6}) = W(L_{n,r})$ holds if and only if $n = r = 8$. That is to say, $W(L_{8,6}) = W(L_{8,8})$.

By Lemma 3.12, one has $L_{n,6} = \frac{n^3 - 37n + 168}{6}$. Hence, together with (8)–(10), it follows that $W(C_r(T_1, T_2, \dots, T_r)) \leq \frac{n^3 - 37n + 168}{6}$. The equality holds if and only if $C_r(T_1, T_2, \dots, T_r) \cong L_{n,6}$ if $n \neq 8$, and $C_r(T_1, T_2, \dots, T_r) \cong L_{8,6}$ or $L_{8,8}$ if $n = 8$, as desired. \square

Lemma 3.15. Let G be a bicyclic graph of order n with diameter $n - 2$. Then

$$\frac{Sz(G)}{W(G)} \geq 1 + \frac{12(n-3)}{n^3 - 13n + 30}$$

with equality if and only if $G \cong B(n, 0)$.

Proof: Note that the diameter of G is $n - 2$, by Lemma 3.1 we obtain $G \cong B(n, s)$, where $0 \leq s \leq \lfloor \frac{n-4}{2} \rfloor$. By a direct calculation, one has

$$\begin{aligned} W(B(n, s)) &= W(P_{n-1}) + (1 + 2 + \dots + (s+1)) + (1 + 2 + \dots + (n-s-3)) + 1 \\ &= \frac{(n-2)(n-1)n}{6} + \frac{(s+1)(s+2)}{2} + \frac{(n-s-3)(n-s-2)}{2} + 1 & (11) \end{aligned}$$

and

$$\begin{aligned}
Sz(B(n, s)) &= ((n-1) + 2(n-2) + \cdots + s(n-s)) + 2(s+2)(n-s-3) + 2(s+1)(n-s+2) \\
&\quad + 1 + ((n-1) + 2(n-2) + (n-s-4)(s+4)) \\
&= \frac{ns(s+1)}{2} - \frac{s(s+1)(2s+1)}{6} + (4s+6)n - 4(s+2)^2 + 1 \\
&\quad + \frac{n(n-s-4)(n-s-3)}{2} - \frac{(n-s-4)(n-s-3)(2n-2s-7)}{6} \\
&= \frac{(n-1)n(n+1)}{6} + ns - s^2 - 4s - 1. \tag{12}
\end{aligned}$$

Note that $0 \leq s \leq \lfloor \frac{n-4}{2} \rfloor - 1$. Hence, $n \geq 2s+6$. Based on (11), one has $W(B(n, s+1)) - W(B(n, s)) = 2s+5-n < 0$, i.e., $W(B(n, s+1)) < W(B(n, s))$. By (12) one has $Sz(B(n, s+1)) - Sz(B(n, s)) = n - (2s+5) > 0$, i.e., $Sz(B(n, s+1)) > Sz(B(n, s))$. Thus, we have

$$\begin{aligned}
W(B(n, \lfloor \frac{n-4}{2} \rfloor)) &< W(B(n, \lfloor \frac{n-4}{2} \rfloor - 1)) < \cdots < W(B(n, 1)) < W(B(n, 0)), \tag{13} \\
Sz(B(n, 0)) &< Sz(B(n, 1)) < \cdots < Sz(B(n, \lfloor \frac{n-4}{2} \rfloor - 1)) < Sz(B(n, \lfloor \frac{n-4}{2} \rfloor)),
\end{aligned}$$

which implies that

$$\frac{Sz(B(n, 0))}{W(B(n, 0))} < \frac{Sz(B(n, 1))}{W(B(n, 1))} < \cdots < \frac{Sz(B(n, \lfloor \frac{n-4}{2} \rfloor - 1))}{W(B(n, \lfloor \frac{n-4}{2} \rfloor - 1))} < \frac{Sz(B(n, \lfloor \frac{n-4}{2} \rfloor))}{W(B(n, \lfloor \frac{n-4}{2} \rfloor))}.$$

Based on Eqs.(11)-(12), one has $W(B(n, 0)) = \frac{n^3-13n+30}{6}$ and $Sz(B(n, 0)) = \frac{n^3-n-6}{6}$. Hence, $\frac{Sz(G)}{W(G)} \geq \frac{Sz(B(n, 0))}{W(B(n, 0))} = \frac{n^3-n-6}{n^3-13n+30} = 1 + \frac{12(n-3)}{n^3-13n+30}$. The equality holds if and only if $G \cong B(n, 0)$. \square

4 Proof of Theorem 2.1

Let \mathcal{U}_n be the set of all unicyclic graph on $n \geq 10$ vertices. Recall that $\mathcal{A}_n = \{L_{n,3}, L_{n,4}, H_n^0, H_n^1, H_n^2, H_n^3, C_3(P_{n-3}, P_2, P_1), C_3(P_{n-4}, P_3, P_1), C_4(P_{n-4}, P_1, P_2, P_1)\}$. By a direct calculation, for $n \geq 10$, one has

$$\begin{aligned}
W(H_n^3) &= W(H_n^2) < W(H_n^1) < W(C_4(P_{n-4}, P_1, P_2, P_1)) < W(C_3(P_{n-4}, P_3, P_1)) < W(H_n^0) \\
&< W(C_3(P_{n-3}, P_2, P_1)) = W(L_{n,4}) < W(L_{n,3}). \tag{14}
\end{aligned}$$

In particular, $W(H_{10}^1) = W(C_3(P_5, P_4, P_1))$ and $W(H_{11}^2) = W(H_{11}^3) = W(C_3(P_6, P_4, P_1))$.

Note that $W(H_n^2) = W(H_n^3) = \frac{n^3-19n+54}{6}$. Hence, in order to complete the proof, it suffices to show that $W(G) < \frac{n^3-19n+54}{6}$ for G in $\mathcal{U}_n \setminus \mathcal{A}_n$ if $n \geq 12$ and for G in $\mathcal{U}_n \setminus (\mathcal{A}_n \cup \{C_3(P_{n-5}, P_4, P_1)\})$ if $n = 10, 11$.

By Lemmas 3.5 and 3.14, one has $W(G) < \frac{n^3-19n+54}{6}$ for all unicyclic graph G of girth $r \geq 5$. Hence, it suffices to consider $r = 3, 4$. Choose such an n -vertex unicyclic graph G of girth $r \leq 4$ in $\mathcal{U}_n \setminus \mathcal{A}_n$ if $n \geq 12$ and in $\mathcal{U}_n \setminus (\mathcal{A}_n \cup \{C_3(P_{n-5}, P_4, P_1)\})$ if $n = 10, 11$ such that $W(G)$ is as large as possible.

In what follows we show that G does not contain at least three non-trivial rooted trees. Otherwise, assume without loss of generality that T_1, T_k are the non-trivial rooted trees with the first two smallest sizes. Recall that for each rooted tree T_i in G , one has $|T_i| = n_i + 1$. Hence, by Lemma 3.4, we have $G \cong C_r(P_{n_1+1}, \dots, P_{n_k+1}, \dots, P_{n_r+1})$. By Lemma 3.13, we have $W(G) < W(H')$ if $\sum_{i \neq 1, k} (n_i + 1)d_G(v_i, v_1) \leq \sum_{i \neq 1, k} (n_i + 1)d_G(v_i, v_k)$ and $W(G) < W(H'')$ otherwise, where

$$\begin{aligned} H' &= C_r(P_1, \dots, P_{n_{k-1}+1}, P_{n_1+n_k+1}, P_{n_{k+1}+1}, \dots, P_{n_r+1}), \\ H'' &= C_r(P_{n_1+n_k+1}, \dots, P_{n_{k-1}+1}, P_1, P_{n_{k+1}+1}, \dots, P_{n_r+1}). \end{aligned}$$

If both H' and H'' are in $\mathcal{U}_n \setminus \mathcal{A}_n$ if $n \geq 12$ or in $\mathcal{U}_n \setminus (\mathcal{A}_n \cup \{C_3(P_{n-5}, P_4, P_1)\})$ if $n = 10, 11$, then we obtain a contradiction to the maximality of $W(G)$. Otherwise, if H' or $H'' \in \mathcal{A}_n$, as H' and H'' have at least two non-trivial rooted trees, then H' or $H'' \in \{C_3(P_{n-3}, P_2, P_1), C_3(P_{n-4}, P_3, P_1), C_4(P_{n-4}, P_1, P_2, P_1)\}$. Note that T_1, T_k are the non-trivial rooted trees with the first two smallest sizes and $n_1 + n_k \geq 2$, we have H' or H'' must be graph $C_3(P_{n-4}, P_3, P_1)$. Together with the fact $n \geq 10$, we have $G \cong C_3(P_{n-4}, P_2, P_2)$. By a simple computing, we have $W(G) = \frac{n^3 - 25n + 96}{6} < \frac{n^3 - 19n + 54}{6}$; if H' or $H'' \cong C_3(P_5, P_4, P_1)$, then G must be the graph $C_3(P_5, P_3, P_2), C_3(P_4, P_4, P_2)$ or $C_3(P_4, P_3, P_3)$. As $W(C_3(P_5, P_3, P_2)) = 135 < 144, W(C_3(P_4, P_4, P_2)) = 133 < 144$ and $W(C_3(P_4, P_3, P_3)) = 129 < 144$, we obtain that $W(G) < 144 = (10^3 - 190 + 54)/6$; if H' or $H'' \cong C_3(P_6, P_4, P_1)$, then G must be the graph $C_3(P_6, P_3, P_2)$ or $C_3(P_4, P_4, P_3)$. Since $W(C_3(P_6, P_3, P_2)) = 184 < 196$ and $W(C_3(P_4, P_4, P_3)) = 176 < 196$, we have $W(G) < 196 = (11^3 - 209 + 54)/6$, as desired.

Therefore, we obtain that G contains at most two non-trivial rooted trees. Bearing in mind that $r = 3$ or $r = 4$. Hence, we proceed by considering the following two possible cases.

Case 1. $r = 3$. In this case, it suffices to consider the following two subcases.

Subcase 1.1. G contains just one non-trivial rooted tree. Assume, without loss of generality, that T_1 is the non-trivial rooted tree. By Lemma 3.8, we have

$$W(G) = W(T_1) + 2D_{T_1}(v_1) + 2n - 3. \quad (15)$$

Note that $G \notin \{L_{n,3}, H_n^0, H_n^1\}$. Hence, we have $T_1 \notin \{P_{n-2}, T_{n-2}(n-5, 1^2), T_{n-2}(n-6, 2, 1)\}$. This implies that the diameter of T_1 is at most $n-4$.

If $d(T_1) = n-4$, as $T_1 \notin \{T_{n-2}(n-5, 1^2), T_{n-2}(n-6, 2, 1)\}$, then $\eta_{T_1}^{n-4}(v_1) \leq 1, \eta_{T_1}^{n-5}(v_1) \leq 1$. Thus,

$$\begin{aligned} D_{T_1}(v_1) &= \eta_{T_1}^1(v_1) + 2\eta_{T_1}^2(v_1) + \dots + (n-5)\eta_{T_1}^{n-5}(v_1) + (n-4)\eta_{T_1}^{n-4}(v_1) \\ &\leq [1 + 2 + \dots + (n-5) + (n-4)] + (n-6) \left(\sum_{i=1}^{n-4} \eta_{T_1}^i(v_1) - n + 4 \right) \\ &\leq [1 + 2 + \dots + (n-5) + (n-4)] + (n-6) \quad (\text{As } \sum_{i=1}^{n-4} \eta_{T_1}^i(v_1) = n-3) \\ &\leq \frac{n^2 - 5n}{2}. \end{aligned}$$

If $d(T_1) \leq n-5$, by Lemma 3.11, we have $D_{T_1}(v_1) \leq \frac{(n-2)^2 - (n-2) - 6}{2} = \frac{n^2 - 5n}{2}$.

Therefore, we obtain

$$2D_{T_1}(v_1) \leq n^2 - 5n.$$

If $|G| = 10$, then $|T_1| = 8$. Note that $T_1 \notin \{P_8, T_8(5, 1^2), T_8(4, 2, 1)\}$. By Lemma 3.7, we have $W(T_1) \leq W(T_8(3^2, 1)) = 75$. By (15) and (4), one has $W(G) \leq 75 + 50 + 17 = 142 < 144$. Hence, (i) holds in this subcase.

Now we consider that $|G| \geq 11$. If $T_1 \cong G''$ (G'' is depicted in Fig. 2), then $2D_{T_1}(v_1) \leq (n-5)(n-4) + 2(n-3) = n^2 - 7n + 14$. By a direct calculation (based on (15)), we have $W(G) \leq \frac{n^3 - 31n + 120}{6} < \frac{n^3 - 19n + 54}{6}$ for $n \geq 11$. If $T_1 \not\cong G''$, by Lemma 3.6 we have $W(T_1) \leq W(T_{n-2}(n-7, 3, 1)) = \frac{n^3 - 6n^2 - 7n + 120}{6}$. Thus, by (15), we have

$$\begin{aligned} W(G) &\leq \frac{n^3 - 6n^2 - 7n + 120}{6} + (n^2 - 5n) + 2n - 3 \\ &= \frac{n^3 - 25n + 102}{6} \\ &< \frac{n^3 - 19n + 54}{6}. \quad (\text{As } n \geq 11) \end{aligned}$$

Hence, (ii) and (iii) hold in this subcase.

Subcase 1.2. G contains just two non-trivial rooted trees, say T_1 and T_2 . Assume, without loss of generality, that $|T_1| \geq |T_2|$.

If $n_2 = 1$, then $n_1 = n - 4$. As $G \not\cong C_3(P_{n-3}, P_2, P_1)$, we obtain that $T_1 \not\cong P_{n-3}$. By Lemma 3.7, we have $W(T_1) \leq W(T_{n-3}(n-6, 1^2)) = \frac{(n-5)(n-4)}{6} + 2$ with equality if and only if $T_1 \cong T_{n-3}(n-6, 1^2)$. Observe that the diameter of T_1 is at most $n - 5$. Hence,

$$\begin{aligned} D_{T_1}(v_1) &= \eta_{T_1}^1(v_1) + 2\eta_{T_1}^2(v_1) + \cdots + (n-5)\eta_{T_1}^{n-5}(v_1) \\ &\leq [1 + 2 + 3 + \cdots + (n-5)] + \left(\sum_{i=1}^{n-5} \eta_{T_1}^i(v_1) - n + 5 \right) (n-5) \quad (16) \\ &= [1 + 2 + 3 + \cdots + (n-5)] + (n-5) \quad (\text{As } \sum_{i=1}^{n-5} \eta_{T_1}^i(v_1) = n-4) \\ &= \frac{(n-5)(n-2)}{2}, \quad (17) \end{aligned}$$

where the equality in (16) holds if and only if $\eta_{T_1}^1(v_1) = \cdots = \eta_{T_1}^{n-6}(v_1) = 1$ and $\eta_{T_1}^{n-5}(v_1) = 2$. This means that $T_1 \cong T_{n-3}(n-6, 1^2)$ and v_1 is a non-unit leaf in $T_{n-3}(n-6, 1^2)$. As v_1 is a cut vertex of G , we obtain (based on Lemma 3.8)

$$\begin{aligned} W(G) &= W(T_1) + W(L_{4,3}) + 3D_{T_1}(v_1) + (n-4)D_{L_{4,3}}(v_1) \\ &\leq \frac{(n-5)(n-4)}{6} + 2 + 8 + 3D_{T_1}(v_1) + 4(n-4) \quad (18) \end{aligned}$$

$$\begin{aligned} &\leq \frac{(n-5)(n-4)}{6} + 2 + 8 + \frac{3(n-5)(n-2)}{2} + 4(n-4) \quad (19) \\ &= \frac{n^3 - 19n + 54}{6}, \end{aligned}$$

where the equality in (18) holds if and only if $T_1 \cong T_{n-3}(n-6, 1^2)$; the equality in (19) holds if and only if $T_1 \cong T_{n-3}(n-6, 1^2)$ and v_1 is a non-unit leaf in $T_{n-3}(n-6, 1^2)$. Thus, $G \cong H_n^3$. As $G \notin \mathcal{A}_n$, we have $G \not\cong H_n^3$, which implies that $W(G) < \frac{n^3-19n+54}{6}$.

If $n_2 = 2$, then $n_1 = n - 5$ and $T_2 \cong P_3$. This implies that $d_G(v_2) = 3$ or 4 . If $d_G(v_2) = 3$, by a similar discussion as in the proof of $n_2 = 1$, we can obtain that $W(G) \leq \frac{n^3-25n+90}{6} < \frac{n^3-19n+54}{6}$ for $n \geq 10$. So we only consider the case $d_G(v_2) = 4$. Note that $|T_1| = n - 4$. By Lemma 3.7, we have $W(T_1) \leq W(P_{n-4}) = \frac{(n-4)(n-5)(n-3)}{6}$. It is easy to see that $D_{T_1}(v_1) \leq 1 + 2 + \dots + (n - 5) = \frac{(n-5)(n-4)}{2}$. As v_1 is a cut vertex of G , by Lemma 3.8 we have

$$\begin{aligned} W(G) &= W(T_1) + W(G[V_{T_2} \cup \{v_1, v_3\}]) + 4D_{T_1}(v_1) + (n-5)D_{G[V_{T_2} \cup \{v_1, v_3\}]}(v_1) \\ &\leq \frac{(n-4)(n-5)(n-3)}{6} + 15 + \frac{(n-5)(n-4)}{2} + 6(n-5) \\ &= \frac{n^3 - 25n + 90}{6} \\ &< \frac{n^3 - 19n + 54}{6}. \quad (\text{Since } n \geq 10) \end{aligned}$$

Now, we consider $n_2 \geq 3$. For $n = 10$, we obtain that $n_1 = 4, n_2 = 3$ directly. As $G \not\cong C_3(P_5, P_4, P_1)$, we have $T_1 \not\cong P_5$ or $T_2 \not\cong P_4$. If $T_1 \not\cong P_5$, then by Lemma 3.7, we have $W(T_1) \leq W(T_5(2, 1^2)) = 18$. Since $G[V_{T_2} \cup \{v_1, v_3\}]$ is a unicyclic graph on 6 vertices, by Lemma 3.3, we have $W(G[V_{T_2} \cup \{v_1, v_3\}]) \leq L_{6,3} = 31$. Observe that the diameters of T_1 and T_2 are at most 3, we can obtain that $D_{T_1}(v_1) \leq 1 + 2 + 3 + 3 = 9$ and $D_{G[V_{T_2} \cup \{v_1, v_3\}]}(v_1) \leq 1 + 1 + 2 + 3 + 4 = 11$. By Lemma 3.8, we have

$$\begin{aligned} W(G) &= W(T_1) + W(G[V_{T_2} \cup \{v_1, v_3\}]) + 3D_{T_1}(v_1) + 4D_{G[V_{T_2} \cup \{v_1, v_3\}]}(v_1) \\ &\leq 18 + 31 + 27 + 44 = 120 < 144. \end{aligned}$$

If $T_1 \cong P_5$, then one must have $T_2 \not\cong P_4$. Thus, $T_2 \cong S_4$ and $d_G(v_2) = 5$ or $d_G(v_2) = 3$. If $d_G(v_2) = 5$, then $W(G) = 126 < 144$; if $d_G(v_2) = 3$, then $W(G) = 138 < 144$. Hence, (i) holds in this subcase.

For $n \geq 11$, by Lemma 3.4, we have $G = C_3(P_{n_1+1}, P_{n_2+1}, P_1)$. Based on the structure of G and the fact $n_1 = n - 3 - n_2$, we have

$$\begin{aligned} W(G) &= W(P_{n_1+n_2+2}) + [1 + 2 + \dots + (n_1 + 1)] + [1 + 2 + \dots + (n_2 + 1)] \\ &= W(P_{n-1}) + [1 + 2 + \dots + (n - n_2 - 2)] + [1 + 2 + \dots + (n_2 + 1)] \\ &= \frac{(n-1)(n-2)n}{6} + \frac{(n-n_2-2)(n-n_2-1)}{2} + \frac{(n_2+1)(n_2+2)}{2} \\ &= n_2^2 - (n-3)n_2 + \frac{n^3 - 7n + 12}{6} \\ &\leq 9 - 3(n-3) + \frac{n^3 - 7n + 12}{6} \quad (\text{As } 3 \leq n_2 \leq \left\lfloor \frac{n-3}{2} \right\rfloor) \end{aligned} \tag{20}$$

$$\begin{aligned} &= \frac{n^3 - 25n + 120}{6} \\ &\leq \frac{n^3 - 19n + 54}{6}, \quad (\text{As } n \geq 11) \end{aligned} \tag{21}$$

where the equality in (20) holds if and only if $n_2 = 3$; whereas the equality in (21) holds if and only if $n = 11$. That is, $G \cong C_3(P_6, P_4, P_1)$. As $G \not\cong C_3(P_6, P_4, P_1)$ for $n = 11$, we have $W(G) < \frac{n^3-19n+54}{6}$. Hence, (ii) and (iii) hold in this subcase.

Hence, by Subcases 1.1 and 1.2 we obtain that (i), (ii) and (iii) hold for Case 1.

Case 2. $r = 4$. In this case, we first consider that G contains just one non-trivial rooted tree, say T_1 . By Lemma 3.8, we have

$$W(G) = W(T_1) + 3D_{T_1}(v_1) + 4n - 8. \quad (22)$$

Note that $G \not\cong L_{n,4}$, we have $T_1 \not\cong P_{n-3}$. Bearing in mind that $n \geq 10$, together with Lemma 3.7 for $n-3 (\geq 7)$, it follows that $W(T_1) \leq W(T_{n-3}(n-6, 1^2)) = \frac{n^3-9n^2+20n+12}{6}$. By (17), we have $D_{T_1}(v_1) \leq \frac{(n-5)(n-2)}{2}$ with equality if and only if $T_1 \cong T_{n-3}(n-6, 1^2)$ and v_1 is a non-unit leaf in $T_{n-3}(n-6, 1^2)$. Thus, by (22), we have

$$W(G) \leq \frac{n^3 - 9n^2 + 20n + 12}{6} + 3D_{T_1}(v_1) + 4n - 8 \quad (23)$$

$$\leq \frac{n^3 - 9n^2 + 20n + 12}{6} + \frac{3(n-5)(n-2)}{2} + 4n - 8 \quad (24)$$

$$= \frac{n^3 - 19n + 54}{6}.$$

By Lemma 3.7, the equality in (23) holds if and only if $T_1 \cong T_{n-3}(n-6, 1^2)$; whereas the equality in (24) holds if and only if $T_1 \cong T_{n-3}(n-6, 1^2)$ and v_1 is a non-unit leaf in $T_{n-3}(n-6, 1^2)$. Hence, $G \cong H_n^2$. As $G \notin \mathcal{A}_n$, we have $G \not\cong H_n^2$, which implies that $W(G) < \frac{n^3-19n+54}{6}$.

Now we consider the remaining case that G contains just two non-trivial rooted trees. By Lemma 3.4, up to isomorphism one has $G = C_4(P_{n_1+1}, P_{n_2+1}, P_1, P_1)$ with $0 \leq n_1 \leq n_2$, or $G = C_4(P_{n_1+1}, P_1, P_{n_3+1}, P_1)$ with $0 \leq n_1 \leq n_3$. For the former case, by a direct calculation, one has $W(G) = 2n_1^2 - 2(n-4)n_1 + \frac{n^3-13n+36}{6} \leq 2 - 2(n-4) + \frac{n^3-13n+36}{6} = \frac{n^3-25n+96}{6} < \frac{n^3-19n+54}{6}$ for $n \geq 10$. For the latter case, bearing in mind that $n_1 \geq 2$, one has $W(G) = n_1^2 - (n-4)n_1 + \frac{n^3-13n+36}{6} \leq 2 - 2(n-4) + \frac{n^3-13n+36}{6} = \frac{n^3-25n+96}{6} < \frac{n^3-19n+54}{6}$ for $n \geq 10$. Here, (i), (ii) and (iii) are true for Case 2.

This completes the proof. \square

5 Proof of Theorem 2.2

In this section, we give the proof of Theorem 2.2. For convenience, let $B_{k,l,t}$ be the bicyclic graph of order n obtained from paths $P_{k+1}, P_{l+1}, P_{t+1}$ by identifying all the left (resp. right) end-vertices as a new vertex. One often calls $B_{k,l,t}$ the θ -graph.

Proof of Theorem 2.2. Let \mathcal{B}_n be the set of all bicyclic graph on $n \geq 6$ vertices. By a direct calculation, for $n \geq 7$, one has

$$W(B_n^{(1)}) < W(B(n, 1)) < W(B_n^{3,3}) < W(B(n, 0)).$$

In particular, $W(B(8, 2)) = W(B_8^{(1)})$ and $W(B_6^{(1)}) < W(B(6, 1)) = W(B_6^{3,3}) < W(B(6, 0))$.




Fig. 3: Graphs F_i used in the proof of Theorem 2.2, $1 \leq i \leq 11$.

It is routine to check that $W(B_n^{(1)}) = (n^3 - 19n + 54)/6$. Hence, in order to complete the proof, it suffices to show that $W(G) < (n^3 - 19n + 54)/6$ for all G in $\mathcal{B}_n \setminus \{B(n, 0), B(n, 1), B_n^{(1)}, B_n^{3,3}\}$ if $n \neq 8$, and for G in $\mathcal{B}_8 \setminus \{B(8, 0), B(8, 1), B_8^{(1)}, B_8^{3,3}, B(8, 2)\}$ if $n = 8$.

Note that the diameter of G is at most $n - 2$. If $d(G) = n - 2$, then $G \cong B(n, s)$ for some $0 \leq s \leq \lfloor \frac{n-4}{2} \rfloor$. As $G \notin \{B(n, 0), B(n, 1)\}$, we have $s \geq 2$, which implies that $n \geq 8$. Note that $G \not\cong B(8, 2)$ for $n = 8$. Thus, $n \geq 9$. Combining with (11) and (13), we obtain $W(G) \leq W(B(n, 2)) = \frac{n^3 - 25n + 102}{6} < \frac{n^3 - 19n + 54}{6}$ for $n \geq 9$. Hence, (ii) and (iii) hold for the case $d(G) = n - 2$.

If $d(G) \leq n - 3$, then we show our result by induction on n . When $n = 6$, we have $W(B_6^{(1)}) = 26$ and with the help of Nauty based on McKay and Piperno (2014), we obtain $\mathcal{B}_6 \setminus \{B(6, 0), B(6, 1), B_6^{(1)}, B_6^{3,3}\} = \{B_6^{3,4}, B_{1,2,4}, B_{1,3,3}, B_{2,2,3}, F_1, F_2, \dots, F_{11}\}$, where F_1, F_2, \dots, F_{11} are depicted in Fig. 3. By a direct calculation, one has

$$\begin{aligned} W(F_1) &= 25, & W(F_4) &= 25, & W(F_7) &= 24, & W(F_{10}) &= 24, & W(B_{1,2,4}) &= 24, \\ W(F_2) &= 23, & W(F_5) &= 24, & W(F_8) &= 25, & W(F_{11}) &= 25, & W(B_{1,3,3}) &= 25, \\ W(F_3) &= 23, & W(F_6) &= 25, & W(F_9) &= 25, & W(B_6^{3,4}) &= 25, & W(B_{2,2,3}) &= 23. \end{aligned}$$

Thus, $W(G) \leq 25 < 26 = W(B_6^{(1)})$. That is, (i) is true for $n = 6$.

Now, consider $n \geq 7$. Assume that our result is true for $n - 1$. In order to complete the proof, it suffices to consider the following two possible cases.

Case 1. G contains pendant vertices. Choose a pendant vertex, say u , in G and let v be its unique neighbor. Clearly, $G - u$ is a bicyclic graph on $n - 1$ vertices. By Corollary 3.9, we have

$$W(G) = W(G - u) + D_{G-u}(v) + n - 1 \tag{25}$$

Note that the diameter $d(G) \leq n - 3$. Hence, $d(G - u) \leq n - 3$.

If $d(G - u) = n - 3$, then by Lemma 3.1, we have $G - u \cong B(n - 1, s)$ with $0 \leq s \leq \lfloor \frac{n-5}{2} \rfloor$. Combining with (13), we obtain $W(G - u) \leq W(B(n - 1, 0)) = \frac{n^3 - 3n^2 - 10n + 42}{6}$. Observe that v isn't a pendant vertex of $G - u$. Otherwise, $d(G) = n - 2$, a contradiction. Hence, $D_{G-u}(v) \leq 1 + 2 + 3 + \dots + (n - 4) + 1 + (n - 5) = \frac{n^2 - 5n + 4}{2}$. By (25), we have

$$\begin{aligned} W(G) &\leq \frac{n^3 - 3n^2 - 10n + 42}{6} + \frac{n^2 - 5n + 4}{2} + n - 1 \\ &= \frac{n^3 - 19n + 48}{6} \\ &< \frac{n^3 - 19n + 54}{6}. \end{aligned}$$

Now, consider that $d(G - u) \leq n - 4$. If $G - u \cong B_{n-1}^{3,3}$, then by a direct calculation, we have $W(G - u) = \frac{n^3 - 3n^2 - 10n + 36}{6}$. Based on the structure of $B_{n-1}^{3,3}$, we obtain $D_{G-u}(v) \leq 1 + 2 + \cdots + (n - 4) + 1 + (n - 4) = \frac{n^2 - 5n + 6}{2}$. By (25), one has $W(G) \leq \frac{n^3 - 3n^2 - 10n + 36}{6} + \frac{n^2 - 5n + 6}{2} + n - 1 = \frac{n^3 - 19n + 48}{6} < \frac{n^3 - 19n + 54}{6}$, the result holds. Note that $d(G - u) \leq n - 4$, by Lemma 3.11, we have

$$D_{G-u}(v) \leq \frac{(n-1)^2 - (n-1) - 6}{2} = \frac{n^2 - 3n - 4}{2}, \quad (26)$$

the equality in (26) if and only if $G - u \cong B_{n-1}^{(1)}$ and v is a non-unit leaf in $B_{n-1}^{(1)}$.

If $G - u \cong B_{n-1}^{(1)}$, then $W(G - u) = \frac{n^3 - 3n^2 - 16n + 72}{6}$. By (25) and (26), one has

$$\begin{aligned} W(G) &\leq \frac{n^3 - 3n^2 - 16n + 72}{6} + \frac{n^2 - 3n - 4}{2} + n - 1 \\ &= \frac{n^3 - 19n + 54}{6}. \end{aligned} \quad (27)$$

By (26), the equality in (27) holds if and only if $G - u \cong B_{n-1}^{(1)}$ and v is a non-unit leaf in $B_{n-1}^{(1)}$. That is to say, $W(G) = \frac{n^3 - 19n + 54}{6}$ holds if and only if $G \cong B_n^{(1)}$, which is impossible. Hence, $W(G) < \frac{n^3 - 19n + 54}{6}$.

Thus, it is left for us to consider $G - u \notin \{B_{n-1}^{3,3}, B_{n-1}^{(1)}\}$. By the induction hypothesis, we have $W(G - u) < \frac{(n-1)^3 - 19(n-1) + 54}{6} = \frac{n^3 - 3n^2 - 16n + 72}{6}$. Combining with (25) and (26), we obtain $W(G) < \frac{n^3 - 3n^2 - 16n + 72}{6} + \frac{n^2 - 3n - 4}{2} + n - 1 = \frac{n^3 - 19n + 54}{6}$, as desired. Therefore, (ii) and (iii) hold for Case 1.

Case 2. G does not contain pendant vertices. In this case, G is either the graph $B_n^{p,q}$ or the θ -graph $B_{k,l,t}$. So we proceed by distinguishing the following two possible subcases.

Subcase 2.1. $G = B_n^{p,q}$. Assume, without loss of generality, that $p \leq q$. Note that $G \not\cong B_n^{3,3}$, we have $q \geq 4$. As $p \geq 3$, by Lemma 3.10, we have $W(G) \leq W(B_n^{3,q})$. In what follows, we show that $W(B_n^{3,q}) < \frac{n^3 - 19n + 54}{6}$.

In fact, by Lemma 3.8, one has

$$W(B_n^{3,q}) = \begin{cases} W(L_{n-2,q}) + n^2 - 3n + 3 + \frac{-q^2 + 2q}{2}, & \text{if } q \text{ is even;} \\ W(L_{n-2,q}) + n^2 - 3n + 3 + \frac{-q^2 + 2q - 1}{2}, & \text{if } q \text{ is odd} \end{cases} \quad (28)$$

and using Lemma 3.12 for $n - 2$ and $r = 4$, one has

$$W(L_{n-2,4}) = \frac{(n-2)^2 - 13(n-2) + 36}{6} = \frac{n^3 - 6n^2 - n + 54}{6}. \quad (29)$$

If $n = 7$, then $W(L_{5,q}) \leq \max\{W(L_{5,4}), W(C_5)\} = 16$. By (28), we have $W(B_7^{3,q}) \leq 16 + 7^2 - 21 + 3 - 4 = 43 < 44 = \frac{n^3 - 19n + 54}{6}$, as desired. Now consider $n \geq 8$. Note that $W(L_{n-2,5}) = \frac{n^3 - 6n^2 - 13n + 132}{6} < \frac{n^3 - 6n^2 - n + 54}{6}$ and $W(L_{n-2,6}) = \frac{n^3 - 6n^2 - 25n + 234}{6} < \frac{n^3 - 6n^2 - n + 54}{6}$. Combining

with (28)-(29), we obtain

$$\begin{aligned}
W(B_n^{3,q}) &\leq W(L_{n-2,q}) + n^2 - 3n + 3 - 4 && \text{(Since } q \geq 4\text{)} \\
&\leq W(L_{n-2,4}) + n^2 - 3n - 1 && \text{(By Lemmas 3.5 and 3.14)} \\
&= \frac{n^3 - 19n + 48}{6} && \text{(By Eq.(29))} \\
&< \frac{n^3 - 19n + 54}{6}.
\end{aligned}$$

Hence, (ii) and (iii) hold for Subcase 2.1.

Subcase 2.2. $G = B_{k,l,t}$. In this subcase, $k + l + t = n + 1$. Assume, without loss of generality, that $k \leq l \leq t$ and G contains just two vertices, say u_1, u_2 , of degree 3. Clearly, each of the rest vertices is of degree 2.

Recall that $\eta_G^i(x) = |\{u | d_G(u, x) = i\}|$ for all $x \in V_G$. As $d(G) \leq \lfloor \frac{n}{2} \rfloor$, we have $\eta_G^i(x) = 0$ for $i > \lfloor \frac{n}{2} \rfloor$. In what follows, we show that if $\eta_G^{i+1}(x) > 0$, then $\eta_G^i(x) \geq 2$.

In fact, if $\eta_G^i(x) < 2$, then $\eta_G^i(x) = 1$. This implies that there exists a unique vertex u such that $d(u, x) = i$. Since $\eta_G^{i+1}(x) > 0$, we obtain that u is a cut vertex of G , a contradiction to the fact that $B_{k,l,t}$ contains no cut vertex. Hence, $\eta_G^i(x) \geq 2$.

Now we show that our result holds for even n . Similarly, we can also show that our result holds for odd n , which is omitted here.

Observe that, for all $x \in V_G \setminus \{u_1, u_2\}$, we have $\eta_G^{\frac{n}{2}}(x) \leq 1$. Otherwise, G has at least $n + 1$ vertices, a contradiction. Similarly, we have $\eta_G^{\frac{n}{2}}(u_1) = \eta_G^{\frac{n}{2}}(u_2) = 0$, $\eta_G^{\frac{n-2}{2}}(u_1) \leq 2$ and $\eta_G^{\frac{n-2}{2}}(u_2) \leq 2$. Thus, for all $x \in V_G \setminus \{u_1, u_2\}$, one has

$$D_G(x) = \sum_{u \in V_G} d_G(u, x) = \sum_{1 \leq i \leq \frac{n}{2}} i \eta_G^i(x) \leq 2 \left(1 + 2 + \cdots + \frac{n-2}{2} \right) + \frac{n}{2} = \frac{n^2}{4}. \quad (30)$$

For $y \in \{u_1, u_2\}$, one has

$$D_G(y) = \sum_{u \in V_G} d_G(u, y) = \sum_{1 \leq i \leq \frac{n-2}{2}} i \eta_G^i(y) \leq 3 + 2 \left(2 + \cdots + \frac{n-2}{2} \right) = \frac{n^2 - 2n + 4}{4}. \quad (31)$$

Together with (1) and (30)-(31), it follows that

$$\begin{aligned}
W(G) &= \frac{1}{2} \sum_{x \neq u_1, u_2} D_G(x) + \frac{1}{2} D_G(u_1) + \frac{1}{2} D_G(u_2) \\
&\leq \frac{n-2}{2} \cdot \frac{n^2}{4} + \frac{n^2 - 2n + 4}{4} \\
&= \frac{n^3 - 4n + 8}{8} \\
&< \frac{n^3 - 19n + 54}{6}. \quad \text{(As } n \geq 7\text{)}
\end{aligned}$$

Hence, (ii) and (iii) hold for Subcase 2.2.

This completes the proof. \square

6 Proofs of Theorems 2.3 and 2.4

In this section, we give the proofs of Theorems 2.3 and 2.4.




Fig. 4: Graphs H_i used in the proof of Theorem 2.3, $1 \leq i \leq 9$.

Proof of Theorem 2.3. If $n = 5$, then with the help of Nauty based on McKay and Piperno (2014) all of such graphs are $C_5, L_{5,4}, B_5^{(1)}, H_1, H_2, \dots, H_9$, where H_1, H_2, \dots, H_9 are depicted in Fig. 4. By a direct calculation, we have

$$\begin{aligned} \frac{Sz(B_5^{(1)})}{W(B_5^{(1)})} &= \frac{8}{7}, & \frac{Sz(C_5)}{W(C_5)} &= \frac{4}{3}, & \frac{Sz(L_{5,4})}{W(L_{5,4})} &= \frac{7}{4}, & \frac{Sz(H_1)}{W(H_1)} &= \frac{19}{15}, & \frac{Sz(H_2)}{W(H_2)} &= \frac{18}{7}, & \frac{Sz(H_3)}{W(H_3)} &= \frac{12}{7}, \\ \frac{Sz(H_4)}{W(H_4)} &= \frac{18}{13}, & \frac{Sz(H_5)}{W(H_5)} &= \frac{29}{13}, & \frac{Sz(H_6)}{W(H_6)} &= \frac{19}{13}, & \frac{Sz(H_7)}{W(H_7)} &= \frac{4}{3}, & \frac{Sz(H_8)}{W(H_8)} &= 2, & \frac{Sz(H_9)}{W(H_9)} &= \frac{15}{11}. \end{aligned}$$

Thus, $B_5^{(1)}$ is the unique extremal graph having minimum value on $\frac{Sz(G)}{W(G)}$. Our result holds for $n = 5$.

Now, we assume that $n \geq 6$. Note that at least one block of G is non-complete. Hence, G contains at least one cycle and its diameter $d(G) \leq n - 2$. For convenience, let r be the girth of graph G . Then if $r \geq 4$, by Theorem 1.7, we have

$$\begin{aligned} \frac{Sz(G)}{W(G)} &\geq \min \left\{ 1 + \frac{24(n-2)}{n^3 - 13n + 36}, 1 + \frac{6(2n-5)}{n^3 - 25n + 90} \right\} \\ &= 1 + \frac{6(2n-5)}{n^3 - 25n + 90} > 1 + \frac{12}{n^3 - 19n + 54}. \end{aligned} \quad (32)$$

So, in what follows, it suffices to consider graph G of girth $r = 3$. In order to complete our proof, we distinguish the following two cases $d(G) = n - 2$ and $d(G) \leq n - 3$.

If $d(G) = n - 2$, then there exists an induced path P_{n-1} in G . Note that $r = 3$ and at least one block is non-complete, we obtain that G contains only one vertex in $V_G \setminus V_{P_{n-1}}$ such that it must be adjacent to just three vertices of P_{n-1} . Thus, G is a bicyclic graph. By Lemma 3.15, we have

$$\frac{Sz(G)}{W(G)} \geq 1 + \frac{12(n-3)}{n^3 - 13n + 30} > 1 + \frac{12}{n^3 - 19n + 54}. \quad (33)$$

If $d(G) \leq n - 3$, then $G \not\cong B_n^{3,3}$ (based on the fact that G contains at least one non-complete block). Note that $W(G) < W(G - e)$ for any non-cut edge $e \in E_G$. Thus, we may repeatedly delete the non-cut edges of G until the resulting graph is a bicyclic graph G_1 such that $W(G) \leq W(G_1)$. If $G_1 \cong B(n, s)$ for some $0 \leq s \leq \lfloor \frac{n-4}{2} \rfloor$, as $G \not\cong B(n, s)$, then there exist some edges $e_1, e_2, \dots, e_k, k \geq 1$, such that




Fig. 5: Graphs (a), (b), (c), (d) used in the proofs of Theorems 2.3 and 2.4.

$G = G_1 + \{e_1, e_2, \dots, e_k\}$, where G_1 is depicted in Fig. 5(a). Now we construct a new bicyclic graph G_2 as follows:

$$G_2 = G_1 - u_1v_1 + e_1.$$

Clearly, G_2 is not in $\{B_n^{3,3}\} \cup \{B(n, s) : s = 0, 1, \dots, \lfloor \frac{n-4}{2} \rfloor\}$. It is routine to check that $G = G_2 + \{u_1v_1, e_2, \dots, e_k\}$. Thus, $W(G) < W(G_2)$.

If $G_1 \cong B_n^{3,3}$, as $G \not\cong B_n^{3,3}$, then there exist some edges e_1, e_2, \dots, e_t , $t \geq 1$, such that $G = G_1 + \{e_1, e_2, \dots, e_t\}$, where G_1 is depicted in Fig. 5(b). Now we construct a new bicyclic graph G_3 as follows:

$$G_3 = G_1 - u_2v_2 + e_1.$$

Clearly, G_3 is not in $\{B_n^{3,3}\} \cup \{B(n, s) : s = 0, 1, \dots, \lfloor \frac{n-4}{2} \rfloor\}$. It is routine to check that $G = G_3 + \{u_2v_2, e_2, \dots, e_t\}$. Thus, $W(G) < W(G_3)$.

Thus, we may assume that G_1 is not in $\{B_n^{3,3}\} \cup \{B(n, s) : s = 0, 1, \dots, \lfloor \frac{n-4}{2} \rfloor\}$. By Theorem 2.2, we have $W(G_1) \leq \frac{n^3-19n+54}{6}$, the equality holds if and only if $G_1 \cong B_n^{(1)}$. Together with $W(G) \leq W(G_1)$, we can obtain that $W(G) \leq \frac{n^3-19n+54}{6}$ with equality if and only if $G \cong B_n^{(1)}$. By Lemma 3.2, we have $Sz(G) - W(G) \geq 2$. Therefore,

$$\begin{aligned} \frac{Sz(G)}{W(G)} - 1 &= \frac{Sz(G) - W(G)}{W(G)} \\ &\geq \frac{6(Sz(G) - W(G))}{n^3 - 19n + 54} \\ &\geq \frac{12}{n^3 - 19n + 54}, \end{aligned} \tag{34}$$

$$\tag{35}$$

where the equality in (34) holds if and only if $G \cong B_n^{(1)}$; while the equality in (35) holds if and only if $G \cong G'$, where G' is depicted in Fig. 2. Hence, $G \cong B_n^{(1)}$.

Therefore, in view of (32), (33) and (35), we have $\frac{Sz(G)}{W(G)} \geq 1 + \frac{12}{n^3-19n+54}$ with equality if and only if $G \cong B_n^{(1)}$, as desired. \square

Now, we determine the second smallest value on $Sz^*(G)/W(G)$ among all connected cyclic graphs.

Proof of Theorem 2.4. It is straightforward to check that $\frac{Sz^*(H_n^2)}{W(H_n^2)} = 1 + \frac{24(n-2)}{n^3-19n+54}$. Note that $\frac{Sz^*(G)}{W(G)} \geq 1 + \frac{3(n^2+4n-6)}{2(n^3-7n+12)}$ if G is a non-bipartite graph; whereas $\frac{Sz^*(G)}{W(G)} \geq 1 + \frac{24(n-2)}{n^3-13n+36}$ if G is a bipartite graph and the equality holds if and only if $G \cong L_{n,4}$. Hence, combining with Theorem 1.4, it suffices

to characterize the bipartite graphs with the smallest value on $Sz^*(G)/W(G)$ for $G \not\cong L_{n,4}$. We may complete our proof by comparing this smallest value with $1 + \frac{3(n^2+4n-6)}{2(n^3-7n+12)}$.

Let G be a bipartite graph with at least one cycle and $G \not\cong L_{n,4}$. In this case, $Sz^*(G) = Sz(G)$. If $G \cong C_4(P_{n-4}, P_1, P_2, P_1)$, then by a direct calculation, we have

$$\frac{Sz^*(G)}{W(G)} = 1 + \frac{36(n-3)}{n^3-19n+66} > 1 + \frac{24(n-2)}{n^3-19n+54}.$$

Now, consider $G \not\cong C_4(P_{n-4}, P_1, P_2, P_1)$. Note that $W(G) < W(G-e)$ for any non-cut edge $e \in E_G$. Thus, we may repeatedly delete the non-cut edges of G until the resulting graph is a unicyclic graph G_1 such that $W(G) \leq W(G_1)$. Clearly, G_1 is bipartite. If $G_1 \cong L_{n,4}$, as $G \not\cong L_{n,4}$, then there exist some edges e_1, e_2, \dots, e_k , $k \geq 1$, such that $G = G_1 + \{e_1, e_2, \dots, e_k\}$, where G_1 is depicted in Fig. 5(c). Then we construct a new unicyclic graph G_2 as follows:

$$G_2 = G_1 - u_3v_3 + e_1.$$

Clearly, G_2 is not in $\{L_{n,4}, C_4(P_{n-4}, P_1, P_2, P_1)\}$. It is routine to check that $G = G_2 + \{u_3v_3, e_2, \dots, e_k\}$. Thus, $W(G) < W(G_2)$.

If $G_1 \cong C_4(P_{n-4}, P_1, P_2, P_1)$, as $G \not\cong C_4(P_{n-4}, P_1, P_2, P_1)$, then there exist some edges e_1, e_2, \dots, e_t , $t \geq 1$, such that $G = G_1 + \{e_1, e_2, \dots, e_t\}$, where G_1 is depicted in Fig. 5(d). Then we construct a new unicyclic graph G_3 as follows:

$$G_3 = G_1 - u_4v_4 + e_1.$$

Clearly, G_3 is not in $\{L_{n,4}, C_4(P_{n-4}, P_1, P_2, P_1)\}$. It is routine to check that $G = G_3 + \{u_4v_4, e_2, \dots, e_t\}$. Thus, $W(G) < W(G_3)$.

Thus, we may assume that G_1 is not in $\{L_{n,4}, C_4(P_{n-4}, P_1, P_2, P_1)\}$. By Theorem 2.1, we have $W(G_1) \leq \frac{n^3-19n+54}{6}$, the equality holds if and only if $G_1 \cong H_n^2, H_n^3$ if $n \geq 12$, $G_1 \cong H_{11}^2, H_{11}^3$ or $C_3(P_6, P_4, P_1)$ if $n = 11$ and $G_1 \cong H_{10}^3$, or H_{10}^3 if $n = 10$. Together with the fact $W(G) \leq W(G_1)$ and G is bipartite, we can obtain that $W(G) \leq \frac{n^3-19n+54}{6}$ with equality if and only if $G \cong H_n^2$ if $n \geq 10$. By Theorem 1.1, we have $Sz^*(G) - W(G) \geq 4n - 8$. Therefore,

$$\begin{aligned} \frac{Sz^*(G)}{W(G)} - 1 &= \frac{Sz^*(G) - W(G)}{W(G)} \\ &\geq \frac{6(Sz(G) - W(G))}{n^3 - 19n + 54} \end{aligned} \quad (36)$$

$$\geq \frac{24(n-2)}{n^3 - 19n + 54}, \quad (37)$$

where the equality in (36) holds if and only if $G \cong H_n^2$ if $n \geq 10$; by Theorem 1.1, the equality in (37) holds if and only if G is composed of a cycle C_4 on 4 vertices, and one tree rooted at a vertex of the C_4 . Hence, $\frac{Sz^*(G)}{W(G)} = 1 + \frac{24(n-2)}{n^3-19n+54}$ with equality if and only if $G \cong H_n^2$.

Note that $\frac{3(n^2+4n-6)}{2(n^3-7n+12)} > \frac{24(n-2)}{n^3-19n+54}$ for $n \geq 10$. Thus, we have $\frac{Sz^*(G)}{W(G)} \geq 1 + \frac{24(n-2)}{n^3-19n+54}$ with equality if and only if $G \cong H_n^2$.

This completes the proof. \square

Acknowledgements

The authors would like to express their sincere gratitude to all of the referees for their insightful comments and suggestions, which led to a number of improvements to this paper.

References

- M. Aouchiche, G. Caporossi, P. Hansen, M. Laffay, AutoGraphiX: a survey, *Electron. Notes Discrete Math.* 22 (2005) 515-520.
- M. Aouchiche, P. Hansen, On a conjecture about the Szeged index, *European J. Combin.* 31 (2010) 1662-1666.
- J.A. Bondy, U.S.R. Murty, *Graph Theory*, in: GTM, vol. 224, Springer, 2008.
- G. Caporossi, P. Hansen, Variable neighborhood search for extremal graphs 1. The AutoGraphiX system, *Discrete Math.* 212 (2000) 29-44.
- L.L. Chen, X.L. Li, M.M. Liu, I. Gutman, On a relation between the Szeged and the Wiener indices of bipartite graphs, *Trans. Comb.* 1 (4) (2012) 43-49.
- L.L. Chen, X.L. Li, M.M. Liu, The (revised) Szeged index and the Wiener index of a non-bipartite graph, *European J. Combin.* 36 (2014) 237-246.
- D.M. Cvetković, M. Boob, H. Sachs, *Spectral of Graphs. Theory and Application*, New York, London, 1979.
- H. Deng, The trees on $n \geq 9$ vertices with the first to seventeenth greatest Wiener indices are chemical trees, *MATCH Commun. Math. Comput. Chem.* 57 (2007) 393-402.
- A. Dobrynin, R. Entringer, I. Gutman, Wiener index of trees: theory and applications, *Acta Appl. Math.*, 66 (2001) 211-249.
- A. Dobrynin, I. Gutman, On a graph invariant related to the sum of all distances in a graph, *Publ. Inst. Math.(Beograd) (N.S.)* 56 (1994) 18-22.
- A. Dobrynin, I. Gutman, Solving a problem connected with distances in graphs, *Graph Theory Notes N. Y.* 28 (1995) 21-23.
- A.A. Dobrymin, I. Gutman, S. Klavžar, P. Žigert, Wiener index of hexagonal systems, *Acta Appl. Math.* 72 (2002) 247-294.
- H. Dong, B. Zhou, Maximum Wiener index of unicyclic graphs with fixed maximum degree, *Ars Combin.* 103 (2012) 407-416.
- Z. Du, A. Ilić, On AGC conjectures regarding average eccentricity, *MATCH Commun. Math. Comput. Chem.* 69 (2013) 579-587.
- R.C. Entringer, D.E. Jackson, D.A. Snyder, Distance in graphs, *Czech. Math. J.* 26 (1976) 283-296.

- I. Gutman, A formula for the Wiener number of trees and its extension to graphs containing cycles, *Graph Theory Notes N. Y.* 27 (1994) 9-15.
- I. Gutman, O.E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer, Berlin, 1986.
- P. Hansen et al. Computers and conjectures in chemical graph theory, in: *Plenary Talk in the International Conference on Mathematical Chemistry*, August 4-7, 2010, Xiamen, China.
- A. Ilić, Note on PI and Szeged indices, *Math. Comput. Modelling* 52 (2010) 1570-1576.
- H. Khodashenas, M.J. Nadjafi-Arani, A.R. Ashrafi, I. Gutman, A new proof of the Szeged-Wiener theorem, *Kragujevac J. Math.* 35 (2011) 165-172.
- S. Klavžar, M.J. Nadjafi-Arani, Wiener index versus Szeged index in networks, *Discrete Appl. Math.* 161 (2013) 1150-1153.
- S. Klavžar, M.J. Nadjafi-Arani, Improved bounds on the difference between the Szeged index and the Wiener index of graphs, *European J. Combin.* 39 (2014) 148-156.
- S. Klavžar, A. Rajapakse, I. Gutman, The Szeged and the Wiener index of graphs, *Appl. Math. Lett.* 9 (1996) 45-49.
- M. Knor, R. Škrekovski, A. Tepeh, Mathematical aspects of Wiener index, *Ars Math. Contemp.* 11 (2016) 327-352.
- S.C. Li, Y.B. Song, On the sum of all distances in bipartite graphs, *Discrete Appl. Math.* 169 (2014) 176-185.
- S.C. Li, H.H. Zhang, Proofs of three conjectures on the quotients of the (revised) Szeged index and the Wiener index and beyond, *Discrete Math.* 340 (2017) 311-324.
- X.L. Li, M.M. Liu, Bicyclic graphs with maximal revised Szeged index, *Discrete Appl. Math.* 161 (2013) 2527-2531.
- R.F. Liu, X. Du, H.C. Jia, Wiener index on traceable and Hamiltonian graphs, *Bull. Aust. Math. Soc.* 94 (2016) 362-372.
- B.D. McKay, A. Piperno, Practical Graph Isomorphism, II, *J. Symbolic Comput.* 60 (2014) 94-112, <http://dx.doi.org/10.1016/j.jsc.2013.09.003>
- S. Mukwembi, T. Vetrík, Wiener index of trees of given order and diameter at most 6, *Bull. Austr. Math. Soc.* 89 (2014) 379-396.
- M.J. Nadjafi-Arani, H. Khodashenas, A.R. Ashrafi, On the differences between Szeged and Wiener indices of graphs, *Discrete Math.* 311 (2011) 2233-2237.
- M.J. Nadjafi-Arani, H. Khodashenas, A.R. Ashrafi, Graphs whose Szeged and Wiener numbers differ by 4 and 5, *Math. Comput. Modelling* 55 (2012) 1644-1648.

- T. Pisanski, M. Randić, Use of the Szeged index and the revised Szeged index for measuring network bipartivity, *Discrete Appl. Math.* 158 (2010) 1936-1944.
- M. Randić, On generalization of Wiener index for cyclic structures, *Acta Chim. Slov.* 49 (2002) 483-496.
- S. Simić, I. Gutman, V. Baltić, Some graphs with extremal Szeged index, *Math. Slovaca* 50 (2000) 1-15.
- H. Wiener, Structural determination of paraffin boiling points, *J. Am. Chem. Soc.* 69 (1947) 17-20.
- R. Xing, B. Zhou, On the revised Szeged index, *Discrete Appl. Math.* 159 (2011) 69-78.
- K.X. Xu, K.C. Das, Extremal unicyclic and bicyclic graphs with respect to Harary index, *Bull. Malays. Math. Sci. Soc.* 36 (2013), 373-383.
- Z.K. Tang, H.Y. Deng, The (n, n) -graphs with the first three extremal Wiener indices, *J. Math. Chem.* 43 (2008) 60-74.
- H.H. Zhang, S.C. Li, L.F. Zhao, On the further relation between the (revised) Szeged index and the Wiener index of graphs, *Discrete Appl. Math.* 206 (2016) 152-164.
- X.D. Zhang, Y. Liu, M.X. Han, Maximum Wiener index of trees with given degree sequence, *MATCH Commun. Math. Comput. Chem.* 64 (2010) 661-682.