

On the number of spanning trees of $K_n^m \pm G$ graphs

Stavros D. Nikolopoulos and Charis Papadopoulos[†]

Department of Computer Science, University of Ioannina, P.O.Box 1186, GR-45110 Ioannina, Greece
 {stavros, charis}@cs.uoi.gr

received June 9, 2005, revised April 14, 2006, accepted July 25, 2006.

The K_n -complement of a graph G , denoted by $K_n - G$, is defined as the graph obtained from the complete graph K_n by removing a set of edges that span G ; if G has n vertices, then $K_n - G$ coincides with the complement \overline{G} of the graph G . In this paper we extend the previous notion and derive determinant based formulas for the number of spanning trees of graphs of the form $K_n^m \pm G$, where K_n^m is the complete multigraph on n vertices with exactly m edges joining every pair of vertices and G is a multigraph spanned by a set of edges of K_n^m ; the graph $K_n^m + G$ (resp. $K_n^m - G$) is obtained from K_n^m by adding (resp. removing) the edges of G . Moreover, we derive determinant based formulas for graphs that result from K_n^m by adding and removing edges of multigraphs spanned by sets of edges of the graph K_n^m . We also prove closed formulas for the number of spanning tree of graphs of the form $K_n^m \pm G$, where G is (i) a complete multipartite graph, and (ii) a multi-star graph. Our results generalize previous results and extend the family of graphs admitting formulas for the number of their spanning trees.

Keywords: Kirchhoff matrix tree theorem, complement spanning tree matrix, spanning trees, K_n -complements, multigraphs.

1 Introduction

The number of spanning trees of a graph G , denoted by $\tau(G)$, is an important, well-studied quantity in graph theory, and appears in a number of applications. Most notable application fields are network reliability [9, 17, 22], enumerating certain chemical isomers [5], and counting the number of Eulerian circuits in a graph [15].

Thus, both for theoretical and for practical purposes, we are interested in deriving formulas for the number of spanning trees of a graph G , and also of the K_n -complement of G ; the K_n -complement of a graph G , denoted by $K_n - G$, is defined as the graph obtained from the complete graph K_n by removing a set of edges (of the graph K_n) that span G ; if G has n vertices, then $K_n - G$ coincides with the complement \overline{G} of the graph G . Many cases have been examined depending on the choice of G . For example, there exist closed formulas for the cases where G is a pairwise disjoint set of edges [24], a chain of edges [16], a cycle [11], a star [20], a multi-star [19, 25], a multi-complete/star graph [7],

[†]Current address: Department of Informatics, University of Bergen, N-5020 Bergen, Norway; charis@ii.uib.no

a labelled molecular graph [5], and more recent results when G is a circulant graph [12, 26], a quasi-threshold graph [18], and so on (see Berge [2] for an exposition of the main results).

In this paper, we extend the previous notion and consider graphs that result from the complete multigraph K_n^m by removing multiple edges; we denote by K_n^m the complete multigraph on n vertices with exactly m edges joining every pair of vertices. Based on the properties of the Kirchhoff matrix, which permits the calculation of the number of spanning trees of any given graph, we derive a determinant based formula for the number of spanning trees of the graph $K_n^m - G$, where G is a subgraph of K_n^m , and, thus, it is a multigraph. Note that, if $m = 1$ then $K_n^m - G$ coincides with the graph $K_n - G$.

We also consider graphs that result from the complete multigraph K_n^m by adding multiple edges. More precisely, we consider multigraphs of the form $K_n^m + G$ that result from the complete multigraph K_n^m by adding a set of edges (of the graph K_n^m) that span G . Again, based on the properties of the Kirchhoff matrix, we derive a determinant based formula for the number of spanning trees of the graph $K_n^m + G$. To the best of our knowledge, not as much seems to be known about the number $\tau(K_n^m + G)$. Bedrosian in [1] considered the number $\tau(K_n + G)$ for some simple configurations of G , i.e., when G forms a cycle, a complete graph, or when its vertex set is quite small. More recently, Golin et al. in [12] derive closed formula for the number $\tau(K_n + G)$ using Chebyshev polynomials, introduced in [4], for the case where G forms a circulant graph.

We denote $K_n^m \pm G$ the family of graphs of the forms $K_n^m + G$ and $K_n^m - G$, and derive a determinant based formula for the number $\tau(K_n^m \pm G)$. Moreover, based on these results, we generalize our formulas and extend the family $K_n^m \pm G$ to the more general family of graphs $F_n^m \pm G$, where F_n^m is the complete multigraph on n vertices with at least $m \geq 1$ edges joining every pair of vertices.

Based on our results—that is, the determinant based formulas for the number of spanning trees of the family of graphs $\tau(K_n^m \pm G)$, and using standard algebraic techniques, we generalize known closed formulas for the number of spanning trees of simple graphs of the form $K_n - G$. In particular, we derive closed formulas for the number of spanning trees $\tau(K_n^m \pm G)$, in the case where G forms (i) a complete multipartite graph, and (ii) a multi-star graph.

We point out that our proposed formulas express the number of spanning trees $\tau(K_n^m \pm G)$ as a function of the determinant of a matrix that can be easily constructed from the adjacency relation of the graph G . Our results generalize previous results and extend the family of graphs of the form $K_n^m \pm G$ admitting formulas for the number of their spanning trees.

2 Preliminaries

We consider finite undirected simple graphs and multigraphs with no loops; the term *multigraph* is used when multiple edges are allowed in a graph. For a graph G , we denote by $V(G)$ and $E(G)$ the vertex set and edge set of G , respectively.

The *multiplicity* of a vertex-pair (v, u) of a graph G , denoted by $\ell_G(vu)$, is the number of edges joining the vertices v and u in G . The minimum multiplicity among all the vertex-pairs of G is denoted $\lambda(G)$ while $\Lambda(G)$ is the largest such number. Thus, if $\lambda(G) > 0$, then every pair of vertices in G is connected with at least $\lambda(G)$ edges; if $\Lambda(G) = 1$, then G contains no multiple edges, that is, G is a *simple* graph (note that a simple graph or a multigraph contains no loops). The *degree* of a vertex v of a graph G , denoted by $d_G(v)$, is the number of edges incident with v in G . The minimum degree among the vertices of G is denoted $\delta(G)$ while $\Delta(G)$ is the largest such number.

It is worth noting that a multigraph can also be viewed as a simple graph whose edges are labeled with

non-zero integers (weights) which correspond to the multiplicity of the edges. Thus, a multigraph can be thought as a weighted graph. In this paper we adopt the standard approach.

Let \mathcal{F} be the family of complete multigraphs on n vertices with multiplicity at least $m \geq 1$, and let $F_n^m \in \mathcal{F}$. Then, $\lambda(F_n^m) = m$, since F_n^m has at least $m \geq 1$ edges joining every pair of its vertices. A complete multigraph on n vertices with exactly m edges joining every pair of its vertices is called *m-complete multigraph* and denoted by K_n^m . Thus, for the m -complete multigraph K_n^m we have that $\lambda(K_n^m) = \Lambda(K_n^m) = m$ and $\delta(K_n^m) = \Delta(K_n^m) = (n - 1)m$. Note that, the 1-complete multigraph is the graph K_n . By definition, every complete multigraph F_n^m contains a subgraph isomorphic to an m -complete multigraph K_n^m .

Let F_n^m be a complete multigraph and let \mathcal{C} be a set of edges of F_n^m such that the graph which is obtained from F_n^m by removing the edges of \mathcal{C} is an m -complete multigraph K_n^m ; the graph spanned by the edges of \mathcal{C} is called a *characteristic graph* of F_n^m and denoted by $\mathcal{H}(F_n^m)$. By definition, a characteristic graph $\mathcal{H}(F_n^m)$ contains no isolated vertex.

Let G and H be two multigraphs. The graph $G + H$ is defined as follows:

$$V(G + H) = V(G) \cup V(H)$$

and

$$vu \in E(G + H) \iff vu \in E(G) \text{ or } vu \in E(H).$$

By definition, both graphs G and H are subgraphs of $G + H$. Moreover, if $v, u \in V(G) \cap V(H)$, then $\ell_{G+H}(vu) = \ell_G(vu) + \ell_H(vu)$.

Let G and H be two multigraphs such that $E(H) \subseteq E(G)$. The graph $G - H$ is defined as the graph obtained from G by removing the edges of H .

Having defined the graphs $G + H$ and $G - H$, it is easy to see that $F_n^m = K_n^m + \mathcal{H}(F_n^m)$ and $K_n^m = F_n^m - \mathcal{H}(F_n^m)$. In general, $\mathcal{H}(F_n^m) \neq F_n^m - K_n^m$; the equality holds if $\mathcal{H}(F_n^m)$ has n vertices.

The *adjacency matrix* of a multigraph G on n vertices, denoted by $A(G)$, is an $n \times n$ matrix with diagonal elements $A(G)[i, i] = 0$ and off-diagonal elements $A(G)[i, j] = \ell_G(v_i v_j)$. The *degree matrix* of the multigraph G , denoted by $D(G)$, is an $n \times n$ matrix with diagonal elements $D(G)[i, i] = d_G(v_i)$ and off-diagonal elements $D(G)[i, j] = 0$. Throughout the paper empty entries in matrices represent 0s.

For an $n \times n$ matrix M , the $(n - 1)$ -st order *minor* μ_j^i is the determinant of the $(n - 1) \times (n - 1)$ matrix obtained from M after having deleted row i and column j ; the i -th *cofactor* equals μ_i^i . The *Kirchhoff matrix* $L(G)$ (also known as the *Laplacian matrix*) for a multigraph G on n vertices is an $n \times n$ matrix with elements

$$L(G)[i, j] = \begin{cases} d_G(v_i) & \text{if } i = j, \\ -\ell_G(v_i v_j) & \text{otherwise} \end{cases}$$

where $d_G(v_i)$ is the degree of vertex v_i in the graph G and $\ell_G(v_i v_j)$ is the number of edges joining the vertices v_i and v_j in G . The matrix $L(G)$ is symmetric, has nonnegative real eigenvalues and its determinant is equal to zero. Note that, $L(G) = D(G) - A(G)$.

The *Kirchhoff matrix tree theorem* [3] is one of the most famous results in graph theory. It provides a formula for the number of spanning trees of a graph G in terms of the cofactors of G 's Kirchhoff matrix; it is stated as follows:

Theorem 2.1 (Kirchhoff Matrix Tree Theorem [3]) For any multigraph G with $L(G)$ defined as above, the cofactors of $L(G)$ have the same value, and this value equals the number of spanning trees $\tau(G)$ of the multigraph G .

The Kirchhoff matrix tree theorem provides a powerful tool for computing the number $\tau(G)$ of spanning trees of a graph G . For this computation, we first form the Kirchhoff matrix $L(G)$ of the graph G and obtain the $(n - 1) \times (n - 1)$ matrix $L_i(G)$ from $L(G)$ by removing its i -th row and column (arbitrary), and then compute the determinant of the matrix $L_i(G)$. We note that the determinant of $L_i(G)$ simply counts the number of spanning trees rooted at vertex $v_i \in V(G)$, which justifies the reason of removing the i -th row and the i -th column from $L(G)$. A combinatorial proof of the Kirchhoff matrix tree theorem can be found in [6].

The number of spanning trees of a graph G can be computed directly (without removing any row or column) in terms of a matrix $L'(G)$ similar to the Kirchhoff matrix $L(G)$, which is associated with the graph G [23], or, alternatively, it can be computed by defining a characteristic polynomial $\det(L(G) - xI)$ on $L(G)$; the latter approach takes into account the computation of the eigenvalues of the matrix $L(G)$ (see [4, 5, 10, 12, 22, 26]).

In our work, we express the number of spanning trees of a graph of the form $K_n^m \pm G$, where G is a subgraph of K_n^m on p vertices, in terms of a $p \times p$ matrix $B(G)$ associated with the graph G , and not in terms of an $n \times n$ matrix $L(K_n^m \pm G)$ associated with the whole graph $K_n^m \pm G$.

3 The $K_n^m \pm G$ graphs

In this section, we consider graphs that result from the m -complete multigraph K_n^m by removing or/and adding multiple edges. We are interested in deriving determinant based formulas for the number of spanning trees of the graphs $K_n^m - G$ and $K_n^m + G$, where G is a multigraph spanned by a set of edges $S \subseteq E(K_n^m)$. To this end, we define a parameter α as follows: (i) $\alpha = 1$, for the case $K_n^m + G$, and (ii) $\alpha = -1$, for the case $K_n^m - G$. In other words, $\alpha = \pm 1$, according to $K_n^m \pm G$. Based on the value of α we conclude with the following result.

Let G be a multigraph spanned by a set of edges of the graph K_n^m . We derive formulas for the number of spanning trees of the graph $K_n^m \pm G$; the graph G has $p \leq n$ vertices and $\Lambda(G) \leq m$.

In order to compute the number $\tau(K_n^m - G)$ we will make use of Theorem 2.1. Thus, we consider the $n \times n$ Kirchhoff matrix $L = L(K_n^m \pm G)$, which has the form:

$$L = \begin{bmatrix} m(n-1) \cdots & -m & & -m & & -m & \cdots & -m \\ \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ -m & \cdots & m(n-1) & & -m & & -m & -m \\ -m & \cdots & -m & m(n-1) + \alpha \cdot d_G(v_1) & & & & -m - \alpha \cdot \ell_G(v_j v_i) \\ -m & \cdots & -m & & m(n-1) + \alpha \cdot d_G(v_2) & & & \\ \vdots & \cdots & \vdots & & & \ddots & & \\ -m & \cdots & -m & -m - \alpha \cdot \ell_G(v_i v_j) & & & m(n-1) + \alpha \cdot d_G(v_p) \end{bmatrix} \quad (1)$$

where $d_G(v_i)$ is the degree of the vertex $v_i \in G$ and $\ell_G(v_i v_j)$ is the multiplicity of the vertices v_i and v_j in G . The entries of the off-diagonal positions $(n - p + i, n - p + j)$ of the matrix L are equal to

$-m - \alpha \cdot \ell_G(v_i v_j)$, $1 \leq i, j \leq p$. Note that, the first $n - p$ rows and columns of L correspond to the $n - p$ vertices of the set $V(K_n^m) - V(G)$ and, thus, they have degree $m(n - 1)$ in $K_n^m \pm G$.

Let L_1 be the $(n - 1) \times (n - 1)$ matrix obtained from L by removing its first row and column. Then, from Theorem 2.1 we have that

$$\tau(K_n^m - G) = \det(L_1).$$

In order to compute the determinant of the matrix L_1 , we add one row and one column to the matrix L_1 ; the resulting $n \times n$ matrix L'_1 has 1 in position $(1, 1)$, $-m$ in positions $(1, j)$, $2 \leq j \leq n$, and 0 in positions $(i, 1)$, $2 \leq i \leq n$. It is easy to see that, $\det(L'_1) = \det(L_1)$. Thus, the $n \times n$ matrix L'_1 has the following form:

$$L'_1 = \begin{bmatrix} 1 & -m & \cdots & -m & & -m & & \cdots & -m \\ 0 & m(n-1) & \cdots & -m & & -m & & \cdots & -m \\ \vdots & \vdots & \ddots & \vdots & & \vdots & & \vdots & \vdots \\ 0 & -m & \cdots & m(n-1) & & -m & & \cdots & -m \\ 0 & -m & \cdots & -m & \cdots & m(n-1) + \alpha \cdot d_G(v_1) & & & -m - \alpha \cdot \ell_G(v_j v_i) \\ 0 & -m & \cdots & -m & & & \cdots & & m(n-1) + \alpha \cdot d_G(v_2) \\ \vdots & \vdots & \cdots & \vdots & & & \ddots & & \vdots \\ 0 & -m & \cdots & -m & & -m - \alpha \cdot \ell_G(v_i v_j) & & & m(n-1) + \alpha \cdot d_G(v_p) \end{bmatrix}.$$

We denote by L''_1 the matrix obtained from L'_1 after multiplying the first row of L'_1 by -1 and adding it to the next $n - 1$ rows. Thus, the determinant of L''_1 is equal to the determinant of L'_1 . Moreover it becomes:

$$\det(L''_1) = \begin{vmatrix} 1 & -m & \cdots & -m & & -m & & \cdots & -m \\ -1 & mn & & & & & & & \\ \vdots & & \ddots & & & & & & \\ -1 & & & mn & & & & & \\ -1 & & & & \cdots & mn + \alpha \cdot d_G(v_1) & & & -\alpha \cdot \ell_G(v_j v_i) \\ -1 & & & & & & \cdots & & mn + \alpha \cdot d_G(v_2) \\ \vdots & & & & & & \ddots & & \vdots \\ -1 & & & & & -\alpha \cdot \ell_G(v_i v_j) & & & mn + \alpha \cdot d_G(v_p) \end{vmatrix},$$

where the entries of the off-diagonal positions $(n - p + i, n - p + j)$ of the matrix L''_1 are equal to $-\alpha \cdot \ell_G(v_i v_j)$, $1 \leq i, j \leq p$. Note that, the first $n - p$ rows of the matrix L''_1 have non-zero elements in positions $(1, i)$ and (i, i) , $2 \leq i \leq n - p$. We observe that the sum of all the elements on each row of L''_1 , except of the first row, is equal to $mn - 1$; recall that, $d_G(v_i) = \sum_{1 \leq j \leq p} \ell_G(v_i v_j)$, for every $v_i \in V(G)$.

Thus, we multiply each column of matrix L''_1 by $\frac{1}{mn}$ and add it to the first column, and we obtain:

$$\det(L''_1) = \begin{vmatrix} \frac{1}{n} & -m & \cdots & -m & & -m & & \cdots & -m \\ 0 & mn & & & & & & & \\ \vdots & & \ddots & & & & & & \\ 0 & & & mn & & & & & \\ 0 & & & \cdots & mn + \alpha \cdot d_G(v_1) & & & & -\alpha \cdot \ell_G(v_j v_i) \\ 0 & & & & \cdots & mn + \alpha \cdot d_G(v_2) & & & \\ \vdots & & & & & & \ddots & & \\ 0 & & & & & & & -\alpha \cdot \ell_G(v_i v_j) & mn + \alpha \cdot d_G(v_p) \end{vmatrix} \quad (2)$$

$$= m \cdot (mn)^{n-p-2} \cdot \det(B),$$

where $B = mnI_p + \alpha L(G)$ is a $p \times p$ matrix; recall that, $L(G)$ is the Kirchhoff matrix of the multigraph G and thus, $L(G) = D(G) \pm A(G)$, where $D(G)$ and $A(G)$ are the degree matrix and the adjacency matrix of G , respectively. Concluding, we obtain the following result.

Theorem 3.1 *Let K_n^m be the m -complete multigraph on n vertices, and let G be a multigraph on p vertices such that $V(G) \subseteq V(K_n^m)$ and $E(G) \subseteq E(K_n^m)$. Then,*

$$\tau(K_n^m \pm G) = m \cdot (mn)^{n-p-2} \det(B),$$

where $B = mnI_p + \alpha \cdot L(G)$ is a $p \times p$ matrix, $\alpha = \pm 1$ according to $K_n^m \pm G$, and $L(G)$ is the Kirchhoff matrix of G .

We note that, for simple graphs K_n and G in case $K_n - G$, Theorem 3.1 has been stated first by Moon in [16] and numerous authors in various guises used it as a constructive tool to obtain formulas for the number of spanning trees of graphs of the type $K_n - G$.

In the previous theorem the graph G is a subgraph of K_n^m , and, thus, it has multiplicity $\Lambda(G) \leq m$. It follows that the graph $K_n^m + G$ has multiplicity $\Lambda(K_n^m + G) \leq 2m$. However, we can relax the previous restriction in the case of the graph $K_n^m + G$. It is not difficult to see that for the matrix L of Equation (1) we have $\lambda(G) \geq 0$. Thus, we can define the graph G to be a multigraph on p vertices such that $V(G) \subseteq V(K_n^m)$. The following theorem holds.

Theorem 3.2 *Let K_n^m be the m -complete multigraph on n vertices, and let G be a multigraph on p vertices such that $V(G) \subseteq V(K_n^m)$. Then,*

$$\tau(K_n^m + G) = m \cdot (mn)^{n-p-2} \cdot \det(B),$$

where $B = mnI_p + L(G)$ is a $p \times p$ matrix, and $L(G)$ is the Kirchhoff matrix of G .

4 The $F_n^m \pm G$ graphs

In this section we derive determinant based formulas for the number $\tau(F_n^m \pm G)$, where F_n^m is a complete multigraph and G is a subgraph of F_n^m . We first take into consideration the graph $K_n^m + G_1 - G_2$ and derive a determinant based formula for the number $\tau(K_n^m + G_1 - G_2)$, and, then, we derive a formula for the number $\tau(F_n^m \pm G)$ using the graph $K_n^m + G_1 - G_2$ and a characteristic graph $\mathcal{H}(F_n^m)$.

4.1 The case $K_n^m + G_1 - G_2$

Here, we consider graphs that result from the m -complete multigraph K_n^m by adding multiple edges of a graph G_1 and removing multiple edges from a graph G_2 . Let G_1 be a multigraph on p_1 vertices, such that $V(G_1) \subseteq V(K_n^m)$, and let G_2 be a multigraph on p_2 vertices, such that $V(G_2) \subseteq V(K_n^m)$ and $E(G_2) \subseteq E(K_n^m + G_1)$. We next focus on the graph $K_n^m + G_1 - G_2$, which is obtained from the m -complete multigraph K_n^m by first adding the edges of the graph G_1 and then removing from the resulting graph $K_n^m + G_1$ the edges of G_2 ; that is, $K_n^m + G_1 - G_2 = (K_n^m + G_1) - G_2$.

It is worth noting that $(K_n^m + G_1) - G_2 \neq K_n^m + (G_1 - G_2)$, since the graph G_2 is not necessarily a subgraph of G_1 . Moreover, $V(G_1) \neq V(G_2)$.

For the pair of multigraphs (G_1, G_2) we define the *union-stable* graphs G_1^* and G_2^* of (G_1, G_2) as follows: G_1^* is the multigraph that results from G_1 by adding in $V(G_1)$ the vertices of the set $V(G_2) - V(G_1)$ and G_2^* is the multigraph that results from G_2 by adding in $V(G_2)$ the vertices of the set $V(G_1) - V(G_2)$. Thus, by definition $V(G_1^*) = V(G_2^*)$.

By definition, both G_1^* and G_2^* are multigraphs on $p = |V(G_1) \cup V(G_2)|$ vertices, with at least $p - p_1$ and $p - p_2$ isolated vertices, respectively. Since $K_n^m + G_1^* - G_2^* = K_n^m + G_1 - G_2$, we focus on the graph $K_n^m + G_1^* - G_2^*$.

Based on Theorem 2.1, we construct the $n \times n$ Kirchoff matrix $L = L(K_n^m + G_1^* - G_2^*)$ associated with the graph $K_n^m + G_1^* - G_2^*$; it is similar to that of the case of $K_n^m \pm G$. The difference here is the $p \times p$ submatrix which is formed by the last p rows and columns of L , where $p = |V(G_1) \cup V(G_2)|$. More precisely, the matrix L has the following form:

$$L = \begin{bmatrix} m(n-1) & -m & \cdots & -m & -m & -m & \cdots & -m \\ -m & m(n-1) & \cdots & -m & -m & -m & \cdots & -m \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots \\ -m & -m & \cdots & m(n-1) & -m & -m & \cdots & -m \\ -m & -m & \cdots & -m & B'(G_1^*, G_2^*)[1, 1] & & & B'(G_1^*, G_2^*)[j, i] \\ -m & -m & \cdots & -m & & B'(G_1^*, G_2^*)[2, 2] & & \\ \vdots & \vdots & \cdots & \vdots & & & \ddots & \\ -m & -m & \cdots & -m & B'(G_1^*, G_2^*)[i, j] & & & B'(G_1^*, G_2^*)[p, p] \end{bmatrix}$$

where the $p \times p$ submatrix $B'(G_1^*, G_2^*)$ has elements

$$B'(G_1^*, G_2^*)[i, j] = \begin{cases} m(n-1) + d_{G_1^*}(v_i) - d_{G_2^*}(v_i) & \text{if } i = j, \\ -m - \ell_{G_1^*}(v_i v_j) + \ell_{G_2^*}(v_i v_j) & \text{otherwise.} \end{cases}$$

We note that $\ell_{G_1^*}(v_i v_j)$ and $\ell_{G_2^*}(v_i v_j)$ are the number of edges of the vertices v_i and v_j in G_1^* and G_2^* , respectively. The entries $d_{G_1^*}(v_i)$ and $d_{G_2^*}(v_i)$, $1 \leq i \leq p$, are the degrees of vertex v_i of G_1^* and G_2^* , respectively. Note that, $V(G_1^*) = V(G_2^*)$.

It is straightforward to apply a technique similar to that we have applied for the computation of the determinant of the matrix L'_1 in the case of $K_n^m - G$. Thus, in the case of $K_n^m + G_1^* - G_2^*$, the determinant

of the matrix L'_1 of Equation (2) becomes

$$\det(L'_1) = \begin{vmatrix} \frac{1}{n} & -m & \cdots & -m & & -m & & \cdots & -m \\ 0 & mn & & & & & & & \\ \vdots & & \ddots & & & & & & \\ 0 & & & mn & & & & & \\ 0 & & & & B(G_1^*, G_2^*)[1, 1] & & & & B(G_1^*, G_2^*)[j, i] \\ 0 & & & & & B(G_1^*, G_2^*)[2, 2] & & & \\ \vdots & & & & & & \ddots & & \\ 0 & & & & B(G_1^*, G_2^*)[i, j] & & & & B(G_1^*, G_2^*)[p, p] \end{vmatrix}$$

$$= m \cdot (mn)^{n-p-2} \cdot \det(B(G_1^*, G_2^*)),$$

where the $p \times p$ submatrix $B(G_1^*, G_2^*)$ has elements

$$B(G_1^*, G_2^*)[i, j] = \begin{cases} mn + d_{G_1^*}(v_i) - d_{G_2^*}(v_i) & \text{if } i = j, \\ -\ell_{G_1^*}(v_i v_j) + \ell_{G_2^*}(v_i v_j) & \text{otherwise.} \end{cases}$$

Since $B(G_1^*, G_2^*) = mnI_p + L(G_1^*) - L(G_2^*)$, we set $B = B(G_1^*, G_2^*)$ and obtain the following result.

Theorem 4.1 *Let K_n^m be the m -complete multigraph on n vertices, and let G_1, G_2 be two multigraphs such that $V(G_1) \subseteq V(K_n^m)$ and $E(G_2) \subseteq E(K_n^m + G_1)$. Then,*

$$\tau(K_n^m + G_1 - G_2) = m \cdot (mn)^{n-p-2} \det(B),$$

where $p = |V(G_1) \cup V(G_2)|$, $B = mnI_p + L(G_1^*) - L(G_2^*)$ is a $p \times p$ matrix, $L(G_1^*)$ and $L(G_2^*)$ are the Kirchhoff matrices of the union-stable graphs G_1^* and G_2^* of (G_1, G_2) , respectively.

4.2 The general case $F_n^m \pm G$

Let F_n^m be a complete multigraph on n vertices and let G be a subgraph of F_n^m . We will show that the previous theorem provides the key idea for computing the number $\tau(F_n^m \pm G)$, where $F_n^m \pm G$ is the graph that results from F_n^m by adding or removing the edges of G . Since $\lambda(F_n^m) > 0$, we have that $F_n^m = K_n^m + \mathcal{H}(F_n^m)$, where $\mathcal{H}(F_n^m)$ is a characteristic graph of F_n^m . Then we have that,

$$F_n^m \pm G = K_n^m + \mathcal{H}(F_n^m) \pm G.$$

The addition of the edges of G in the graph F_n^m , implies that $F_n^m + G = K_n^m + G'$, where the graph $G' = \mathcal{H}(F_n^m) + G$. Thus, for the computation of the number $\tau(F_n^m + G)$ we can apply Theorem 3.2. On the other hand, in the case of removal the edges of G from the graph F_n^m , for the computation of the number $\tau(F_n^m - G)$ we can apply Theorem 4.1, since $F_n^m - G = K_n^m + \mathcal{H}(F_n^m) - G$. Concluding we have the following result.

Lemma 4.1 *Let F_n^m be a complete multigraph on n vertices and $\mathcal{H}(F_n^m)$ be a characteristic graph of F_n^m , and let G be a subgraph of F_n^m . Then,*

$$\tau(F_n^m \pm G) = m \cdot (mn)^{n-p-2} \det(B),$$




Fig. 1: (a) A complete multipartite graph $K_{1,2,3}$ and (b) a multi-star graph $K_4(0, 2, 2, 3)$.

where $p = |V(\mathcal{H}(F_n^m)) \cup V(G)|$, $B = mnI_p + L(\mathcal{H}(F_n^m)^*) \pm L(G^*)$ is a $p \times p$ matrix and $L(\mathcal{H}(F_n^m)^*)$ and $L(G^*)$ are the Kirchhoff matrices of the union-stable graphs $\mathcal{H}(F_n^m)^*$ and G^* of $(\mathcal{H}(F_n^m), G)$, respectively.

Note that, we consider the graph $F_n^m \pm G$ and therefore G must be a subgraph of F_n^m . However in the case of the $F_n^m + G$ graph, similar to Theorem 3.2, it is obvious that G can be a graph spanned by any set of edges joining the vertices of F_n^m .

5 Classes of graphs

In this section, we generalize known closed formulas for the number of spanning trees of families of graphs of the form $K_n - G$. As already mentioned in the introduction there exist many cases for the $\tau(K_n - G)$, depending on the choice of G . The purpose of this section is to prove closed formulas for $\tau(K_n^m \pm G)$, by applying similar techniques to that of the case of $K_n - G$. Thus we derive closed formulas for the number of spanning trees $\tau(K_n^m \pm G)$, in the cases where G forms (i) a complete multipartite graph, and (ii) a multi-star graph.

5.1 Complete multipartite graphs

A graph is defined to be a *complete multipartite* (or complete k -partite) if there is a partition of its vertex set into k disjoint sets such that no two vertices of the same set are adjacent and every pair of vertices of different sets are adjacent. We denote a complete multipartite graph on p vertices by K_{m_1, m_2, \dots, m_k} , where $p = m_1 + m_2 + \dots + m_k$; see Figure 1(a). We note that the number of spanning trees of a complete multipartite graph has been considered by several authors in the past [8, 14].

Let $G = K_{m_1, m_2, \dots, m_k}$ be a complete multipartite graph on p vertices. In [21] it has been proved that the number of spanning trees of $K_n - G$ is given by the following formula:

$$\tau(K_n - G) = n^{n-p-1} (n-p)^{k-1} \prod_{i=1}^k (n - (p - m_i))^{m_i-1},$$

where p is the number of vertices of G .

In this section, we extend the previous result by deriving a closed formula for the number of their spanning trees for the graphs $K_n^m \pm G$, where G a complete multipartite graph on $p \leq n$ vertices. From Theorem 3.1, we construct the $p \times p$ matrix $B(G)$ and add one row and column to the matrix $B(G)$; the

resulting $(p + 1) \times (p + 1)$ matrix $B'(G)$ has 1 in position $(1, 1)$, α in positions $(1, j)$, $2 \leq j \leq p + 1$, and 0 in positions $(i, 1)$, $2 \leq i \leq p + 1$; recall that, $\alpha = \pm 1$. Thus, the resulting matrix $B'(G)$ has the following form:

$$B'(G) = \begin{bmatrix} 1 & \alpha & \alpha & \cdots & \alpha \\ & M_1 & -\alpha & \cdots & -\alpha \\ & -\alpha & M_2 & \cdots & -\alpha \\ & \vdots & \vdots & \ddots & \vdots \\ & -\alpha & -\alpha & \cdots & M_k \end{bmatrix},$$

where the diagonal $m_i \times m_i$ submatrices M_i have diagonal elements $mn + \alpha \cdot (p - m_i)$, $1 \leq i \leq k$. Note that, $\det(B(G)) = \det(B'(G))$.

In order to compute the determinant of the matrix $B'(G)$ we add the first row to the next p rows. Let $B''(G)$ be the resulting matrix. It follows that the determinant of $B'(G)$ is equal to the determinant of $B''(G)$. We multiply the $2, 3, \dots, p + 1$ columns of the matrix $B''(G)$ by $-1/(mn + \alpha \cdot p)$ and add them to the first column; note that, the sum of each row of the matrix $B''(G)$ is equal to $mn + \alpha \cdot p$. Thus, the determinant of matrix $B''(G)$ becomes:

$$\begin{aligned} \det(B''(G)) &= \begin{vmatrix} 1 - \frac{\alpha \cdot p}{mn + \alpha \cdot p} & \alpha & \alpha & \cdots & \alpha \\ & M'_1 & & & \\ & & M'_2 & & \\ & & & \ddots & \\ & & & & M'_k \end{vmatrix} \\ &= \frac{mn}{mn + \alpha \cdot p} \cdot \det(M'_1) \cdot \det(M'_2) \cdot \cdots \cdot \det(M'_k), \end{aligned} \tag{3}$$

where the $m_i \times m_i$ submatrices M'_i , $1 \leq i \leq k$, have the following form:

$$M'_i = \begin{vmatrix} mn + \alpha \cdot (p - m_i) + \alpha & \alpha & \cdots & \alpha \\ \alpha & mn + \alpha \cdot (p - m_i) + \alpha & \cdots & \alpha \\ \vdots & \vdots & \ddots & \vdots \\ \alpha & \alpha & \cdots & mn + \alpha \cdot (p - m_i) + \alpha \end{vmatrix}$$

For the determinant of matrix M'_i we multiply the first row by -1 and add it to the next $m_i - 1$ rows. Then, we add the columns of matrix M'_i to the first column. Observing that $mn + \alpha \cdot (p - m_i) + \alpha \cdot m_i = mn + \alpha \cdot p$, we obtain

$$\det(M'_i) = (mn + \alpha \cdot p) \cdot (mn + \alpha \cdot (p - m_i))^{m_i - 1}.$$

Thus, from Equation (3) we have the following result.

Theorem 5.1 *Let $G = K_{m_1, m_2, \dots, m_k}$ be a complete multipartite graph on $p = m_1 + m_2 + \cdots + m_k$ vertices. Then,*

$$\tau(K_n^m \pm G) = m \cdot (mn)^{n-p-1} (mn + \alpha \cdot p)^{k-1} \prod_{i=1}^k (mn + \alpha \cdot (p - m_i))^{m_i - 1},$$

where $p \leq n$ and $\alpha = \pm 1$ according to $K_n^m \pm G$.

Remark 5.1. The class of complete multipartite graphs contains the class of c-split graphs (complete split graphs); a graph is defined to be a c-split graph if there is a partition of its vertex set into a stable set S and a complete set K and every vertex in S is adjacent to all the vertices in K [13].

Thus, a c-split graph G on p vertices and $V(G) = K + S$ is a complete multipartite graph K_{m_1, m_2, \dots, m_k} with $m_1 = |S|$, $m_2 = m_3 = \dots = m_k = 1$ and $k = |K| + 1$. A closed formula for the number of spanning trees of the graph $K_n - G$ was proposed in [18], where G is a c-split graph.

Let G be a c-split graph on p vertices and let $V(G) = K + S$. Then, from Theorem 5.1 we obtain that the number of spanning trees of the graphs $K_n^m \pm G$ is given by the following closed formula:

$$\tau(K_n^m \pm G) = m \cdot (mn)^{n-p-1} (mn + \alpha \cdot |K|)^{|S|-1} (mn + \alpha \cdot p)^{|K|},$$

where $p = |K| + |S|$ and $p \leq n$. \square

5.2 Multi-star graphs

A multi-star graph, denoted by $K_r(b_1, b_2, \dots, b_r)$, consists of a complete graph K_r with vertices labelled v_1, v_2, \dots, v_r , and b_i vertices of degree one, which are incident with vertex v_i , $1 \leq i \leq r$ [7, 19, 25]; see Figure 1(b).

Let $G = K_r(b_1, b_2, \dots, b_r)$ be a multi-star graph on $p = r + b_1 + b_2 + \dots + b_r$ vertices. It has been proved [7, 19, 25] that the number of spanning trees of the graph $K_n - G$ is given by the following closed formula:

$$\tau(K_n - G) = n^{n-p-2} (n-1)^{p-r} \left(1 + \sum_{i=1}^r \frac{1}{q_i - 1} \right) \cdot \prod_{i=1}^r (q_i - 1),$$

where $q_i = n - (r - 1 + b_i) - \frac{b_i}{n-1}$.

In this section, based on Theorem 3.1, we generalize the previous result by deriving a closed formula for the number of spanning trees of the graphs $K_n^m \pm G$, where G is a multi-star on $p \leq n$ vertices. Let K_r be the complete graph of the multi-star graph G and let v_1, v_2, \dots, v_r be its vertices. The vertex set consisting of the vertex v_i and the b_i vertices of degree one which are incident with vertex v_i induces a star on $b_i + 1$ vertices, $1 \leq i \leq r$. We construct a $(b_i + 1) \times (b_i + 1)$ matrix M_i which corresponds to the star with center vertex v_i ; it has the following form:

$$M_i = \begin{bmatrix} mn + \alpha & & & -\alpha \\ & mn + \alpha & & -\alpha \\ & & \vdots & \\ -\alpha & -\alpha & \dots & mn + \alpha \cdot (r - 1 + b_i) \end{bmatrix},$$

where $\alpha = \pm 1$.

In order to compute the determinant of the matrix M_i we first multiply the first row by -1 and add it to the next $b_i - 1$ rows. We then add the b_i columns to the first column. Finally, we multiply the first column

Theorem 5.2 Let $G = K_r(b_1, b_2, \dots, b_r)$ be a multi-star graph on $p = r + b_1 + b_2 + \dots + b_r$ vertices. Then,

$$\tau(K_n^m \pm G) = m \cdot (mn)^{n-p-2} (mn + \alpha)^{p-r} \left(1 - \alpha \cdot \sum_{i=1}^r \frac{1}{q_i + \alpha} \right) \cdot \prod_{i=1}^r (q_i + \alpha),$$

where $p \leq n$, $q_i = mn + \alpha \cdot (r - 1 + b_i) - \frac{b_i}{mn + \alpha}$ and $\alpha = \pm 1$ according to $K_n^m \pm G$.

6 Concluding remarks

In this paper we derived determinant based formulas for the number of spanning trees of the family of graphs of the form $K_n^m \pm G$, and also for the more general family of graphs $F_n^m \pm G$, where K_n^m (resp. F_n^m) is the complete multigraph on n vertices with exactly (resp. at least) m edges joining every pair of vertices and G is a multigraph spanned by a set of edges of K_n^m (resp. K_n^m). Based on these determinant based formulas, we prove closed formulas for the number of spanning trees $\tau(K_n^m \pm G)$, in the case where G is (i) a complete multipartite graph, and (ii) a multi-star graph.

In light of our results, it would be interesting to consider the problem of proving closed formulas for the number of spanning tree $\tau(K_n^m \pm G)$ in the cases where G belongs to other classes of simple graphs or multigraphs. Moreover, instead of a multigraph G that we have taken into account, we do not know whether a generalization for an arbitrarily weighted graph G holds.

The problem of maximizing the number of spanning trees was solved for some families of graphs of the form $K_n - G$, where G is a multi-star graph, a union of paths and cycles, etc. (see [7, 11, 19, 22]). Thus, an interesting open problem is that of maximizing the number of spanning trees of graphs of the form $K_n^m \pm G$.

Acknowledgements

The authors would like to express their thanks to the anonymous referees whose suggestions helped improve the presentation of the paper.

References

- [1] S.D. Bedrosian, Generating formulas for the number of trees in a graph, *J. Franklin Inst.* **277** (1964) 313–326.
- [2] C. Berge, *Graphs and hypergraphs*, North-Holland, 1973.
- [3] N. Biggs, *Algebraic Graph Theory*, Cambridge University Press, London, 1974.
- [4] F.T. Boesch and H. Prodinger, Spanning tree formulas and Chebyshev polynomials, *Graphs and Combinatorics* **2** (1986) 191–200.
- [5] T.J.N. Brown, R.B. Mallion, P. Pollak, and A. Roth, Some methods for counting the spanning trees in labelled molecular graphs, examined in relation to certain fullerenes, *Discrete Appl. Math.* **67** (1996) 51–66.
- [6] S. Chaiken, A combinatorial proof of the all minors matrix tree theorem, *SIAM J. Algebraic Discrete Methods* **3** (1982) 319–329.

- [7] K.-L. Chung and W.-M. Yan, On the number of spanning trees of a multi-complete/star related graph, *Inform. Process. Lett.* **76** (2000) 113–119.
- [8] L. Clark, On the enumeration of spanning trees of the complete multipartite graph, *Bull. Inst. Combin. Appl.* **38** (2003) 50–60.
- [9] C.J. Colbourn, *The combinatorics of network reliability*, Oxford University Press, New York, 1980.
- [10] D.M. Cvetković, M. Doob and H. Sachs, *Spectra of graphs*, Academic Press, New York, 1980.
- [11] B. Gilbert and W. Myrvold, Maximizing spanning trees in almost complete graphs, *Networks* **30** (1997) 23–30.
- [12] M.J. Golin, X. Yong and Y. Zhang, Chebyshev polynomials and spanning tree formulas for circulant and related graphs, *Discrete Math.* **298** (2005) 334–364.
- [13] M.C. Golumbic, *Algorithmic graph theory and perfect graphs*, Academic Press, 1980.
- [14] R.P. Lewis, The number of spanning trees of a complete multipartite graph, *Discrete Math.* **197/198** (1999) 537–541.
- [15] F. Harary, *Graph theory*, Addison-Wesley, Reading, MA, 1969.
- [16] W. Moon, Enumerating labeled trees, in: F. Harary (Ed.), *Graph Theory and Theoretical Physics*, 261–271, 1967.
- [17] W. Myrvold, K.H. Cheung, L.B. Page, and J.E. Perry, Uniformly-most reliable networks do not always exist, *Networks* **21** (1991) 417–419.
- [18] S.D. Nikolopoulos and C. Papadopoulos, The number of spanning trees in K_n -complements of quasi-threshold graphs, *Graphs and Combinatorics* **20** (2004) 383–397.
- [19] S.D. Nikolopoulos and P. Rondogiannis, On the number of spanning trees of multi-star related graphs, *Inform. Process. Lett.* **65** (1998) 183–188.
- [20] P.V. O’Neil, The number of trees in a certain network, *Notices Amer. Math. Soc.* **10** (1963) 569.
- [21] P.V. O’Neil, Enumeration of spanning trees in certain graphs, *IEEE Trans. Circuit Theory* CT-17 (1970) 250.
- [22] L. Petingi, F. Boesch and C. Suffel, On the characterization of graphs with maximum number of spanning trees, *Discrete Math.* **179** (1998) 155–166.
- [23] H.N.V. Temperley, On the mutual cancellation of cluster integrals in Mayer’s fugacity series, *Proc. Phys. Soc.* **83** (1964) 3–16.
- [24] L. Weinberg, Number of trees in a graph, *Proc. IRE* 46 (1958), 1954–1955.
- [25] W.-M. Yan, W. Myrvold and K.-L. Chung, A formula for the number of spanning trees of a multi-star related graph, *Inform. Process. Lett.* **68** (1998) 295–298.
- [26] Y. Zhang, X. Yong and M.J. Golin, The number of spanning trees in circulant graphs, *Discrete Math.* **223** (2000) 337–350.