

Graphs of Edge-Intersecting and Non-Splitting One Bend Paths in a Grid *

Arman Boyacı¹ Tınaz Ekim¹ Mordechai Shalom^{1,2 †} Shmuel Zaks³

¹ Department of Industrial Engineering, Bogazici University, Istanbul, Turkey

² TelHai College, Upper Galilee, 12210, Israel

³ Department of Computer Science, Technion, Haifa, Israel

received 22nd Dec. 2015, revised 15th Sep. 2016, accepted 5th May 2017.

The families EPT (resp. EPG) Edge Intersection Graphs of Paths in a tree (resp. in a grid) are well studied graph classes. Recently we introduced the graph classes Edge-Intersecting and Non-Splitting Paths in a Tree (ENPT), and in a Grid (ENPG). It was shown that ENPG contains an infinite hierarchy of subclasses that are obtained by restricting the number of bends in the paths. Motivated by this result, in this work we focus on one bend ENPG graphs. We show that one bend ENPG graphs are properly included in two bend ENPG graphs. We also show that trees and cycles are one bend ENPG graphs, and characterize the split graphs and co-bipartite graphs that are one bend ENPG. We prove that the recognition problem of one bend ENPG split graphs is NP-complete even in a very restricted subfamily of split graphs. Last we provide a linear time recognition algorithm for one bend ENPG co-bipartite graphs.

Keywords: Intersection Graphs, Path Graphs, EPT Graphs, EPG Graphs

1 Introduction

1.1 Background

Given a host graph H and a set \mathcal{P} of paths in H , the Edge Intersection Graph of Paths (EP graph) of \mathcal{P} is denoted by $EP(\mathcal{P})$. The graph $EP(\mathcal{P})$ has a vertex for each path in \mathcal{P} , and two vertices of $EP(\mathcal{P})$ are adjacent if the corresponding two paths intersect in at least one edge. A graph G is EP if there exist a graph H and a set \mathcal{P} of paths in H such that $G = EP(\mathcal{P})$. In this case, we say that $\langle H, \mathcal{P} \rangle$ is an EP representation of G . We also denote by EP the family of all graphs G that are EP.

The main application area of EP graphs is communication networks. Messages to be delivered are sent through routes of a communication network. Whenever two paths use the same link on the communication network, we say that they conflict. Noting that this conflict model is equivalent to an EP graph, several optimization problems in communication networks (such as message scheduling) can be seen as graph problems (such as vertex coloring) in the corresponding EP graph.

*This work was supported in part by TUBITAK PIA BOSPHORUS Grant No. 111M303.

†Part of this work is accomplished while this author was visiting Bogazici University, Department of Industrial Engineering, under the TUBITAK 2221 Program whose support is gratefully acknowledged.

In many applications it turns out that the host graphs are restricted to certain families such as paths, cycles, trees, grids, etc. Several known graph classes are obtained with such restrictions: when the host graph is restricted to paths, cycles, trees and grids, we obtain interval graphs, circular-arc graphs, Edge Intersection Graph of Paths in a Tree (EPT) (see Golumbic and Jamison (1985a)), and Edge Intersection Graph of Paths in a Grid (EPG) (see Golumbic et al. (2009)), respectively.

Given a representation $\langle T, \mathcal{P} \rangle$ where T is a tree and \mathcal{P} is a set of paths of T , the graph of edge intersecting and non-splitting paths of $\langle T, \mathcal{P} \rangle$ (denoted by $\text{ENPT}(\mathcal{P})$) is defined as follows in Boyacı et al. (2015a): $\text{ENPT}(\mathcal{P})$ has a vertex v for each path P_v of \mathcal{P} and two vertices u, v of this graph are adjacent if the paths P_u and P_v edge-intersect and do not split (that is, their union is a path). We note that $\text{ENPT}(\mathcal{P})$ is a subgraph of $\text{EPT}(\mathcal{P})$. The motivation to study these graphs arises from all-optical Wavelength Division Multiplexing (WDM) networks in which two streams of signals can be transmitted using the same wavelength only if the paths corresponding to these streams do not split from each other (see Boyacı et al. (2015a) for a more detailed discussion). A graph G is an ENPT graph if there is a tree T and a set of paths \mathcal{P} of T such that $G = \text{ENPT}(\mathcal{P})$. Clearly, when T is a path, $\text{EPT}(\mathcal{P}) = \text{ENPT}(\mathcal{P})$ and this graph is an interval graph. Therefore, interval graphs are included in the class ENPT. In Boyacı et al. (2015b) we obtain the so-called ENP graphs by extending this definition to the case where the host graph is not necessarily a tree. In the same work, it has been shown that $\text{ENP} = \text{ENPG}$ where ENPG is the family of ENP graphs where the host graphs are restricted to grids. Whenever the host graph is a grid, it is common to use the following notion: a *bend* of a path on a grid is an internal point in which the path changes direction. An ENPG graph is B_k -ENPG if it has a representation in which every path has at most k bends.

1.2 Related Work

While ENPT and ENPG graphs have been recently introduced, EPT and EPG graphs are well studied in the literature. The recognition of EPT graphs is NP-complete (Golumbic and Jamison (1985b)), whereas one can solve in polynomial time the maximum clique (Golumbic and Jamison (1985b)) and the maximum stable set (Tarjan (1985)) problems in this class.

Several recent papers consider the edge intersection graphs of paths on a grid. Since all graphs are EPG (see Golumbic et al. (2009)), most of the studies focus on the sub-classes of EPG obtained by limiting the number of bends in each path. An EPG graph is B_k -EPG if it admits a representation in which every path has at most k bends. The work of Biedl and Stern (2010) investigates the minimum number k such that G has a B_k -EPG representation for some special graph classes. The work of Golumbic et al. (2009) studies the B_1 -EPG graphs. In particular it is shown that every tree is B_1 -EPG, and a characterization of C_4 representations is given. In Biedl and Stern (2010) the existence of an outer-planar graph which is not B_1 -EPG is shown. The recognition problem of B_1 -EPG graphs is shown to be NP-complete in Heldt et al. (2014). Similarly, in the class of B_1 -EPG, the minimum coloring and the maximum stable set problems are NP-complete (Epstein et al. (2013)), however one can solve in polynomial time the maximum clique problem (Epstein et al. (2013)). Asinowski and Ries (2012) give a characterization of graphs that are both B_1 -EPG and belong to some subclasses of chordal graphs. Recently, Cameron et al. (2016) consider subclasses of B_1 -EPG obtained by restricting the representations to contain only certain subsets of the four possible single bend rectilinear paths. It is shown that for each possible non-empty subset of these four shapes, the recognition of the corresponding subclass of B_1 -EPG is an NP-complete problem.

In Boyacı et al. (2015a) we defined the family of ENPT graphs and investigated the representations

of induced cycles. These representations turn out to be much more complex than their counterpart in the EPT graphs (discussed in Golombic and Jamison (1985a)). In Boyacı et al. (2015b) we extended this definition to the general case in which the host graph is not necessarily a tree. We showed that the family of ENP graphs coincides with the family of ENPG graphs, and that unlike EPG graphs, not every graph is ENPG. We also showed that, in a way similar to the family of EPG graphs, the sub families B_k -ENPG of ENPG contains an infinite subset totally ordered by proper inclusion.

1.3 Our Contribution

In this work, we consider B_1 -ENPG graphs. In Section 2 we present definitions and preliminary results among which we show that cycles and trees are B_1 -ENPG graphs. In Section 3 we show that the B_1 -ENPG recognition problem is NP-complete even for a very restricted subfamily of split graphs, i.e. graphs whose vertex sets can be partitioned into a clique and an independent set. In Section 4 we show that B_1 -ENPG graphs can be recognized in polynomial time within the family of co-bipartite graphs. As a byproduct, we also show that, unlike B_k -EPG graphs, B_k -ENPG graphs do not necessarily admit a representation where every path has exactly k bends. We summarize and point to further research directions in Section 5.

2 Preliminaries

Given a simple graph (no loops or parallel edges) $G = (V(G), E(G))$ and a vertex v of G , we denote by $N_G(v)$ the set of neighbors of v in G , and by $d_G(v) = |N_G(v)|$ the degree of v in G . A graph is called d -regular if every vertex v has $d(v) = d$. Whenever there is no ambiguity we omit the subscript G and write $d(v)$ and $N(v)$. Given a graph G and $U \subseteq V(G)$, $N_U(v) \stackrel{def}{=} N_G(v) \cap U$. Two adjacent (resp. non-adjacent) vertices u, v of G are *twins* (resp. *false twins*) if $N_G(u) \setminus \{v\} = N_G(v) \setminus \{u\}$. For a graph G and $U \subseteq V(G)$, we denote by $G[U]$ the subgraph of G induced by U .

A vertex set $U \subseteq V(G)$ is a clique (resp. stable set) (of G) if every pair of vertices in U is adjacent (resp. non-adjacent). A graph G is a *split graph* if $V(G)$ can be partitioned into a clique and a stable set. A graph G is *co-bipartite* if $V(G)$ can be partitioned into two cliques. Note that these partitions are not necessarily unique. We denote bipartite, co-bipartite and split graphs as $X(V_1, V_2, E)$ where

- a) $X = B$ (resp. C, S) whenever G is bipartite (resp. co-bipartite, split),
- b) $V_1 \cap V_2 = \emptyset, V_1 \cup V_2 = V(G)$,
- c) for bipartite graphs V_1, V_2 are stable sets,
- d) for co-bipartite graphs V_1 and V_2 are cliques,
- e) for split graphs V_1 is a clique and V_2 is a stable set, and
- f) $E \subseteq V_1 \times V_2$ (in other words E does not contain the cliques' edges).

Unless otherwise stated we assume that G is connected and both V_1 and V_2 are non-empty.

In this work every single path is simple, i.e. without duplicate vertices. However, if a union of paths is a path, the resulting path is not necessarily simple. For example, consider a graph on 5 vertices v_1, v_2, v_3, v_4, v_5 and 5 edges $e_1 = v_1v_2, e_2 = v_2v_3, e_3 = v_3v_4, e_4 = v_4v_2, e_5 = v_2v_5$. Each of the

paths $P_1 = \{e_1e_2e_3\}$ and $P_2 = \{e_3, e_4, e_5\}$ is simple. On the other hand, though $P_1 \cup P_2$ is a path, it is not simple. Whenever v is an internal vertex of a path P , we sometimes say that P *crosses* v . Given two paths P, P' , a *split* of P, P' is a vertex with degree at least 3 in $P \cup P'$. We denote by $split(P, P')$ the set of all splits of P and P' . When $split(P, P') \neq \emptyset$ we say that P and P' are *splitting*. Whenever P and P' edge intersect and $split(P, P') = \emptyset$ we say that P and P' are *non-splitting* and denote this by $P \sim P'$. Clearly, for any two paths P and P' exactly one of the following holds:

- i) P and P' are edge disjoint,
- ii) P and P' are splitting,
- iii) $P \sim P'$.

A two-dimensional *grid graph*, also known as a square grid graph, is an $m \times n$ lattice graph that is the Cartesian product graph of two paths P and P' of respectively length n and m . Such a grid has vertex set $V = [n] \times [m]$. A *bend* of a path P in a grid H is an internal vertex of P whose incident edges (in the path) have different directions, i.e. one vertical and one horizontal.

Let \mathcal{P} be a set of paths in a graph H . The graphs $EP(\mathcal{P})$ and $ENP(\mathcal{P})$ are such that $V(ENP(\mathcal{P})) = V(EP(\mathcal{P})) = V$, and there is a one-to-one correspondence between \mathcal{P} and V , i.e. $\mathcal{P} = \{P_v : v \in V\}$. Given two paths $P_u, P_v \in \mathcal{P}$, $\{u, v\}$ is an edge of $EP(\mathcal{P})$ if and only if P_u and P_v have a common edge (cases (ii) and (iii)), whereas $\{u, v\}$ is an edge of $ENP(\mathcal{P})$ if and only if $P_u \sim P_v$ (case (iii)). Clearly, $E(ENP(\mathcal{P})) \subseteq E(EP(\mathcal{P}))$. A graph G is ENP if there is a graph H and a set of paths \mathcal{P} of H such that $G = ENP(\mathcal{P})$. In this case $\langle H, \mathcal{P} \rangle$ is an ENP *representation* of G . When H is a tree (resp. grid) $EP(\mathcal{P})$ is an EPT (resp. EPG) graph, and $ENP(\mathcal{P})$ is an ENPT (resp. ENPG) graph; these graphs are denoted also as $EPT(\mathcal{P})$, $EPG(\mathcal{P})$, $ENPT(\mathcal{P})$ and $ENPG(\mathcal{P})$, respectively. We say that two representations are *equivalent* if they are representations of the same graph.

Let $\langle H, \mathcal{P} \rangle$ be a representation of an ENP graph G . For each edge e of H , \mathcal{P}_e denotes the set of the paths of \mathcal{P} containing the edge e , i.e. $\mathcal{P}_e \stackrel{def}{=} \{P \in \mathcal{P} \mid e \in P\}$. For a subset $U \subseteq V(G)$ we define $\mathcal{P}_U \stackrel{def}{=} \{P_v \in \mathcal{P} : v \in U\}$. Following standard notations, $\cup \mathcal{P}_U \stackrel{def}{=} \cup_{P \in \mathcal{P}_U} P$.

Given two paths P and P' of a graph, a *segment* of $P \cap P'$ is a maximal path that constitutes a sub-path of both P and P' . Clearly, $P \cap P'$ is the union of edge disjoint segments. We denote the set of these segments by $\mathcal{S}(P, P')$.

Throughout the paper, whenever a representation $\langle H, \mathcal{P} \rangle$ of an ENPG graph is given, we assume the host graph H is a grid on sufficiently many vertices each of which is denoted by an ordered pair of integers.

The following Proposition that is proven in Boyacı et al. (2015b) is the starting point of many of our results.

Proposition 2.1 *Boyacı et al. (2015b) Let K be a clique of a B_1 -ENPG graph G with a representation $\langle H, \mathcal{P} \rangle$. Then $\cup \mathcal{P}_K$ is a path with at most 2 bends. Moreover, there is an edge $e_K \in E(H)$ such that every path of \mathcal{P}_K contains e_K .*

Note that whenever $\cup \mathcal{P}_K$ has two bends, e_K lies between these two bends. Based on the above proposition, given two cliques K, K' of a B_1 -ENPG graph we denote $\mathcal{S}(K, K') \stackrel{def}{=} \mathcal{S}(\cup \mathcal{P}_K, \cup \mathcal{P}_{K'})$.

By the following two observations, in the sequel we focus on connected twin-free graphs.

Observation 2.1 *Let G be a graph and G' obtained from G by removing a twin vertex until no twins remain. Then, G is B_k -ENPG if and only if G' is B_k -ENPG.*




Fig. 1: (a) A B_1 -EPG representation of C_4 , (b) A B_1 -EPG representation of C_{11} .

Observation 2.2 A graph G is B_k -ENPG if and only if every connected component of G is B_k -ENPG.

We first observe that some well-known graph classes are included in B_1 -ENPG.

Proposition 2.2

- i) Every cycle is B_1 -ENPG.
- ii) Every tree is B_1 -ENPG, and it has a representation $\langle H, \mathcal{P} \rangle$ where $\cup \mathcal{P}$ is a tree.

Proof:

- i) For $k = 3$ three identical paths consisting of one edge constitutes a B_1 -ENPG representation of C_3 . For $k = 4$ Figure 1 (a) depicts a B_1 -ENPG representation of C_4 . Finally for any $k > 4$, we can construct a C_k as shown in Figure 1 (b) for the case $k = 11$.
- ii) Given a representation $\langle H, \mathcal{P} \rangle$ of a B_1 -ENPG graph G and $U \subseteq V(G)$, we denote by R_U the bounding rectangle of \mathcal{P}_U . Let T be a tree with a root r . We prove the following claim by induction on the structure of T (see Figure 2). The tree T has a B_1 -ENPG representation $\langle H, \mathcal{P} \rangle$ in which the corners of the bounding rectangle R_T can be renamed as a_T, b_T, c_T, d_T in counterclockwise order such that i) every path of \mathcal{P} has exactly one bend, ii) b_T is a bend of P_r , iii) a_T is an endpoint of P_r , iv) the line between a_T and d_T is used exclusively by P_r , v) $\cup \mathcal{P}$ is a tree.

If T is an isolated vertex, any path with one bend is a representation of T . Moreover, it is easy to verify that it satisfies conditions i) through v).

Otherwise let T_1, \dots, T_k be the subtrees of T obtained by the removal of r , with roots r_1, \dots, r_k respectively. By the inductive hypothesis every such subtree T_i has a representation with bounding box $a_{T_i}, b_{T_i}, c_{T_i}, d_{T_i}$ satisfying conditions i) through iv). We now build a representation of T satisfying the same conditions. We shift and rotate the representations of T_1, \dots, T_k so that the bounding rectangles do not intersect and the vertices $a_{T_1}, b_{T_1}, a_{T_2}, b_{T_2}, \dots, a_{T_k}, b_{T_k}$ are on the same horizontal line and in this order (See Figure 2). We extend the paths P_{r_2}, \dots, P_{r_k} representing the roots of the trees T_2, \dots, T_k such that the endpoint a_{T_i} of P_{r_i} is moved to a_{T_1} .

Since a_{T_i} is used exclusively by P_{r_i} this modification does not cause P_{r_i} to split from a path of $\mathcal{P}_{V(T_i)}$. Therefore, the individual trees T_1, \dots, T_K are properly represented. Clearly, if two paths from different subtrees T_i, T_j ($i < j$) intersect, then one of the intersecting paths must be P_{r_j} . The path P_{r_j} intersects the bounding rectangle of T_i only at the path between a_i and b_i . As every path of $\mathcal{P}_{V(T_i)}$, in particular one intersecting P_{r_j} has one bend, such a path splits from P_{r_j} . Therefore,




Fig. 2: A construction for B_1 -ENPG representation of trees.

for any pair of vertices $(v_i, v_j) \in T_i \times T_j$ we have that v_i and v_j are non-adjacent in $\text{ENPG}(\mathcal{P})$, as required.

We rename the corners of the bounding rectangle R_T such that $b_T = a_{T_1}$. We now add the path P_r from b_{T_1} to a_T with a bend at b_T . The conditions i), ii), iii) are satisfied. We extend P_r by one edge at a_T to make sure that the line between a_T and d_T is exclusively used by P_r , thus satisfying condition iv). The extension of the paths P_{r_2}, \dots, P_{r_k} does not add new edges to regions bounded by $a_{T_i}, b_{T_i}, c_{T_i}, d_{T_i}$ and they don't introduce cycles between this regions. Moreover, since the line between a_{T_1} and d_{T_1} is used only at a_{T_1} the path P_r does not introduce any cycles either. Therefore, $\cup \mathcal{P}$ is a tree, i.e. condition v) is satisfied.

The path P_r intersects only R_{T_1} . This intersection is the path between b_{T_1} and d_{T_1} bending at a_{T_1} . Every path that intersects P_r and does not split from it must bend at a_{T_1} . As a_{T_1} is used exclusively by P_{r_1} , P_{r_1} is the only path that possibly satisfies $P_{r_1} \sim P_r$. We now observe that $P_{r_i} \sim P_r$ for every $i \in [k]$. Therefore r is adjacent to the root of T_j in $\text{ENPG}(\mathcal{P})$, as required.

□

3 Split Graphs

In this section, we present a characterization theorem (Theorem 3.1) for B_1 -ENPG split graphs. In Sections 3.1 and 3.2 we proceed with some properties of these graphs implied by this theorem. An interesting implication of one of these properties is that the family of B_1 -ENPG is properly included in the family of B_2 -ENPG graphs. Finally, using Theorem 3.1, we prove in Section 3.3 that the recognition problem of B_1 -ENPG graphs is NP-complete even in a very restricted subfamily of split graphs. Throughout this section, G is a split graph $S(K, S, E)$ unless indicated otherwise. We assume without loss of generality that K is maximal, i.e. that no vertex in S is adjacent to all vertices of K , and G is connected (in particular S does not contain isolated vertices).

3.1 Characterization of B_1 -ENPG Split Graphs

Consult Figure 3 for the following discussion.




Fig. 3: Regions of $\cup \mathcal{P}_K$ and possible representations of a vertex in S for a B_1 -ENPG split graph $S(K, S, E)$.

Let G be a B_1 -ENPG split graph $S(K, S, E)$ with a representation $\langle H, \mathcal{P} \rangle$. By Proposition 2.1, we know that $\cup \mathcal{P}_K$ is a path with at most two bends, such that there is an edge e_K contained in every path of \mathcal{P}_K . Moreover, if $\cup \mathcal{P}_K$ contains two bends then e_K is between the two bends. Assume without loss of generality that e_K is a horizontal edge. Therefore, $\cup \mathcal{P}_K$ consists of a horizontal segment L_H between two vertices b_L, b_R and two vertical segments (each of which is possibly empty). The subgrid $L_H \setminus e_K$ consists of two horizontal subsegments $L_{L,H}, L_{R,H}$. Finally, $\cup \mathcal{P}_K \setminus L_H$ consists of two vertical segments $L_{L,V}$ and $L_{R,V}$. The segment $L_{L,Y}$ (resp. $L_{R,Y}$) is on the left (resp. right) of e_K for every $Y \in \{H, V\}$.

For $(X, Y) \in \{L, R\} \times \{H, V\}$, let $K_{X,Y}$ be the set of vertices v of K such that P_v has an endpoint in $L_{X,Y}$. Every path of \mathcal{P}_K has its left (resp. right) endpoint on $L_{L,H} \cup L_{L,V}$ (resp. $L_{R,H} \cup L_{R,V}$) since it contains e_K . Therefore, $\{K_{X,H}, K_{X,V}\}$ is a partition of K for every $X \in \{L, R\}$. For every $(X, Y) \in \{L, R\} \times \{H, V\}$, let $\sigma_{X,Y}$ be the permutation of $K_{X,Y}$ obtained by ordering the endpoints of \mathcal{P}_K in $L_{X,Y}$ in increasing distance from e_K . Moreover, $K_{L,V} \cap K_{R,V} = \emptyset$ since otherwise this implies a path containing both b_L and b_R as bends.

The following theorem characterizes the B_1 -ENPG split graphs. It further provides a canonical representation for them using the above mentioned partitions and by partitioning the vertices of S according to their neighborhoods.

Theorem 3.1 *A connected split graph $G = S(K, S, E)$ is B_1 -ENPG if and only if there are two partitions $\{K_{L,H}, K_{L,V}\}, \{K_{R,H}, K_{R,V}\}$ of K such that $K_{L,V} \cap K_{R,V} = \emptyset$, there is a permutation $\sigma_{X,Y}$ of $K_{X,Y}$ for every $(X, Y) \in \{L, R\} \times \{H, V\}$, and a partition $\mathcal{S} = \{S_{X,H}, S_{X,V}, S_{X,HH}, S_{X,HV} \mid X \in \{L, R\}\}$ of S such that the following hold.*

- i) If $s \in S_{X,Y}$ then $N(s)$ is an interval σ_s of $\sigma_{X,Y}$.
- ii) If $s \in S_{X,HH}$ then $N(s)$ consists of the intersection of a prefix σ_s of $\sigma_{X,H}$ with $K_{\bar{X},H}$ where $\bar{X} = \{L, R\} \setminus X$.
- iii) If $s \in S_{X,HV}$ then $N(s)$ is the union of a suffix σ_s of $\sigma_{X,H}$ with $K_{X,V}$.

iv) If $s \in S_{X,H} \cup S_{X,HH}$ then there is at most one $s' \in S_{X,HV}$ such that the interval σ_s of $\sigma_{X,H}$ and the suffix $\sigma_{s'}$ of $\sigma_{X,H}$ overlap.

v) If $S_{X,HH} \neq \emptyset$ then $|S_{\bar{X},HV}| \leq 1$ where $\bar{X} = \{L, R\} \setminus X$.

vi) $K_{X,V} \subseteq K_{\bar{X},H}$ where $\bar{X} = \{L, R\} \setminus X$.

Proof: (\Rightarrow) We fix a B_1 -ENPG representation of G and consider the sets $K_{X,Y}$ and their permutations $\sigma_{X,Y}$ defined by this representation.

i, ii iii) For each vertex $s \in S$ we will determine its membership to one of the sets of the partition \mathcal{S} depending on its representation P_s . Suppose that there exists a vertex $s \in S$ such that $|\mathcal{S}(P_s, \cup \mathcal{P}_K)| > 1$. Then $P_s \cup \cup \mathcal{P}_K$ contains a cycle, therefore at least 4 bends. But P_s has at most one bend and $\cup \mathcal{P}_K$ has at most two bends, a contradiction. Therefore, $\mathcal{S}(P_s, \cup \mathcal{P}_K)$ consists of one segment Q_s .

Given a vertex $s \in S$, we consider two disjoint and complementary cases for P_s .

1. $e_K \notin Q_s$: Let c_s (resp. f_s) be the vertex of Q_s closer to (resp. farther from) e_K . Assume without loss of generality that $Q_s \subseteq L_L$ where $L_L = L_{L,H} \cup L_{L,V}$ (we define similarly $L_R = L_{R,H} \cup L_{R,V}$). We observe that P_s does not split from L_L at c_s , since otherwise P_s splits from every path of \mathcal{P}_K that it intersects, implying that s is an isolated vertex. We further consider three subcases:
 - (a) $Q_s \subseteq L_{L,V}$: If P_s splits from L_L then $N(s)$ consists of the vertices v of K such that the left endpoint of P_v is between c_s and f_s . Therefore, $N(s)$ is an interval of $\sigma_{L,V}$. If P_s does not split from L_L then $N(s)$ consists of the vertices v of K such that the left endpoint of P_v is farther than c_s on $L_{L,V}$ (with respect to b_L). Therefore, $N(s)$ is an interval of $\sigma_{L,V}$. In both cases we set $s \in S_{L,V}$.
 - (b) b_l is an internal vertex of Q_s : In this case P_s does not split from L_L . Then $N(s)$ consists of the vertices v of K such that the left endpoint of P_v is on the left of c_s on L_L . Therefore, $N(s)$ consists of the union of a suffix of $\sigma_{L,H}$ with $K_{L,V}$, and we set $s \in S_{L,HV}$.
 - (c) $Q_s \subseteq L_{L,H}$:
 - i. P_s splits from L_L . In this case $N(s)$ consists of the vertices v of K such that the left endpoint of P_v is between c_s and f_s . Therefore, $N(s)$ is an interval of $\sigma_{L,H}$ and we set $s \in S_{L,H}$.
 - ii. P_s does not split from L_L . In this case $N(s)$ consists of the vertices v of K such that the left endpoint of P_v is on the left of c_s on L_L . Therefore, $N(s)$ consists of the union of a suffix of $\sigma_{L,H}$ with $K_{L,V}$ and we set $s \in S_{L,HV}$.
2. $e_K \in Q_s$: In this case, let l_s, r_s be the endpoints of Q_s on L_L and L_R respectively. The path P_s splits from $\cup \mathcal{P}_K$, since otherwise s is adjacent to every vertex of the clique, contradicting the maximality of K . Since P_s intersects every path of \mathcal{P}_K , $N(s)$ consists of the vertices v of K such that P_v does not split from P_s . We consider two subcases:
 - (a) P_s splits from exactly one of L_L and L_R : Assume without loss of generality that P_s splits from L_L but not from L_R . We observe that $l_s \in L_{L,H}$. In this case $N(s)$ is the set of vertices v of K such that the left endpoint of P_v is closer than l_s to e_K which corresponds to a prefix of $\sigma_{L,H}$. In this case we set $s \in S_{L,H}$.

- (b) P_s splits from both of L_L, L_R : In this case at least one of the endpoints of Q_s is a bend of $\cup \mathcal{P}_K$, i.e., $\{l_s, r_s\} \cap \{b_L, b_R\} \neq \emptyset$. Assume without loss of generality that $r_s = b_R$. Then $N(s)$ consists of those vertices v of K such that the left endpoint of P_v is closer to e_K than l_s and the right endpoint of P_v is in $L_{R,H}$. This is exactly a prefix of $\sigma_{L,H}$ intersected with $K_{R,H}$, thus we set $s \in S_{L,HH}$.

iv) Assume, for a contradiction that for some $X \in \{L, R\}$, say L , the condition does not hold, i.e. there is a vertex $s \in S_{L,H} \cup S_{L,HH}$ and two vertices $s', s'' \in S_{L,HV}$ such that the interval σ_s of $\sigma_{L,H}$ corresponding to s overlaps both of the suffixes $\sigma_{s'}, \sigma_{s''}$ corresponding to s', s'' respectively. By the above case analysis we know that P_s is either of type 2 or of type 1.c.i, and that $P_{s'}$ and $P_{s''}$ are of one of the types 1.b, 1.c.ii. Note that $\sigma_{s'}$ and $\sigma_{s''}$ are determined by the right endpoints of the corresponding paths $P_{s'}$ and $P_{s''}$. Since $\sigma_{s'}$ and σ_s overlap, Q_s contains the right endpoint of $P_{s'}$. Therefore these two paths intersect. Moreover $P_{s'}$ contains the left endpoint of Q_s (which is the bend point of P_s) otherwise $P_s \sim P_{s'}$. By the same arguments $P_{s''}$ also contains the left endpoint of Q_s and therefore $P_{s'}$ intersects $P_{s''}$. Moreover, since none of them splits from L_L we have $P_{s'} \sim P_{s''}$, i.e. s and s' are adjacent in G , a contradiction.

v) Assume, for a contradiction that for some $X \in \{L, R\}$, say L , the condition does not hold, i.e. there exists $s \in S_{R,HH}$ and $s', s'' \in S_{L,HV}$. Then both $P_{s'}$ and $P_{s''}$ are of one of the types 1.b, 1.c.ii. Moreover, P_s is of type 2.b with $l_s = b_L$. We proceed as in the previous case to get a contradiction.

vi) This immediately follow from the already observed fact that $K_{L,V} \cap K_{R,V} = \emptyset$.

(\Leftarrow) Suppose that the partitions and the permutations stated in the claim exist. We construct a representation $\langle H, \mathcal{P} \rangle$ as follows (see Figure 4). The host graph H is a $(2 + 4|S|(|K| + 1))$ by $2|S| \cdot |K|$ vertices grid, where each vertex is represented by an ordered pair from $[-1 - 2|S|(|K| + 1), 1 + 2|S|(|K| + 1)] \times [0, 2|S||K|]$. For $(X, Y) \in \{L, R\} \times \{H, V\}$, let $k_{X,Y} = |K_{X,Y}|$. The coordinates of b_L and b_R are respectively $(-1 - 2|S|(k_{L,H} + 1), 0)$ and $(1 + 2|S|(k_{L,H} + 1), 0)$. The horizontal line between b_L and b_R is called L_H . For $(X, Y) \in \{L, R\}$, $L_{X,V}$ is a vertical line of length $2|S|(k_{X,V})$ starting at b_X . We choose $k_{X,Y}$ vertices on each line $L_{X,Y}$ such that their distances from each other and from each of $b_L, b_R, (-1, 0), (1, 0)$ is at least $2|S|$. We label these vertices as $w_{X,\sigma_{X,Y}(1)}, \dots, w_{X,\sigma_{X,Y}(k_{X,Y})}$ in increasing order of their distances from the origin. Every vertex $v \in K$ is represented by a path $P_v \subseteq \cup L_H \cup L_{L,V} \cup L_{R,V}$ between $w_{L,v}$ and $w_{R,v}$. Since $K_{X,V} \subseteq K_{\bar{X},H}$ every such path has at most one bend. Since $e_K = (-1, 0)(1, 0)$ is contained in every such path, these paths constitute a proper representation of the clique K .

We proceed with the representation of the vertices of S . Let $W_X = \{w_{X,1}, \dots, w_{X,|K|}\}$. The endpoints of paths $Q_s, s \in S$ will be chosen between two vertices of $W \cup \{b_X\}$ so that they are all distinct. We first determine the representations of the vertices of $S \setminus \{S_{L,HV} \cup S_{R,HV}\}$ such that all are one bend paths with distinct bends on the endpoint of Q_s farther from the origin. We determine the endpoints of Q_s according to the permutation σ_s and as close to the origin as possible. Since all these paths have distinct bends, they represent an independent set. Last, we represent the vertices $s' \in \{S_{L,HV} \cup S_{R,HV}\}$. For each such vertex its representation will be a path $P_{s'} \subseteq \cup \mathcal{P}_K$. We choose the endpoint closer to the origin according to the suffix $\sigma_{s'}$. Let $O_{s'}$ be the set of vertices such that for all vertices $s \in O_{s'}$, σ_s and $\sigma_{s'}$ overlap. The other endpoint is chosen as the endpoint closest to the origin that is farther from the origin than all the endpoints of the paths P_s where $s \in O_{s'}$. Conditions iv and v guarantee that after this is done, every path $P_{s'}$ that intersects with P_s splits from it. \square

From the proof of Theorem 3.1 we obtain the following corollary where a *caterpillar* is a tree in which




Fig. 4: Construction of a representation $\langle H, \mathcal{P} \rangle$ for a split graph satisfying conditions i)–vi) of Theorem 3.1.

all vertices are within distance 1 of a central path.

Corollary 3.1

- i) Every connected B_1 -ENPG split graph $G = S(K, S, E)$ has a representation $\langle H, \mathcal{P} \rangle$ such that $\cup \mathcal{P}$ is a caterpillar with central path $\cup \mathcal{P}_K$ and maximum degree 3.
- ii) B_1 -ENPG \cap SPLIT \subseteq ENPT \cap SPLIT.

3.2 Two Consequences of The Characterization of B_1 -ENPG Split Graphs

Throughout this section, we use the notation introduced in the previous section. X (resp. Y) denotes an element of $\{L, R\}$ (resp. $\{H, V\}$), and $\bar{X} = \{L, R\} \setminus \{X\}$. Given a B_1 -ENPG split graph $G = S(K, S, E)$, we denote by $K_{X,Y}, S_{X,Y}, S_{X,HV}, S_{X,HH}$ and $\sigma_{X,Y}$ the sets and permutations whose existence are guaranteed by Theorem 3.1. Furthermore $\sigma_X = \sigma_{X,H} \cdot \sigma_{X,V}$ is the permutation of K obtained by the concatenation of the two permutations $\sigma_{X,H}$ and $\sigma_{X,V}$. We define $S_{X,Y,d}$ as the set of vertices of $S_{X,Y}$ having degree d in G . The notations $S_{X,HV,d}$, and $S_{X,HH,d}$ are defined similarly.

The following inequalities are easy to show using the definitions of the sets and counting the number of prefixes, suffixes, or intervals of a given permutation having a given length.

Proposition 3.1 *If $G = S(K, S, E)$ is a twin-free and (false twin)-free B_1 -ENPG split graph, then*

$$|S_{X,Y,d}| \leq \max\{|K_{X,Y}| + 1 - d, 0\} \quad (1)$$

$$|S_{X,HV,d}| \leq \begin{cases} 1 & \text{if } d > |K_{X,V}| \\ 0 & \text{otherwise} \end{cases} \quad (2)$$

$$|S_{X,V,d} \cup S_{X,HV,d}| \leq \begin{cases} 1 & \text{if } d > |K_{X,V}| \\ |K_{X,V}| + 1 - d & \text{otherwise} \end{cases} \quad (3)$$

$$|S_{X,HH,d}| \leq \begin{cases} 1 & \text{if } d \leq |K_{L,H} \cap K_{R,H}| \leq 1 \\ 0 & \text{otherwise} \end{cases} \quad (4)$$

$$|S_d| \leq 2(|K| + 2 - d). \quad (5)$$

Proof: (1) If $s \in S_{X,Y,d}$ then $N(s)$ is an interval σ_s of $\sigma_{X,Y}$ of size d . Since G is (false twin)-free $N(s) \neq N(s')$ whenever $s \neq s'$. Therefore, $|S_{X,Y,d}|$ is at most the number of such intervals which is given by the right hand side of the inequality.

(2) If $s \in S_{X,HV,d}$ then $N(s) = K_{X,V} \cup \sigma_s$ where σ_s is a suffix of $\sigma_{X,H}$. Therefore, $d > |K_{X,V}|$ and σ_s is the unique suffix of $\sigma_{X,H}$ of size $d - |K_{X,V}|$.

(3) Follows from (1) and (2).

(4) If $s \in S_{X,HH,d}$ then $N(s) = K_{\bar{X},H} \cap \sigma_s$ where σ_s is a prefix of $\sigma_{X,H}$ of size d . If $d > |K_{L,H} \cap K_{R,H}|$ no such prefix exists, otherwise there is exactly one such prefix.

(5) By summing up (2), (3), and also (1) for $Y = H$ and finally multiplying by two for the two possible values of X . \square

Summing up (5) for all the possible values of $d \in [|K|]$ we get the following corollary.

Corollary 3.2 *If $G = S(K, S, E)$ is a (false twin)-free B_1 -ENPG split graph, then $|S|$ is $\mathcal{O}(|K|^2)$.*

Using similar arguments one can show that if $G = S(K, S, E)$ is twin-free then $|K|$ is $\mathcal{O}(|S|^2)$ implying that $|S|$ is $\Omega(\sqrt{|K|})$. More specifically, one should consider the set of endpoints or bend points of the paths \mathcal{P}_S all of which are in $\cup \mathcal{P}_K$ and observe that no four such points that are pairwise consecutive in $\cup \mathcal{P}_K$ may surround the endpoints of two paths $P_u, P_v \in \mathcal{P}_K$ since otherwise u and v are twins.

Theorem 3.2 *The following strict inclusions hold:*

- $B_1\text{-ENPG} \cap \text{SPLIT} \subsetneq B_2\text{-ENPG} \cap \text{SPLIT}$.
- $B_1\text{-ENPG} \cap \text{SPLIT} \subsetneq \text{ENPT} \cap \text{SPLIT}$.

Proof: By the definition of a B_k -ENPG graph, we have $B_1\text{-ENPG} \cap \text{SPLIT} \subseteq B_2\text{-ENPG} \cap \text{SPLIT}$ and by Corollary 3.1, we have $B_1\text{-ENPG} \cap \text{SPLIT} \subseteq \text{ENPT} \cap \text{SPLIT}$. In the following, we provide a split graph with a representation which is both B_2 -ENPG and ENPT. We show that this split graph is not B_1 -ENPG.

Let $K = [0, 10]$, σ_L be the identity permutation $(0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10)$ on K and σ_R be the permutation $(0, 5, 10, 4, 9, 3, 8, 2, 7, 1, 6)$ on K . Let $G = S(K, S, E)$ where S contains 23 vertices: one for every pair that is consecutive in one of σ_L, σ_R (there are 10 in every permutation and we note that these pairs are distinct) and one for each of the pairs $\{0, 2\}, \{0, 3\}, \{0, 4\}$ (which are not consecutive in




Fig. 5: The B_2 -ENPG representation of a non- B_1 -ENPG split graph used in the proof of Theorem 3.2. The paths representing vertices of K are not drawn. They are implied by the numbers of the vertices: for every $i \in [0, 10]$, P_i is the shortest path between the two vertices labeled i . The paths with two bends, intersects every path of \mathcal{P}_K , but splits from every path having an endpoint after its bend points. Therefore, these path represent vertices with neighborhood $\{0, 2\}$, $\{0, 3\}$, $\{0, 4\}$.

any of these permutations). Every vertex in S is adjacent to the corresponding pair in K . Note that G is (false-twin)-free and $|S_2| = |S| = 23 > 22 = 2(|K| + 2 - 2)$. By Proposition 3.1 (5), G is not B_1 -ENPG. Figure 5 depicts a set of paths that constitute a B_2 -ENPG representation and an ENPT representation of G . \square

3.3 NP-completeness of B_1 -ENPG split graph recognition

We now proceed with the NP-completeness of B_1 -ENPG recognition in split graphs. We first present a preliminary result that can be useful per se.

A graph is d -regular if all its vertices have degree d . A 3-regular graph is also termed *cubic*. A *diamond* is the graph $K_4 - e$ obtained by removing an edge from a clique on four vertices. Clearly, if the edge set of a graph G can be partitioned into two Hamiltonian cycles, then G is 4-regular. However, in the opposite direction we have the following:

Theorem 3.3 *The problem of determining whether the edge set of a diamond-free 4-regular graph can be partitioned into two Hamiltonian cycles is NP-complete.*

Proof: We prove by reduction from the Hamiltonicity problem of cubic bipartite graphs which is known to be NP-complete (Akiyama et al. (1980)). Let G be a cubic bipartite graph whose Hamiltonicity has to be decided, and let $H = L(G)$ be its line graph. H is clearly 4-regular. In the sequel we will show that H is also diamond-free. In addition, we know that G is Hamiltonian if and only if the edge set of its line graph H can be partitioned into two Hamiltonian cycles (Kotzig (1957)). This concludes the proof. It remains to show that H is diamond-free.

Suppose for a contradiction that H contains a diamond on vertices $\{e_1, e_2, e_3, e_4\}$ that are pairwise adjacent except for the pair e_1, e_4 . Then $\{e_1, e_2, e_3\}$ and $\{e_2, e_3, e_4\}$ are two triangles of H . Every triangle of H corresponds to either a triangle of G , or to three edges of G incident to a common vertex. Since G is bipartite, only the latter case is possible. Then e_1, e_2, e_3 (resp. e_2, e_3, e_4) are edges of G incident to a vertex v (resp. v'). Since G is cubic we have $v \neq v'$. We conclude that $e_2 = e_3 = vv'$, a contradiction. \square

We are now ready to prove the main result of this section.

Theorem 3.4 *The recognition problem of B_1 -ENPG graphs is NP-complete even when restricted to split graphs.*

Proof: The proof is by reduction from the problem of decomposing a 4-regular, diamond-free graph into two Hamiltonian cycles which is shown to be NP-complete in Theorem 3.3. Given a 4-regular graph G on $n + 1$ vertices, we remove an arbitrary vertex v of G and obtain the graph $G' = G - v$ on n vertices all of which having degree 4, except the four neighbours $\{v_1, v_2, v_3, v_4\}$ of v each of which having degree 3. We construct the split graph $G'' = S(K, S, E)$ where $K = V(G')$, $S = E(G') \cup \{s_1, s_2, s_3, s_4\}$. Furthermore, the neighborhood of a vertex $s \in S$ is determined as follows. If $s = s_i$ for some $i \in [4]$ then $N_{G''}(s) = K - v_i$, otherwise s is an edge uv of G' in which case $N_{G''}(s) = \{u, v\}$. It remains to show that G'' is B_1 -ENPG if and only if $E(G)$ can be partitioned into two Hamiltonian cycles.

Assume that $E(G)$ can be partitioned into two Hamiltonian cycles C_L, C_R . This induces a partition of $E(G')$ into two paths Q_L and Q_R which in turn induces a partition of $S - \{s_1, s_2, s_3, s_4\}$ into $S_{L,H} = E(Q_L)$ and $S_{R,H} = E(Q_R)$. Note that the endpoints of Q_L and Q_R are the degree 3 vertices of G' , i.e. $\{v_1, v_2, v_3, v_4\}$. Let without loss of generality v_1, v_2 (resp. v_3, v_4) be the endpoints of Q_L (resp. Q_R). For $X \in \{L, R\}$ let $K_{X,H} = K = V(G')$ and $K_{X,V} = \emptyset$. We set $\sigma_{X,H}$ as the order of the vertices of G' in Q_X (which is a permutation of the vertices of $K = V(G')$). Then v_1 and v_2 (resp. v_3 and v_4) are the first and last vertices of the permutation $\sigma_{L,H}$ (resp. $\sigma_{R,H}$). For $X \in \{L, R\}$ we set $S_{X,V} = S_{X,HH} = S_{X,HV} = \emptyset$. We now verify that these settings satisfy the conditions of Theorem 3.1. Conditions ii), iii), iv), v) and vi) easily follow since the sets $S_{X,V}, S_{X,HV}, S_{X,HH}$ and $K_{X,V}$ are empty. As for Condition i) we consider two cases. If $s = s_i$ for some $i \in [4]$ then $N_{G''}(s) = K - v_i$ is an interval of $\sigma_{X,H}$ for some $X \in \{L, R\}$ since v_i is either the first or the last vertex of one of these permutations. If s is an edge $uv \in E(Q_X)$ of G' then u and v are consecutive in the permutation $\sigma_{X,H}$. Since all the conditions are satisfied, we conclude that G'' is B_1 -ENPG.

Now assume that G'' is B_1 -ENPG. For $X \in \{L, R\}, Y \in \{H, V\}$, let $K_{X,Y}, \sigma_{X,Y}, S_{X,Y}, S_{X,HV}$ and $S_{X,HH}$ be sets and permutations whose existence are guaranteed by Theorem 3.1. We first show that $|K_{X,V}| \leq 1$. Assume for a contradiction that $|K_{X,V}| > 1$ for some $X \in \{L, R\}$, say $X = L$. Then we have $|K_{R,H}| > 1$, and $|K_{L,H}|, |K_{R,V}| < n - 1$. By Proposition 3.1, these imply $S_{L,H,n-1} = S_{L,HH,n-1} = S_{R,HH,n-1} = \emptyset$. Moreover, $|S_{R,H,n-1}| \leq 2$ and this may hold with equality only when $K_{R,H} = K$. Finally, we have $|S_{X,V,n-1} \cup S_{X,HV,n-1}| \leq 1$. Summing up all inequalities we obtain $|S_{n-1}| \leq 4$. We recall that all the vertices of S have degree 2 except the four special vertices with degree $n - 1$. Therefore, $S_{n-1} = \{s_1, s_2, s_3, s_4\}$, and we conclude that all the inequalities hold with equality. In particular $|S_{R,H,n-1}| = 2$, implying $K_{R,H} = K$ and $K_{R,V} = \emptyset$. Then we have $S_{R,V,n-1} \cup S_{R,HV,n-1} = \emptyset$, i.e. one of the inequalities is strict, a contradiction. Therefore, $|K_{X,V}| \leq 1$, implying

$$S_{X,V,2} = \emptyset. \quad (6)$$

Recall that $\sigma_X = \sigma_{X,H} \cdot \sigma_{X,V}$. We now show that the set of the first and last vertices of σ_L and σ_R is $\{v_1, v_2, v_3, v_4\}$. Let $i \in [4]$ and consider each one of the cases $s_i \in S_{X,H}, s_i \in S_{X,HH}$ and $s_i \in S_{X,HV}$ (the case $s_i \in S_{X,V}$ is impossible since $|K_{X,V}| \leq 1$). It is easy to verify for every case that v_i is either the first or the last vertex of σ_X . By the pigeonhole principle we conclude that the set of first and last vertices of σ_L and σ_R is $\{v_1, v_2, v_3, v_4\}$. We assume without loss of generality that v_1 (resp. v_2) is the first (resp. last) vertex of σ_L and that v_3 (resp. v_4) is the first (resp. last) vertex of σ_R .

Our next step is to show that $S_{X,HH,2} = \emptyset$. Assume for a contradiction that this does not hold, and let s be a vertex (without loss of generality) of $S_{L,HH,2}$. Then σ_s is a prefix with two vertices of




Fig. 6: Two path sets $\mathcal{P}_u, \mathcal{P}_v$ meet at a path S with endpoints u and v .

$\sigma_{L,H} \cap K_{R,H} = \sigma_{L,H} \setminus K_{R,V} = \sigma_{L,H} - v_4$. Clearly, the first vertex of σ_s is v_1 . If the second vertex w of $\sigma_{L,H}$ is not v_4 , then $\sigma_s = v_1w$ is an interval of $\sigma_{X,H}$ implying that $s \in S_{L,H}$, a contradiction. Therefore, $w = v_4$, and $\sigma_s = v_1x$ where x is the third vertex of $\sigma_{L,H}$. We conclude that $S_{L,HH,2} = \{v_1x\}$. Recall that v_4 has three incident edges in G' . Since v_4 is the leftmost vertex of σ_R , none of these edges is in $S_{R,HH} \cup S_{L,VH}$. Moreover, at most one of them is in $S_{R,H} \cup S_{R,HV}$. Therefore, at least two of them are in $S_{L,H}$. Then these edges are necessarily v_4v_1 and v_4x . We conclude that $\{v_1, v_4, x\}$ induces a triangle in G' . In other words $\{v_1, v_4, x, v\}$ induces a diamond on G , a contradiction. Therefore

$$S_{X,HH,2} = \emptyset. \quad (7)$$

Finally, if $K_{X,V} = \emptyset$ we have $S_{X,HV} = \emptyset$ and $|S_{X,H,2}| \leq n - 1$. Otherwise, $|K_{X,V}| = 1, |K_{X,H}| = n - 1$ and we have $|S_{X,HV,2}| \leq 1$ and $|S_{X,H,2}| \leq n - 2$. In both cases we have

$$|S_{X,H,2} \cup S_{X,HV,2}| \leq n - 1. \quad (8)$$

Combining (6), (7) and (8) we obtain $|S_2| \leq 2(n - 1)$. Since $|S_2| = |E(G')| = |E(G)| - 4 = 2(n - 1)$, all the inequalities must hold with equality, in particular $|S_{X,H,2} \cup S_{X,HV,2}| = n - 1$. Therefore, every two consecutive vertices in σ_X are adjacent in G' . In other words, the permutation σ_X corresponds to a path Q_X of G' . The endpoints of Q_L (resp. Q_R) are v_1 and v_2 (resp. v_3 and v_4). Adding two edges incident to v to each Q_X , we get two edge disjoint Hamiltonian cycles of G . \square

4 Co-bipartite Graphs

In Section 4.1 we characterize B_1 -ENPG co-bipartite graphs. We show that there are two types of representations for B_1 -ENPG co-bipartite graphs. For each type of representation, we characterize their corresponding graphs. These characterizations imply a polynomial-time recognition algorithm. In Section 4.2 we present an efficient (linear-time) implementation of the algorithm.

4.1 Characterization of B_1 -ENPG Co-bipartite Graphs

We proceed with definitions and two related lemmas (Lemma 4.1, Lemma 4.2) that will be used in each of the above mentioned characterizations.

Let S be a path of a graph H with endpoints u, v . Two sets $\mathcal{P}_u, \mathcal{P}_v$ of paths *meet at* S if for $x \in \{u, v\}$ (a) every path of \mathcal{P}_x contains x (b) every path of \mathcal{P}_x has an endpoint that is a vertex of S different than x , and (c) a pair of paths $P_u \in \mathcal{P}_u, P_v \in \mathcal{P}_v$ may intersect only in S (see Figure 6).

A graph $G = (V, E)$ is a *difference graph* (equivalently *bipartite chain graph*) if every $v_i \in V$ can be assigned a real number a_i and there exists a positive real number T such that (a) $|a_i| < T$ for all i and

(b) $\{v_i, v_j\} \in E$ if and only if $|a_i - a_j| \geq T$. Every difference graph is bipartite where the bipartition is according to the sign of a_i .

Theorem 4.1 *Hammer et al. (1990)* If $G = (V, E)$ is a bipartite graph with bipartition $V = X \cup Y$ then the following statements are equivalent:

i) G is a difference graph.

ii) Let $\delta_1 < \delta_2 < \dots < \delta_s$ be distinct nonzero degrees in X , and $\delta_0 = 0$. Let $\sigma_1 < \sigma_2 < \dots < \sigma_t$ be distinct nonzero degrees in Y , and $\sigma_0 = 0$. Let $X = X_0 \cup X_1 \cup \dots \cup X_s$, $Y = Y_0 \cup Y_1 \cup \dots \cup Y_t$, where $X_i = \{x \in X \mid d(x) = \delta_i\}$, $Y_j = \{y \in Y \mid d(y) = \delta_j\}$. Then $s = t$ and for $x \in X_i$, $y \in Y_j$, $\{x, y\} \in E$ if and only if $i + j > t$.

Theorem 4.2 *Hammer et al. (1990)* A graph is a difference graph if and only if it is bipartite and $2K_2$ -free.

Lemma 4.1 Let $G_B = B(K, K', E)$ a difference graph, and t be the number of distinct nonzero degrees of vertices of K in G_B . Let H be a grid and S be a path of H with length at least $t + 2$ and no bends. Then there is a B_1 -ENPG representation $\langle H, \mathcal{P} \rangle$ of $G_C = C(K, K', E)$ such that \mathcal{P}_K and $\mathcal{P}_{K'}$ meet at S .

Proof: Let $\delta_1 < \delta_2 < \dots < \delta_s$ (resp. $\sigma_1 < \sigma_2 < \dots < \sigma_t$) be the distinct nonzero degrees in K (resp in K') in G_B . By Theorem 4.1 we have $s = t$. Let $-1, 0, 1, \dots, t + 1$ be $t + 3$ vertices of S such that 0 and $t + 2$ are the endpoints of S and they appear in this order on S . Let x (resp. x') be a vertex of K (resp. K'), and let i be such that $d_{G_B}(x) = \delta_i$ (resp. $d_{G_B}(x') = \sigma_{i'}$). The path P_x (resp. $P_{x'}$) is constructed between vertices -1 and i (resp. $t - j$ and $t + 1$).

With this construction $\mathcal{P}_K, \mathcal{P}_{K'}$ represent the cliques K and K' , moreover they meet at S . By the construction two paths $P_x \in \mathcal{P}_K, P_{x'} \in \mathcal{P}_{K'}$ intersect if and only if $i + j > t$. By Theorem 4.1 x and x' are adjacent if and only if $i + j > t$. Therefore, \mathcal{P} is a representation of $G = C(K, K', E)$. \square

Lemma 4.2

i) If two sets $\mathcal{P}_K, \mathcal{P}_{K'}$ of one-bend paths meet at a path S then $G_B = B(K, K', E)$ is a difference graph.
 ii) If a cobipartite graph $G = C(K, K', E)$ is an interval graph, then $G_B = B(K, K', E)$ is a difference graph.

Proof:

i) Let u, v be the endpoints of S . Let $T = |E(S)| + 1$ and r_i (resp. l_j) be the endpoint of the path $P_i \in \mathcal{P}_K$ (resp. $P_j \in \mathcal{P}_{K'}$) among the internal vertices of S . Let $a_i = |E(p_S(u, r_i))|$ (resp. $a_j = -|E(p_S(l_j, v))|$) where $p_T(x, y)$ is the unique path between vertices x and y of a tree T . By definition, $|a_i| \leq |E(S)| < T$ for every $i \in K \cup K'$. Two paths $P_i \in \mathcal{P}_K, P_j \in \mathcal{P}_{K'}$ have an edge in common if and only if $|a_i - a_j| \geq |E(S)| + 1 = T$. Therefore, G_B is a difference graph.

ii) Fix an interval representation of G . For $X \in \{K, K'\}$ let e_X be the edge of the representation that is common to all the paths \mathcal{P}_X representing the clique X . We can assume without loss of generality that e_K and $e_{K'}$ are the leftmost and rightmost edges of the representation. We now subdivide e_K and $e_{K'}$ by adding new vertices v_K and $v_{K'}$ respectively. Finally, if a path contains both e_K and $e_{K'}$

we truncate one edge from its end so that it contains v_K but not $v_{K'}$. In the new representation, \mathcal{P}_K and $\mathcal{P}_{K'}$ meet at the segment between v_K and $v_{K'}$.

□

Two representations $\langle H, \mathcal{P} \rangle$ and $\langle H', \mathcal{P}' \rangle$ are *bend-equivalent* if they are representations of the same graph G and the paths $P_v \in \mathcal{P}$ and $P'_v \in \mathcal{P}'$ representing the same vertex v of G have the same number of bends. We proceed with the following lemma that classifies all the B_1 -ENPG representations of a co-bipartite graph into two types.

Lemma 4.3 *Let $G = C(K, K', E)$ be a connected B_1 -ENPG co-bipartite graph with a representation $\langle H, \mathcal{P} \rangle$. Then G has a bend-equivalent representation $\langle H, \mathcal{P}' \rangle$ that satisfies exactly one of the following*

- i) $|\mathcal{S}(\cup\mathcal{P}'_K, \cup\mathcal{P}'_{K'})| = 1$ and $\cup\mathcal{P}'$ is a tree with maximum degree at most 3 with at most two vertices of degree 3 as depicted in Figure 7 (a).
- ii) $|\mathcal{S}(\cup\mathcal{P}'_K, \cup\mathcal{P}'_{K'})| = 2$ and the paths $\cup\mathcal{P}'_K$ and $\cup\mathcal{P}'_{K'}$ intersect as depicted in Figure 7 (b).

Proof: By Proposition 2.1, $\cup\mathcal{P}_K$ and $\cup\mathcal{P}_{K'}$ are two paths with at most 2 bends each. Let e_K (resp. $e_{K'}$) be an arbitrary edge of $\cap\mathcal{P}_K$ (resp. $\cap\mathcal{P}_{K'}$). The paths $\cup\mathcal{P}_K$ and $\cup\mathcal{P}_{K'}$ intersect in at least one edge, because otherwise G is not connected. Therefore, $|\mathcal{S}(\cup\mathcal{P}_K, \cup\mathcal{P}_{K'})| \geq 1$. We consider two disjoint cases:

- $|\mathcal{S}(\cup\mathcal{P}_K, \cup\mathcal{P}_{K'})| = 1$. Let $T = \cup\mathcal{P}$ and S be the unique segment of $\mathcal{S}(\cup\mathcal{P}_K, \cup\mathcal{P}_{K'})$. The collection T is clearly a tree, since any cycle in T is a cycle in one of $\cup\mathcal{P}_K, \cup\mathcal{P}_{K'}$. Any vertex of degree at least 3 in T is an endpoint of S , therefore there are at most 2 such vertices. On the other hand an endpoint of S has degree at most 3. Therefore $\Delta(T) \leq 3$ and there are at most 2 vertices of degree 3 in T .

Let u and v be the two endpoints of S . Let also e_u, e_v (respectively e'_u, e'_v) be the (at most four) edges not in S but belonging to $\cup\mathcal{P}_K$ (respectively $\cup\mathcal{P}'_{K'}$) and incident respectively to u and v . Now, shrink all the paths on the branches of T starting with e_u, e_v, e'_u, e'_v and not containing S to respectively the edges e_u, e_v, e'_u, e'_v . Clearly, this transformation does not create or remove any split between two paths and does not remove any intersections since all paths intersecting on one of these branches now intersect on the shrunken edge. To maintain bend-equivalence, we add one more edge to every path that loses a bend during the shrinkage (the same edge for all paths of the same branch). This new representation $\langle H, \mathcal{P}' \rangle$ is bend-equivalent to $\langle H, \mathcal{P} \rangle$ and $T' = \cup\mathcal{P}'$ is a tree with the claimed properties.

- $|\mathcal{S}(\cup\mathcal{P}_K, \cup\mathcal{P}_{K'})| \geq 2$. We claim that $\cup\mathcal{P}_K \cap \cup\mathcal{P}_{K'}$ contains only horizontal edges, or only vertical edges. Indeed, assume that there is a vertical edge e_V and a horizontal edge e_H in $\cup\mathcal{P}_K \cap \cup\mathcal{P}_{K'}$. We observe that there is a unique one bend path with e_V and e_H its end edges, and that any other connecting these edges contains at least three bends. Therefore, both $\cup\mathcal{P}_K$ and $\cup\mathcal{P}_{K'}$ contain this path. We conclude that e_V and e_H are in the same segment. As any other edge is either horizontal or vertical, we can proceed similarly for all the edges of $\cup\mathcal{P}_K \cap \cup\mathcal{P}_{K'}$ and prove that they all belong to the same segment, contradicting the fact that we have at least 2 segments. Assume without loss of generality that all the edges of $\cup\mathcal{P}_K \cap \cup\mathcal{P}_{K'}$ are vertical. Then every segment is a vertical path. No two segments can be on the same vertical line, because this will require at least one of $\cup\mathcal{P}_K,$




Fig. 7: Two types of B_1 -ENPG representation of connected co-bipartite graphs: (a) Type I: $|\mathcal{S}(K, K')| = 1$, $\cup\mathcal{P}$ is isomorphic to a tree T with $\Delta(T) \leq 3$ and at most two vertices u, v having degree 3, (b) Type II: $|\mathcal{S}(K, K')| = 2$, \mathcal{P}_K (resp. $\mathcal{P}_{K'}$) has exactly two bend points u, v (resp. u', v')

$\cup\mathcal{P}_{K'}$ to contain four bends. Moreover, three vertical segments in distinct vertical lines imply that \mathcal{P}_K and $\mathcal{P}_{K'}$ contain at least four bends each. Therefore, there are exactly 2 vertical segments and \mathcal{P}_K (also $\mathcal{P}_{K'}$) has exactly two bends.

Let u, v (resp. u', v') be the bends of $\cup\mathcal{P}_K$ (resp. $\cup\mathcal{P}_{K'}$). Then $\mathcal{S}(\cup\mathcal{P}_K, \cup\mathcal{P}_{K'})$ where S_u (resp. S_v) is on the same vertical line as u and u' (resp. v and v'). Moreover e_K (resp. $e_{K'}$) is between u and v (resp. u' and v') since otherwise we would have paths crossing both u and v (resp. u' and v') and thus 2 bends. If both the pairs u, u' and v, v' are on different sides of respectively S_u and S_v (as in Figure 7 (b)) then let $H' = H$ and $\mathcal{P}' = \mathcal{P}$ be the desired representation. Now consider the situation where u and u' are on the same side of S_u . Every path intersecting S_u crosses the same endpoint of S_u say without loss of generality u , implying that if a pair of paths from distinct cliques intersect at S_u , they split at this endpoint. Then remove S_u from every path of $\mathcal{P}_{K'}$ to obtain a bend-equivalent representation that contains one segment. The resulting representation can be transformed into a bend-equivalent representation $\langle H, \mathcal{P}' \rangle$ as described the previous bullet.

□

Based on Lemma 4.3, a B_1 -ENPG representation of a connected co-bipartite graph $G = C(K, K', E)$ is *Type I* (resp. *Type II*) if $|\mathcal{S}(K, K')| = 1$ (resp. $|\mathcal{S}(K, K')| = 2$).

We proceed with the characterization of B_1 -ENPG graphs having a Type II representation that turns out to be simpler than the characterization of the others. In the following lemma, a *trivial* connected component is an isolated vertex.

Lemma 4.4 *A connected twin-free co-bipartite graph $G = C(K, K', E)$ has a Type II B_1 -ENPG representation if and only if the bipartite graph $G_B = B(K, K', E)$ contains at most two non-trivial connected components each of which is a difference graph.*

Proof: (\Rightarrow) Let $\langle H, \mathcal{P} \rangle$ be a Type II B_1 -ENPG representation of G and u, v (resp. u', v') be the bends of $\cup\mathcal{P}$ (resp. $\cup\mathcal{P}'$) as depicted in Figure 7 b). For $x \in \{u, v\}$, let S_x be the segment contained in the path between x and x' . The paths of \mathcal{P} not intersecting with any of S_u, S_v correspond to isolated vertices of G_B . Since G is twin-free, there is at most one such path in \mathcal{P}_K (resp. $\mathcal{P}_{K'}$).

Each one of the remaining paths intersects exactly one of S_u, S_v , as otherwise such a path would contain two bends. For $X \in \{K, K'\}$ and $x \in \{u, v\}$ let \mathcal{P}_{X_x} be the paths of \mathcal{P}_X intersecting S_x . Then \mathcal{P}_{K_x} and $\mathcal{P}_{K'_x}$ meet at S_x . By Lemma 4.2, $G_B[K_x \cup K'_x]$ is a difference graph.

(\Leftarrow) It is sufficient to construct a Type II representation for the maximal case, i.e. G_B contains exactly two trivial connected components and two non-trivial connected components. Let $w \in K$ and $w' \in K'$ be the trivial connected components and $B(K_u, K'_u, E_u), B(K_v, K'_v, E_v)$ be the non-trivial connected components of G_B . We construct a rectangle as depicted in Figure 7 b) having vertical lines with $\max(\min(|K_u|, |K'_u|), \min(|K_v|, |K'_v|)) + 2$ edges, and horizontal lines with one edge $e_K = \{u, v\}$ and $e_{K'} = \{u', v'\}$. For $X \in \{K, K'\}$, and $x \in \{u, v\}$ the paths \mathcal{P}_{X_x} start with e_X and enter segment S_x . The other endpoints of the paths will be in the segment S_x . Then, for $x \in \{u, v\}$, \mathcal{P}_{K_x} and $\mathcal{P}_{K'_x}$ meet at S_x . Since $B(K_x, K'_x, E_x)$ is a difference graph, by Lemma 4.1, the endpoints can be determined such that $\mathcal{P}_{K_x} \cup \mathcal{P}_{K'_x}$ is a representation of $B(K_x, K'_x, E_x)$. The path P_w (resp. $P_{w'}$) consists of the edge e_K (resp. $e_{K'}$). It is easy to verify that this is a representation of G . \square

We proceed with the characterization of the B_1 -ENPG graphs with a Type I representation. For this purpose we resort to the following definitions from Fouquet et al. (2004).

Let $G = B(V, V', E)$ be a bipartite graph and $M \subseteq V \cup V'$. A vertex $v \in V \setminus M$ (resp. $v \in V' \setminus M$) *distinguishes* M if it has a neighbour in $M \cap V'$ (resp. $M \cap V$) and a non-neighbour in $M \cap V'$ (resp. $M \cap V$). A non-empty subset M of $V \cup V'$ is a *bimodule* of G if no vertex distinguishes M . It follows from the definition that $V \cup V'$ is a bimodule of G , and so are all the singletons and all the pairs of vertices with exactly one from V . These bimodules are the *trivial* bimodules of G .

A *zed* is a graph isomorphic to a P_4 or any induced subgraph of it. We note that a trivial bimodule different from $V \cup V'$ is a zed.

Lemma 4.5 *A connected twin-free co-bipartite graph $G = C(K, K', E)$ has a Type I B_1 -ENPG representation if and only if there is a set of vertices Z of G such that*

- i) Z is a zed of G ,
- ii) Z is a bimodule of $G_B = B(K, K', E)$, and
- iii) $G_B \setminus Z$ is a difference graph.

Moreover, if Z is a set of vertices of minimum size that satisfies i)-iii) and Z is a set of two non-adjacent vertices of G , then for the unique segment S of $\mathcal{S}(\cup K, \cup K')$ the following hold in every representation $\langle H, \mathcal{P} \rangle$:

- a) S is contained in at least one of the paths of \mathcal{P}_Z ,
- b) the endpoints of S have degree 3 in $\cup \mathcal{P}$ and these endpoints constitute $\text{split}(\cup \mathcal{P}_K, \cup \mathcal{P}_{K'})$.

Proof: (\Rightarrow) Let $\langle H, \mathcal{P} \rangle$ be a Type I B_1 -ENPG representation of G . By Lemma 4.3, $|\mathcal{S}(K, K')| = 1$ and $\cup \mathcal{P}$ is a tree. Let u, v be the endpoints of the unique segment S of $\mathcal{S}(K, K')$. We consider the following disjoint cases:

- $\{e_K, e_{K'}\} \not\subseteq E(S)$: Without loss of generality, suppose that $e_K \notin E(S)$ and u is closer to e_K than v . Consider two paths $P_{x'}, P_{y'} \in \mathcal{P}_{K'}$ that cross u . We observe that these paths are indistinguishable by the paths of \mathcal{P}_K . Namely, every path of \mathcal{P}_K either does not intersect any one of




Fig. 8: (a) Four special paths corresponding to a zed (b) The type of vertices and edge relations of a B_1 -ENPG co-bipartite graph having a Type I representation. K^{\emptyset} (resp. K'^{\emptyset}) is the set of vertices corresponding to the paths of \mathcal{P}_K (resp. $\mathcal{P}_{K'}$) crossing neither u nor v .

$P_{x'}, P_{y'}$, or intersects both and splits from both at u . Therefore the corresponding vertices x', y' are twins. As G is twin-free we conclude that there is at most one path of $\mathcal{P}_{K'}$ that crosses u . Similarly, consider two paths $P_x, P_y \in \mathcal{P}_K$ that cross v . These paths cross also u since e_K is an edge of both paths. Therefore, every path of $\mathcal{P}_{K'}$ either does not intersect any one of P_x, P_y , or intersects both and splits from both at either u or v , or intersects both and does not split from any of them. We conclude that there is at most one path of \mathcal{P}_K that crosses v . Let $\mathcal{P}_{Z'}$ be a set of these at most two paths. Namely, $\mathcal{P}_{Z'}$ consists of all the paths of $\mathcal{P}_{K'}$ crossing u and all the paths of \mathcal{P}_K that cross v . We now observe that $\cup(\mathcal{P} \setminus \mathcal{P}_{Z'})$ is either a path or the union of two edge-disjoint paths. In both cases no two paths split from each other, and their adjacency is determined only by the intersections. Therefore, the resulting graph $G \setminus Z'$ is an interval graph implying that $G_B \setminus Z'$ is a difference graph. We note that the path $P_{x'} \in \mathcal{P}_{K'}$ that crosses u is an isolated vertex of G_B , therefore for $Z = Z' \setminus \{x'\}$ we have that $G_B \setminus Z$ is a difference graph too, i.e. Z satisfies iii). Since $|Z| \leq 1$, Z satisfies i) and ii) trivially. This completes the proof of the first part of the claim. As for the second part, since $|Z| \leq 1$, any set of minimum size satisfying the conditions has at most one vertex. Therefore, the second part of the claim holds vacuously.

- $\{e_K, e_{K'}\} \subseteq E(S)$: We first note that we can assume $e_K \neq e_{K'}$ since otherwise we can subdivide this edge into two and rename the new edges as e_K and $e_{K'}$. Assume without loss of generality that e_K is closer to u than $e_{K'}$, (see Figure 8). Consider two paths $P_{x'}, P_{y'} \in \mathcal{P}_{K'}$ that cross u but not v . We observe that these paths are indistinguishable by the paths of \mathcal{P}_K . Therefore, the corresponding vertices are twins. As G is twin-free we conclude that there is at most one path $P_{K'}^u$ of $\mathcal{P}_{K'}$ that

crosses u and does not cross v . Similarly there is at most one path P_K^v of \mathcal{P}_K that crosses v but does not cross u , at most one path $P_{K'}^{u,v}$ of $\mathcal{P}_{K'}$ that crosses both u and v , and at most one path $P_K^{u,v}$ of \mathcal{P}_K that crosses both u and v . Let \mathcal{P}_Z be the set of these at most four paths. As in the previous case, $\cup(\mathcal{P} \setminus \mathcal{P}_Z)$ is a path, thus $G_B \setminus Z$ is a difference graph, i.e. Z satisfies iii). Assuming that all the four paths exist, it is easy to verify that their corresponding vertices $K^v, K'^u, K^{u,v}, K'^{u,v}$ constitute a P_4 with endpoints $K^{u,v}, K'^{u,v}$. Therefore, Z is a zed, i.e. Z satisfies i). Finally, we observe that P_K^v and $P_K^{u,v}$ are distinguishable only by $P_{K'}^u \in \mathcal{P}_Z$. In other words, they are indistinguishable by paths from $\mathcal{P}_{K'} \setminus \mathcal{P}_Z$. By symmetry, we conclude that Z is a bimodule of G_B , i.e. it satisfies ii). This concludes the proof of the first part of the claim. To prove the second part, assume by contradiction that there is a minimal set Z satisfying i)- iii) consisting of two vertices and none of the corresponding paths contains the segment S . Then these paths are $P_{K'}^u$ and P_K^v . We now observe that $P_{K'}^u \sim P_K^v$, i.e. K^v and K'^u are adjacent in G , contradicting the assumption that the vertices of Z are non-adjacent in G . This concludes the proof of a). If both paths contain S , then these paths are P_K^{uv} and $P_{K'}^{uv}$ and we have $split(\cup \mathcal{P}_K, \cup \mathcal{P}_{K'}) \supseteq split(P_K^{uv}, P_{K'}^{uv}) = \{u, v\}$, proving b) for this case. Otherwise, one of the paths does not contain S . Let, without loss of generality this path be $P_{K'}^u$. Then no path of $\mathcal{P}_{K'}$ crosses v . We conclude that $\cup(\mathcal{P} \setminus \{P_{K'}^u\})$ is a path, implying that the corresponding vertices induce a difference graph on G_B , contradicting the assumption that Z is a minimal set satisfying i)-iii).

(\Leftarrow) Given a zed Z of G satisfying the conditions of the lemma, we construct a Type I representation $\langle H, \mathcal{P} \rangle$ as follows. Without loss of generality we assume that Z is a P_4 with endpoints $y \in K, y' \in K'$ and internal vertices $x \in K, x' \in K'$. Let $\ell = \min(|K|, |K'|) + 2$. The graph H is a 3 by $\ell + 3$ vertices grid where each vertex is represented by an ordered pair from $[-1, \ell + 1] \times [-1, 1]$. The path P_x (resp. P_y) is between $(0, 0)$ (resp. $(-1, 0)$) and $(\ell, 1)$ with a bend at $(\ell, 0)$. The path $P_{x'}$ (resp. $P_{y'}$) is between $(\ell, 0)$ (resp. $(\ell + 1, 0)$) and $(0, -1)$ with a bend at $(0, 0)$. It is easy to verify that this correctly represents Z . The representation of the difference graph $G_B \setminus Z$ is two sets of paths that meet at the line segment between $(0, 0)$ and $(\ell, 0)$. By Lemma 4.1, the endpoints of the paths within this segment can be determined in accordance with the difference graph $G_B \setminus Z$. The other endpoints of these paths are determined so as to satisfy the adjacencies of vertices of Z with other vertices, as follows: The other endpoint of every path of $\mathcal{P}_{K' \cap N_G(y)}$ (resp. $\mathcal{P}_{K' \setminus N_G(y)}$) is $(\ell, 0)$ (resp. $(\ell + 1, 0)$). The other endpoint of every path of $\mathcal{P}_{K \cap N_G(y')}$ (resp. $\mathcal{P}_{K \setminus N_G(y')}$) is $(0, 0)$ (resp. $(-1, 0)$). \square

By Lemmata 4.4 and 4.5 we have the following Theorem.

Theorem 4.3 *Let $G = C(K, K', E)$ be a connected, twin-free co-bipartite graph, and $G_B = B(K, K', E)$. Then, $G \in \mathbf{B}_1$ -ENPG if and only if at least one of the following holds:*

- i) G_B contains at most two non-trivial connected components each of which is a difference graph.
- ii) G contains a zed Z that is a bimodule of G_B such that $G_B \setminus Z$ is a difference graph.

Since all the properties mentioned in Theorem 4.3 can be tested in polynomial time we have the following corollary.

Corollary 4.1 \mathbf{B}_1 -ENPG co-bipartite graphs can be recognized in polynomial time.

4.2 Efficient Recognition Algorithm

In this section we describe an efficient algorithm, namely Algorithm 1, to recognize whether a co-bipartite graph is B_1 -ENPG using the characterization of Theorem 4.3. In Algorithm 1, `ISYPEI` is a function taking as input a connected twin-free cobipartite graph and a subset Z of vertices to decide if there is $Z' \supseteq Z$ for the graph being B_1 -ENPG of Type I. Similarly, `ISYPEII` takes a connected twin-free cobipartite graph G and returns "YES" if G is B_1 -ENPG of Type II, and "NO" otherwise. As for function `FINDBIMODULEZED`, it takes a twin-free cobipartite graph G and a Z of G to return the minimum superset of Z that is a zed of G and a bimodule of G_B , if any. Lastly, the function `ISDIFFERENCE` in Algorithm 1 takes a bipartite graph G and either indicates that G is a difference graph or provides a $2K_2$ certifying that G is not a difference graph.

Theorem 4.4 *Given a co-bipartite graph $G = C(K, K', E)$, Algorithm 1 decides in time $O(|K| + |K'| + |E|)$ whether G is B_1 -ENPG.*

Proof: Let $n = |K| + |K'|$, $m = |E|$. Let $T_{diff}(n, m)$ be the running time of `ISDIFFERENCE` on a graph with n vertices and m edges, and let $T_{bm}(n, m)$ be the running time of `FINDBIMODULEZED` that finds a minimum zed of G that is a bimodule of G_B and contains a given zed Z . Finally let $\alpha(n, m) \stackrel{def}{=} T_{diff}(n, m) + T_{bm}(n, m)$.

The correctness of the algorithm follows from Observations 2.1, 2.2, Lemma 4.3 and from the correctness of the functions `ISYPEI` and `ISYPEII` that we prove in the sequel.

The correctness of `ISYPEI` is based on Lemma 4.5. A subset Z of vertices of G satisfying i)-iii) of Lemma 4.5 is termed as a *certificate* through this proof. We now show that given a twin-free cobipartite graph G and $Z \subseteq V(G)$, `ISYPEI` returns "YES" if and only if there exists a certificate $Z' \supseteq Z$. Moreover, we show that its running time is at most $5^{5-|Z|}\alpha(n, m)$ when $|Z| \leq 4$ and constant otherwise.

We first observe that if Z is not a zed, then no superset of Z is a zed, and the algorithm returns correctly "NO" in constant time at line 8. Therefore, our claim is correct whenever Z is not a zed. We proceed by induction on $5 - |Z|$. If $5 - |Z| = 0$, then Z is not a zed and the algorithm returns "NO" in constant time. In the sequel we assume that Z is a zed. In this case, Z is verified to be a zed by `ISYPEI` in constant time and `ISYPEI` proceeds to line 9 to find (in time $T_{bm}(n, m)$) the minimal bimodule Z' of G_B that contains Z and is a zed of G . We consider three cases according to the branching of `ISYPEI`.

- $Z' = Z$ (i.e. Z is a bimodule of G_B), **and $G_B \setminus Z$ is a difference graph:** $G_B \setminus Z$ is verified to be a difference graph by `ISYPEI` at line 11. It returns "YES" which is correct by Lemma 4.5 since Z is a certificate. The running time is $\alpha(n, m)$, and the result follows since $1 \leq 5^{5-|Z|}$.
- $Z' = Z$ (i.e. Z is a bimodule of G_B), **but $G_B \setminus Z$ is not a difference graph:** As $G_B \setminus Z$ is not a difference graph, there is a set $U \subseteq K \cup K' \setminus Z$ such that $G_B[U]$ is a $2K_2$. Every certificate $Z' \supseteq Z$ must contain at least one vertex of U because otherwise $G_B \setminus Z'$ contains $G_B[U]$ which is a $2K_2$. Therefore, `ISYPEI` proceeds recursively calling `ISYPEI` on $(G, Z \cup \{u\})$ for each $u \in U$. The algorithm returns "YES" if and only if one of the guesses succeeds. Then, the total running time is at most $\alpha(n, m) + 4 \cdot 5^{5-(|Z|+1)}\alpha(n, m) < (1 + 4 \cdot 5^{4-|Z|})\alpha(n, m)$. Since $1 \leq 5^{4-|Z|}$ we conclude that the running time is at most $5^{5-|Z|}\alpha(n, m)$.
- $Z' \neq Z$ (i.e. Z is not a bimodule of G_B): If Z' exists, the definition of a bimodule implies that any certificate that contains Z has to contain Z' . Therefore, `ISYPEI`(G, Z') is invoked and its result is

returned. Otherwise, no certificate contains Z and "NO" is returned. The running time of ISTYPEI is $T_{bm}(n, m) + 5^{5-|Z'|}\alpha(n, m) < (1 + 5^{5-|Z'|})\alpha(n, m) \leq 5^{5-|Z|}\alpha(n, m)$.

Since ISTYPEI is invoked initially at line 3 with $Z = \emptyset$, together with Lemma 4.5 this implies that the algorithm recognizes correctly graphs having a Type I representation. Moreover, the running time of line 3 is $5^{5-|\emptyset|}\alpha(n, m) = O(\alpha(n, m))$.

The correctness of ISTYPEII follows directly from Lemma 4.4. The connected components of G_B can be calculated in $O(n + m)$ time using breadth first search. Therefore, the running time of ISTYPEII is $O(T_{diff}(n, m)) = O(\alpha(n, m))$.

We now calculate the running time of the algorithm. All the twins of a graph can be removed in time $O(n + m)$ using partition refinement, i.e. starting from the trivial partition consisting of one set, and iteratively refining this partition using the closed neighborhoods of the vertices (see Habib et al. (1999)). Each set of the resulting partition constitutes a set of twins. Summarizing, we get that the running time of Algorithm 1 is $O(\alpha(n, m)) = O(T_{diff}(n, m) + T_{bm}(n, m))$.

$T_{diff}(n, m)$ is $O(n + m)$ (see Heggernes and Kratsch (2006)). It remains to prove the correctness of `FINDBIMODULEZED` and calculate its running time $T_{bm}(n, m)$. We consider the case where Z contains at most one vertex from each one of K and K' and the complementing case where Z contains at least two vertices from K separately.

- $Z = \emptyset$ or Z is a singleton or Z is a pair of vertices of $K \times K'$. By definition, Z is both a zed of G and a bimodule of G_B . Therefore, Z is the minimal bimodule of G_B that is a zed of G , and contains Z . In this case `FINDBIMODULEZED` return Z in constant time.
- Without loss of generality $Z \cap K$ contains at least two vertices u_1, u_2 . We note that $Z \cap K = \{u_1, u_2\}$, because otherwise Z contains a K_3 contradicting the fact that it is a zed. Let Z' be the superset of Z obtained by adding to it all the vertices that distinguish u_1 and u_2 . Formally, $Z' \stackrel{def}{=} (N_{G_B}(u_1) \Delta N_{G_B}(u_2)) \cup Z$. If Z' is not a zed we can return that no superset of Z is both a zed of G and a bimodule of G_B . Now, let Z' be a zed and let $U' = Z' \cap K'$. If $|U'| \leq 1$ then Z' is the minimal subset that contains Z and is both a zed of G and a bimodule of G_B . If $|U'| > 2$ then Z' is not a zed. Assume $|U'| = 2$ and let $U' = \{u'_1, u'_2\}$. We now add to Z' , the set of vertices of K that distinguish U' to get Z'' . If $Z'' = Z'$ then Z' is the minimal superset of Z that is both a zed of G and a bimodule of G_B . Otherwise every bimodule that contains Z' has to contain also Z'' . However $|Z'' \cap K| > |Z \cap K| = 2$, implying that Z'' contains a K_3 , and is thus not a zed. In this case, we conclude that there is no superset of Z as required.

As for the running time, we observe that all the operations can be performed in constant time except lines 30 and 35 that take time $O(|K'|)$ and $O(|K|)$, respectively. Therefore, the running time $T_{bm}(n, m)$ of `FINDBIMODULEZED` is at most $O(|K| + |K'|) = O(n)$. We conclude that the running time of Algorithm 1 is $O(T_{diff}(n, m) + T_{bm}(n, m)) = O(n + m)$. \square

We conclude with an interesting remark, pointing to a fundamental difference between EPG and ENPG graphs. A graph is B_k -EPG if it has an EPG representation $\langle H, \mathcal{P} \rangle$ such that every path of \mathcal{P} has at most k bends. It is known that given a B_k -EPG representation it is always possible to modify the paths such that every path has exactly k bend; indeed, if there is a path with less than k bends, one can subdivide the edges of the host grid (and consequently all the paths containing the related edges) as needed to introduce

new bends until it has exactly k bends, without creating any new intersection or split Golumbic (2015). The following proposition states that this does not hold for B_k -ENPG graphs.

Proposition 4.1 *Every B_1 -ENPG representation of a graph $G = C(K, K', E)$ such that $G_B = B(K, K', E)$ is isomorphic to $3K_2$ contains at least one path with zero bend.*

Proof: Consider a set Z consisting of two non-adjacent vertices of G . Then Z is a trivial bimodule of G_B and a zed of G . Moreover, by Theorem 4.2 $G_B \setminus Z$ is a difference graph since it does not contain a $2K_2$. Therefore, Z satisfies conditions i)-iii) of Lemma 4.5. Then G is B_1 -ENPG.

Let $\langle H, \mathcal{P} \rangle$ be a B_1 -ENPG representation of G . Since G_B has three non-trivial connected components, by Lemma 4.4, $\langle H, \mathcal{P} \rangle$ is a Type I representation. For any single vertex v of G , the graph $G_B \setminus \{v\}$ contains a $2K_2$ therefore fails to satisfy condition iii). We conclude that Z is a set of minimum size satisfying the conditions i)-iii) of Lemma 4.5. Moreover, Z consists of two non-adjacent vertices of G . Therefore, the unique segment S of $\mathcal{S}(K, K')$ has the properties a) and b) mentioned in the same Lemma.

Let $Z = \{x, y'\}$ where $x \in K$ and $y' \in K'$, and let y and x' be the unique neighbors in G_B of x and y' respectively. Let also u, v be the endpoints of S . By property a, without loss of generality P_x contains S . Therefore, $P_{x'}$ is contained in S as otherwise it would split from P_x in at least one of u, v , contradicting the fact that x and x' are adjacent. By property b of the lemma, u and v are split points. To conclude the claim, we now show that $P_{x'}$ has no bends. Assume by contradiction that $P_{x'}$ has a bend w . Then w is a bend of S and also of P_x . Therefore, P_x does not bend neither at u nor in v as otherwise it would contain 2 bends. We conclude that both u and v are bends of $\cup \mathcal{P}_{K'}$. Clearly, w is also a bend of $\cup \mathcal{P}_{K'}$. Then $\cup \mathcal{P}_{K'}$ has 3 bends, contradicting Proposition 2.1. \square

5 Summary and Future Work

In Boyacı et al. (2015b) we showed that ENPG contains an infinite hierarchy of subclasses that are obtained by restricting the number of bends. In this work we showed that B_1 -ENPG graphs are properly included in B_2 -ENPG graphs. The question whether B_2 -ENPG \subsetneq B_3 -ENPG \subsetneq \dots remains open.

In this work, we studied the intersection of B_1 -ENPG with some special chordal graphs. We showed that the recognition problem of B_1 -ENPG graphs in NP-complete even for a very restricted sub family of split graphs. On the other hand we showed that this recognition problem is polynomial-time solvable within the family of co-bipartite graphs. A forbidden subgraph characterization of B_1 -ENPG co-bipartite graphs is also work in progress.

We also showed that unlike B_k -EPG graphs that always have a representation in which every path has exactly k bends, some B_1 -ENPG graphs can not be represented using only paths having (exactly) one bend. One can define and study the graphs of edge intersecting non-splitting paths with exactly k bends. Another possible direction is to follow the approach of Cameron et al. (2016) and consider B_1 -ENPG representations restricted to subsets of the four possible rectilinear paths with one bend.

We showed that trees and cycles are B_1 -ENPG. The characterization of their representations is work in progress. A natural extension of such a characterization is to investigate the relationship of B_1 -ENPG graphs and cactus graphs. Another possible extension is to use the characterization of the special case of C_4 to characterize induced sub-grids. A non-trivial characterization would imply that not every bipartite graph is B_1 -ENPG. Therefore, it would be natural to consider the recognition problem of B_1 -ENPG bipartite graphs. The following interpretation of our results suggests that the latter problem is NP-hard:

Algorithm 1 B_1 -ENPG \cap Co-bipartite Recognition**Require:** A co-bipartite graph $G = C(K, K', E)$

- 1: **if** G is not connected **then return** "YES" $\triangleright G$ has a trivial B_1 -ENPG representation.
- 2: Make G twin-free using modular decomposition.
- 3: **if** $\text{ISTYPEI}(G, \emptyset)$ **then return** "YES".
- 4: **if** $\text{ISTYPEII}(G)$ **then return** "YES".
- 5: **return** "NO".

6: **function** $\text{ISTYPEI}(G = C(K, K', E), Z)$ **Require:** G is connected, twin-free, $Z \subseteq V(G)$ **Ensure:** returns whether there is a certificate $Z' \supseteq Z$ for G being Type I

- 7: $G_B \leftarrow B(K, K', E)$.
- 8: **if** $G[Z]$ is not a zed **then return** "NO".
- 9: $Z' \leftarrow \text{FINDBIMODULEZED}(G, Z)$.
- 10: **if** $Z' = Z$ **then** $\triangleright Z$ is a zed of G and also a bimodule of G_B
- 11: **if** $\text{ISDIFFERENCE}(G_B \setminus Z)$ **then return** "YES".
- 12: Let $U \subseteq (K \cup K') \setminus Z$ such that $G_B[U]$ is a $2K_2$.
- 13: **for** $u \in U$ **do**
- 14: **if** $\text{ISTYPEI}(G, Z \cup \{u\})$ **then return** "YES".
- 15: **return** "NO".
- 16: **else**
- 17: **if** $Z' \neq \text{NULL}$ **then return** $\text{ISTYPEI}(G, Z')$.
- 18: **else return** "NO".

19: **function** $\text{ISTYPEII}(G = C(K, K', E))$ **Require:** G is connected, twin-free**Ensure:** returns whether G has a Type II representation

- 20: $G_B \leftarrow B(K, K', E)$.
- 21: Remove all isolated vertices from G_B . \triangleright There are at most two of them
- 22: Calculate the connected components G_1, \dots, G_k of G_B .
- 23: **if** $k > 2$ **then return** "NO".
- 24: **if** not $\text{ISDIFFERENCE}(G_1)$ **then return** "NO".
- 25: **if** not $\text{ISDIFFERENCE}(G_2)$ **then return** "NO".
- 26: **return** "YES".

27: **function** $\text{FINDBIMODULEZED}(G = C(K, K', E), Z)$ **Require:** G is twin-free, Z is a zed of G **Ensure:** Returns the minimum superset of Z that is a zed of G and a bimodule of G_B

- 28: **if** $|Z \cap K| \leq 1$ and $|Z \cap K'| \leq 1$ **then return** Z .
- 29: Let without loss of generality $Z \cap K = \{u_1, u_2\}$.
- 30: $Z' \leftarrow (N_{G_B}(u_1) \Delta N_{G_B}(u_2)) \cup Z$.
- 31: **if** Z' is not a zed **then return** NULL.
- 32: $U' \leftarrow Z' \cap K'$.
- 33: **if** $|U'| \leq 1$ **then return** Z' .
- 34: Let without loss of generality $U' = \{u'_1, u'_2\}$.
- 35: $Z'' \leftarrow (N_{G_B}(u'_1) \Delta N_{G_B}(u'_2)) \cup Z'$.
- 36: **if** $Z'' = Z'$ **then return** Z'
- 37: **else return** NULL.

38: **function** $\text{ISDIFFERENCE}(G)$ \triangleright Heggernes and Kratsch (2006)**Require:** G is bipartite**Ensure:** Returns "YES" if G is a difference graph and a $2K_2$ of G otherwise.

A clique provides substantial information on the representation, and when the graph is partitioned into two cliques we are able to recognize B_1 -ENPG graphs. However, the absence of one such clique (in case of split graphs) already makes the problem NP-hard. In case of bipartite graphs both of the cliques are absent.

References

- T. Akiyama, T. Nishizeki, and N. Saito. NP-completeness of the hamiltonian cycle problem for bipartite graphs. *Journal of Information Processing*, 3(2):73 – 76, 1980.
- A. Asinowski and B. Ries. Some properties of edge intersection graphs of single-bend paths on a grid. *Discrete Mathematics*, 312(2):427 – 440, 2012. ISSN 0012-365X. doi: <http://dx.doi.org/10.1016/j.disc.2011.10.005>.
- T. C. Biedl and M. Stern. On edge-intersection graphs of k-bend paths in grids. *Discrete Mathematics & Theoretical Computer Science*, 12(1):1–12, 2010.
- A. Boyacı, T. Ekim, M. Shalom, and S. Zaks. Graphs of edge-intersecting non-splitting paths in a tree: Towards hole representations-Part I. *Discrete Applied Mathematics*, 2015a. doi: [10.1016/j.dam.2015.07.024](https://doi.org/10.1016/j.dam.2015.07.024). available online.
- A. Boyacı, T. Ekim, M. Shalom, and S. Zaks. Graphs of edge-intersecting and non-splitting paths. *Theoretical Computer Science*, 629:40–50, 2015b. doi: [10.1016/j.tcs.2015.10.004](https://doi.org/10.1016/j.tcs.2015.10.004).
- K. Cameron, S. Chaplick, and C. T. Hoàng. Edge intersection graphs of L-shaped paths in grids. *Discrete Applied Mathematics*, 210:185 – 194, 2016. doi: <http://dx.doi.org/10.1016/j.dam.2015.01.039>.
- D. Epstein, M. Golumbic, and G. Morgenstern. Approximation algorithms for B_1 -EPG graphs. *Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics)*, 8037 LNCS:328–340, 2013.
- J.-L. Fouquet, M. Habib, F. de Montgolfier, and J.-M. Vanherpe. Bimodular decomposition of bipartite graphs. In *30th International Workshop on Graph Theoretic Concepts in Computer Science, WG*, pages 117–128, 2004. doi: [10.1007/978-3-540-30559-0_10](https://doi.org/10.1007/978-3-540-30559-0_10).
- M. Golumbic. Personal communication, 2015.
- M. C. Golumbic and R. E. Jamison. The edge intersection graphs of paths in a tree. *Journal of Combinatorial Theory, Series B*, 38(1):8 – 22, 1985a. ISSN 0095-8956. doi: [10.1016/0095-8956\(85\)90088-7](https://doi.org/10.1016/0095-8956(85)90088-7).
- M. C. Golumbic and R. E. Jamison. Edge and vertex intersection of paths in a tree. *Discrete Mathematics*, 55(2):151 – 159, 1985b. ISSN 0012-365X. doi: [10.1016/0012-365X\(85\)90043-3](https://doi.org/10.1016/0012-365X(85)90043-3).
- M. C. Golumbic, M. Lipshteyn, and M. Stern. Edge intersection graphs of single bend paths on a grid. *Networks*, 54(3):130–138, 2009.
- M. Habib, C. Paul, and L. Viennot. Partition refinement techniques: An interesting algorithmic tool kit. *Int. J. Found. Comput. Sci.*, 10(2):147–170, 1999. doi: [10.1142/S0129054199000125](https://doi.org/10.1142/S0129054199000125). URL <http://dx.doi.org/10.1142/S0129054199000125>.

- P. L. Hammer, U. N. Peled, and X. Sun. Difference graphs. *Discrete Appl. Math.*, 28(1):35–44, July 1990. ISSN 0166-218X. doi: 10.1016/0166-218X(90)90092-Q.
- P. Heggernes and D. Kratsch. Linear-time certifying algorithms for recognizing split graphs and related graph classes. *Reports in Informatics*, 328, 2006.
- D. Heldt, K. Knauer, and T. Ueckerdt. Edge-intersection graphs of grid paths: the bend-number. *Discrete Applied Mathematics*, 167:144–162, 2014. doi: 10.1016/j.dam.2013.10.035.
- A. Kotzig. Aus der Theorie der endlichen regulären Graphen dritten und vierten Grades. *Časopis Pěst. Mat.*, (82):76–92, 1957.
- R. E. Tarjan. Decomposition by clique separators. *Discrete Mathematics*, 55(2):221 – 232, 1985. ISSN 0012-365X. doi: 10.1016/0012-365X(85)90051-2.