

NONCOMMUTATIVE SYMMETRIC FUNCTIONS ASSOCIATED WITH A CODE, LAZARD ELIMINATION, AND WITT VECTORS

Jean-Gabriel Luque, Jean-Yves Thibon

► **To cite this version:**

Jean-Gabriel Luque, Jean-Yves Thibon. NONCOMMUTATIVE SYMMETRIC FUNCTIONS ASSOCIATED WITH A CODE, LAZARD ELIMINATION, AND WITT VECTORS. *Discrete Mathematics and Theoretical Computer Science, DMTCS*, 2007, 9 (2), pp.59-72. hal-00084906

HAL Id: hal-00084906

<https://hal.archives-ouvertes.fr/hal-00084906>

Submitted on 10 Jul 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

NONCOMMUTATIVE SYMMETRIC FUNCTIONS ASSOCIATED WITH A CODE, LAZARD ELIMINATION, AND WITT VECTORS

JEAN-GABRIEL LUQUE AND JEAN-YVES THIBON

IGM, Laboratoire d'informatique
77454 Marne-la-Vallée Cedex 2, France.
{luque,jyt}@univ-mlv.fr

July 11, 2006

Abstract

The construction of the universal ring of Witt vectors is related to Lazard's factorizations of free monoids by means of a noncommutative analogue. This is done by associating to a code a specialization of noncommutative symmetric functions.

Keywords: Witt vectors, codes, symmetric functions, factorizations of free monoids

1 Introduction

Among Ernst Witt's many contributions to mathematics, one finds two apparently unrelated ideas, both published in 1937 in two consecutive issues of Crelle's journal. The first one, the ring of Witt vectors [19] (a generalisation of p -adic numbers), solves a problem in commutative algebra, whilst the second one, the introduction of the free Lie algebra [20], definitely pertains to noncommutative mathematics.

The aim of this note is to point out a close connection between both constructions, through the notion of noncommutative symmetric functions. What comes out is that the natural noncommutative analogues of the symmetric functions classically associated to the construction of universal Witt vectors can be related to another classical topic in the combinatorics of free Lie algebras: Lazard's factorizations of free monoids. This relation manifests itself when the elementary noncommutative symmetric functions are specialized by means of a code.

Our notations will be those of [11] and [7]. If S is a set of words, we denote by $\underline{S} = \sum_{w \in S} w$ its characteristic series.

2 Noncommutative Witt symmetric functions

Let us first recall the construction of the universal ring $W(R)$ of Witt vectors associated to a commutative ring R . This is not Witt's original construction, but a simpler one that he found much later, and that he communicated to S. Lang, who published it as a series of exercises in his famous textbook of Algebra [9].

The ring $W(R)$ can be characterized by the following properties [4] (see also [8, 5]):

(W1) As a set, $W(R) = \{\mathbf{a} = (a_n)_{n \geq 1}, a_n \in R\}$, and for any ring homomorphism $f : R \rightarrow R'$, the map $W(f) : \mathbf{a} \mapsto (f(a_n))_{n \geq 1}$ is a ring homomorphism.

(W2) The maps $w_n : \mathbf{a} \mapsto \sum_{d|n} da_d^{n/d}$ are ring homomorphisms $W(R) \rightarrow R$.

Operations on Witt vectors are better understood in terms of symmetric functions. Let $X = \{x_n, n \geq 1\}$ be an infinite set of commuting indeterminates (called here an *alphabet*), and following [14], define symmetric functions $q_n(X)$ by

$$\prod_{n \geq 1} \frac{1}{1 - t^n q_n(X)} = \sigma_t(X) := \sum_{n \geq 1} t^n h_n(X) \quad (1)$$

where the complete homogeneous functions $h_n(X)$ are defined by $\sigma_t(X) = \prod_{n \geq 1} (1 - tx_n)^{-1}$ (*cf.* [11]). The q_n 's are connected to the power-sums $p_n(X) =$

$\sum_i x_i^n$ by

$$p_n = \sum_{d|n} dq_d^{n/d} \quad (2)$$

and condition (W2) can be regarded as expressing the familiar properties of power sums

$$\begin{aligned} p_n(X + Y) &= p_n(X) + p_n(Y) \\ p_n(XY) &= p_n(X)p_n(Y) \end{aligned}$$

where we use the λ -ring notation (an alphabet is identified with the formal sum of its elements). The transformation (1) is also used in [2, 12].

Reutenauer [14] studied the symmetric functions $q_n(X)$ and made the conjecture that for $n \geq 2$, $(-q_n)$ is Schur positive. This conjecture was proved in [17], in a generalised noncommutative form. Denoting as in [7] the noncommutative complete homogeneous symmetric functions of an alphabet A by $S_n(A)$, one introduces noncommutative Witt symmetric functions $Q_n(A)$ by the identity

$$\prod_{n \geq 1}^{\rightarrow} \frac{1}{1 - t^n Q_n(A)} = \sigma_t(A) := \sum_{n \geq 1} t^n S_n(A) \quad (3)$$

Then, it is proved in [17] that for $n \geq 2$, $(-Q_n)$ is a positive, multiplicity free, sum of noncommutative ribbon Schur functions R_I .

3 Noncommutative symmetric functions associated to a code

Let A be an alphabet and $C \subset A^+$ a *code*, *i.e.* a minimal generating set of a free submonoid C^* in A^* . Suppose we have a decomposition of C as a disjoint union

$$C = \coprod_{n \geq 1} C_n \quad (4)$$

of possibly empty subsets C_n . In what follows, we shall in general assume that $C_n = C \cap A^n$, but this restriction is not necessary. In any case, we will consider that the elements of C_n have degree n .

We will denote by $l(w)$ the length of a word w . Recall that, as a graded algebra, $\mathbf{Sym} = \mathbb{Q}\langle \Lambda_1, \Lambda_2, \dots \rangle$ where Λ_n has degree n . We can define a

specialization $\mathbf{Sym}[C]$ of the algebra of noncommutative symmetric functions by setting

$$\Lambda_n[C] = (-1)^{n-1} \underline{C}_n. \quad (5)$$

With this choice of signs, the complete symmetric functions are then given by

$$S_n[C] = \sum_{w \in (C^*)_n} w, \quad (6)$$

the sum of all elements of degree n in C^* . We call C -Witt symmetric function the value $Q_i[C]$ of Q_i under this specialization. Let us give the first $Q_i[C]$ for three examples

Example 1 The Fibonacci prefix code $C = \{b, ab\}$.

$$\begin{aligned} Q_1[C] &= b \\ Q_2[C] &= ab \\ Q_3[C] &= ab^2 \\ Q_4[C] &= ab^3 \\ Q_5[C] &= ab^2ab + ab^4 \\ Q_6[C] &= ab^3ab + ab^5 \\ Q_7[C] &= ab^6 + ab^2(ab)^2 + ab^4ab + ab^3ab^2 \end{aligned}$$

Example 2 The infinite prefix code $C = ba^*$.

$$\begin{aligned} Q_1[C] &= b \\ Q_2[C] &= ba \\ Q_3[C] &= ba^2 + bab \\ Q_4[C] &= bab^2 + ba^3 + ba^2b \\ Q_5[C] &= bab^2a + bab^3 + ba^2b^2 + ba^2b^2 + ba^4 + ba^2ba \\ Q_6[C] &= ba^4b + ba^3b^2 + ba^2b^2a + bab^3a + bab^4 + ba^3ba + ba^5 + ba^2b^3 \end{aligned}$$

Example 3 The Dyck code D (for the Dyck language $\underline{D}^* = 1 + a\underline{D}^*b\underline{D}^*$.)

$$\begin{aligned} Q_1[D] &= Q_3[D] = Q_5[D] = Q_7[D] = 0 \\ Q_2[D] &= ab \\ Q_4[D] &= aabb \\ Q_6[D] &= aaabbb + aababb + aabbab \\ Q_8[D] &= aaaabbbb + aaababbb + aabaabbb + aaabbbab + aaabbabb \\ &+ aabbabab + aabababb + aababbab \end{aligned}$$

On these examples, we remark that each Q_i is multiplicity free and is the characteristic series of a code. In the following section, we prove that it is always the case and give a characterization of these codes in terms of Lazard's factorizations.

4 Witt symmetric functions and factorization

4.1 Lazard elimination process

We recall that a factorization of a monoid \mathbb{M} is an ordered family of monoids $\mathbb{F} = (\mathbb{M}_i)_{i \in I}$ such that each element $m \in \mathbb{M}$ admits an unique decomposition

$$m = m_{i_1} \cdots m_{i_k} \quad (7)$$

where $i_1 > \cdots > i_k$ and $m_{i_1} \in \mathbb{M}_{i_1}, \dots, m_{i_k} \in \mathbb{M}_{i_k}$. In the case of free monoids $\mathbb{M} = A^*$, this property can be stated in terms of generating series

$$\sum_{w \in A^*} w = \prod_i^{\leftarrow} \sum_{w \in \mathbb{M}_i} w \quad (8)$$

A submonoid $\mathbb{M}' \subset \mathbb{M}$ can be characterized by its generating set $\mathbb{M}' \setminus \mathbb{M}'^2$. In the sequel, a factorization will be denoted by the sequence of the generating sets of its components, and for a factorization $\mathbb{F} = (C_i)_{i \in I}$ of A^* into free submonoids, (8) reads

$$\frac{1}{1 - \underline{A}} = \prod_i^{\leftarrow} \frac{1}{1 - \underline{C}_i}. \quad (9)$$

Factorisations of free monoids have been extensively studied and the reader can refer to [3, 18] for a survey. A simple but relevant example is a *Lazard bisection*: considering a subalphabet $B \subset A$, one has

$$A^* = B^*((A \setminus B)B^*)^*. \quad (10)$$

The pair $(B, (A \setminus B)B^*)$ is a factorization of A^* (*cf.* [18, 3], and [15] for applications to free Lie algebras). Now, from (10) we can obtain a trisection (*i.e.* a factorization in three submonoids) by iterating the process on the left or the right factor, and so on. Factorisations which can be generated applying only Lazard bisections on the right factors will be called *Lazard right compositions*. This is a simple particular case of a *locally finite right regular*

factorization [18]. Let A be an alphabet (possibly infinite) and $\omega : A \rightarrow \mathbb{N} \setminus 0$ a weight function. This function can be extended uniquely as a morphism $\omega : A^* \rightarrow (\mathbb{N}, +)$. We can associate a factorization $\mathbb{F}(A, \omega)$ to this weight as follows. Let $(Z_i)_{i \geq 1}$ and $(C_i)_{i \geq 1}$ be the sequences of codes defined by the recurrence relations

1. $Z_1 = A$
2. for each integer $i > 0$, $C_i = Z_i \cap \{w \in A^* \mid \omega(w) = i\}$.
3. for each integer $i > 0$, $Z_{i+1} = (Z_i \setminus C_i)C_i^*$

The sequence $\mathbb{F}(A, \omega) = (C_i)_{i \in I}$ obtained by omitting the empty elements in $(C_i)_{i \geq 1}$ is a Lazard right composition. Since each code $C \in A^*$ admits the length of words as weight function, we define the *right length factorization* $\mathbb{F}(C^*)$ of C^* by

$$\mathbb{F}(C^*) = \mathbb{F}(C, l). \quad (11)$$

We can remark that a code is homogeneous if and only if $\mathbb{F}(C^*) = (C)$.

4.2 Computation of the C -Witt symmetric functions

The equality between formal series (3) can be rewritten as

$$\frac{1}{1 - \underline{C}} = \frac{1}{1 - Q_1} \frac{1}{1 - Q_2} \frac{1}{1 - Q_3} \cdots, \quad (12)$$

and in this section, we prove that this factorization of series is the weight right factorization of C^* .

In a more general setting, T. Scharf and one of the authors [17], gave a recursive algorithm for computing the Q'_n 's. Recall that the algebra **Sym** of noncommutative symmetric functions is the free associative algebra $\mathbb{Q}\langle S_1, S_2, \dots \rangle$ generated by an infinite sequence of non commuting variables S_n , graded by the weight function $w(S_n) = n$. If $I = (i_1, \dots, i_n)$, one defines

$$\tilde{S}^I = (-1)^n S_{i_1} \cdots S_{i_n} \quad (13)$$

The Q_i can be computed following the rules given in [17]:

1. $F_1 = -\sum_i \tilde{S}_i$,
2. $F_{n+1} = F_n + Q_n(1 - F_n)$,

3. Q_n is the term of weight n in F_n multiplied by -1 .

Setting $Z_n = 1 - (1 - F_n)^{-1}$, we obtain

$$F_{n+1} = 1 - (1 - Z_{n+1})^{-1} \quad (14)$$

and

$$\begin{aligned} F_n + Q_n(1 - F_n) &= 1 - (1 - Z_n)^{-1} + Q_n(1 - Z_n)^{-1} \\ &= 1 - (1 - Q_n)(1 - Z_n)^{-1}. \end{aligned} \quad (15)$$

This implies

$$\begin{aligned} Z_{n+1} &= 1 - (1 - Z_n)(1 - Q_n)^{-1} \\ &= (Z_n - Q_n)(1 - Q_n)^{-1}. \end{aligned} \quad (16)$$

Following [17], each Q_i is multiplicity free on the \tilde{S}_I .

Let $Z_i[C]$ and $F_i[C]$ the values of Z_i and F_i under the specialization $S_n = S_n[C]$. The definition of noncommutative complete functions gives

$$\sigma_1[C] = \sum S_n[C] = \frac{1}{1 - \underline{C}}. \quad (17)$$

Hence, one has

$$Z_1[C] = 1 - (\sigma_1[C])^{-1} = \underline{C}, \quad (18)$$

$$Q_1[C] = S_1[C] = \underline{C}_1, \text{ and} \quad (19)$$

$$Z_1[C] - Q_1[C] = \underline{C} - \underline{C}_1. \quad (20)$$

It follows that $Z_1[C]$ and $Q_1[C]$ are the characteristic series of the codes $Z_1 = C$ and $Q_1 = C_1$. By induction, for each $n > 0$ the series $Z_{n+1}[C]$ and $Q_{n+1}[C]$ are the characteristic series of $Z_{n+1} = (Z_n \setminus Q_n)Q_n^*$ and $Q_n^* = Z_n \cap A^{\leq n}$. Hence, the following statement holds.

Proposition 1 *Let C be a code. Each C -Witt symmetric function $Q_i[C]$ is the characteristic series of a code, and the sequence obtained by deleting the empty set from $(Q_1, Q_2, \dots, Q_n, \dots)$ is the right length factorization of C^* .*

Example 4 The sequence $(a, Q_1[ba^*], Q_2[ba^*], \dots)$ is a factorization of A^* . The same method is applicable to compute homogeneous factorizations of

non-homogeneous alphabets. For example, considering the alphabet $A = \mathbb{N} \setminus 0$ with the weight $\omega = id$, one finds

$$\begin{aligned} Q_1[A] &= 1 \\ Q_2[A] &= 2 \\ Q_3[A] &= 21 + 3 \\ Q_4[A] &= 211 + 31 + 4 \\ Q_5[A] &= 2111 + 212 + 311 + 32 + 41 + 5 \\ Q_6[A] &= 21111 + 51 + 2112 + 6 + 3111 + 312 + 42 + 411 \end{aligned}$$

It is easy to see that this can be obtained from example 2 by the morphism $ba^n \rightarrow n + 1$.

One has the following decomposition

Corollary 2 *Let $w \in C^*$, then either $w \in C$ either there exist $w_1, w_2 \in C^*$ such that $w = w_1 w_2$, with $\omega(w_2) < \omega(w_1) < \omega(w)$. Furthermore, if $w_1 \notin C$ then $w_1 = w'_1 w''_1$ with $w'_1, w''_1 \in C^*$ and $\omega(w''_1) \leq \omega(w_2)$*

This follows from the standard bracketing process of regular factorizations, which is described in [18].

4.3 Noncommutative elementary symmetric functions and Lazard elimination

The link between Lazard elimination and noncommutative Witt symmetric functions can be better understood in terms of elementary symmetric functions. The generating function of noncommutative elementary symmetric function is

$$\lambda_t = \sum_{k \geq 0} \Lambda_k t^k. \quad (21)$$

These are related to noncommutative complete functions by

$$\sigma_t = \frac{1}{\lambda_{-t}} = \frac{1}{1 - \Lambda_1 t + \Lambda_2 t^2 - \dots + (-1)^n \Lambda_n t^n + \dots}. \quad (22)$$

If we set $\tilde{\Lambda}_n = (-1)^{n+1} \Lambda_n$, the series σ_1 can be considered as the characteristic series of the free monoid $\mathbf{\Lambda}^* = \{\tilde{\Lambda}_1, \tilde{\Lambda}_2, \dots, \tilde{\Lambda}_n \dots\}^*$. We endow this monoid with the weight function defined by $\omega(\tilde{\Lambda}_n) = n$. Then,

Theorem 3 *One has*

$$\mathbb{F}(\mathbf{\Lambda}, \omega) = (Q_1[\mathbf{\Lambda}], Q_2[\mathbf{\Lambda}], \dots) = (Q_1, Q_2, \dots) \quad (23)$$

This provides a simple algorithm for computing the decomposition of the noncommutative symmetric Witt functions on the basis of elementary symmetric functions. Let us give the computation of the first Q_i 's,

$$\begin{aligned} Q_1 &= \Lambda_1 \\ Q_2 &= -\Lambda_2 \\ Q_3 &= -\Lambda_2\Lambda_1 + \Lambda_3 \\ Q_4 &= -\Lambda_2\Lambda_1\Lambda_1 + \Lambda_3\Lambda_1 - \Lambda_4 \\ Q_5 &= -\Lambda_2\Lambda_1^3 + \Lambda_2\Lambda_1\Lambda_2 + \Lambda_3\Lambda_1\Lambda_1 - \Lambda_3\Lambda_2 - \Lambda_4\Lambda_1 + \Lambda_5 \\ Q_6 &= -\Lambda_2\Lambda_1^4 + \Lambda_5\Lambda_1 + \Lambda_2\Lambda_1\Lambda_1\Lambda_2 - \Lambda_6 + \Lambda_3\Lambda_1^3 - \Lambda_3\Lambda_1\Lambda_2 + \Lambda_4\Lambda_2 \\ &\quad - \Lambda_4\Lambda_1\Lambda_1 \\ Q_7 &= -\Lambda_3\Lambda_1^2\Lambda_2 + \Lambda_3\Lambda_1^4 - \Lambda_3\Lambda_1\Lambda_2\Lambda_1 + \Lambda_3\Lambda_1\Lambda_3 + \Lambda_3\Lambda_2^2 - \Lambda_4\Lambda_1^3 \\ &\quad + \Lambda_4\Lambda_1\Lambda_2 - \Lambda_4\Lambda_3 + \Lambda_4\Lambda_2\Lambda_1 + \Lambda_5\Lambda_1^2 - \Lambda_5\Lambda_2 - \Lambda_6\Lambda_1 + \Lambda_7 \\ &\quad - \Lambda_2\Lambda_1^5 + \Lambda_2\Lambda_1^3\Lambda_2 + \Lambda_2\Lambda_1^2\Lambda_2\Lambda_1 - \Lambda_2\Lambda_1^2\Lambda_3 - \Lambda_2\Lambda_1\Lambda_2^2 \end{aligned}$$

(compare Examples 2 and 4).

The decomposition of the elementary functions on the basis $Q_I = Q_{i_1} \cdots Q_{i_n}$ is obtained by inspection of the series

$$\frac{1}{\sigma_t} = \lambda_{-t} = 1 - \sum_n \tilde{\Lambda}_n t^n = \prod_i^{\leftarrow} (1 - Q_i t^i). \quad (24)$$

One finds

$$\tilde{\Lambda}_n = \sum_k (-1)^k \sum_{\substack{i_1 > \dots > i_k \\ i_1 + \dots + i_k = n}} Q_{i_1} \cdots Q_{i_k} \quad (25)$$

5 Examples involving Gaudier's *-multinomials

5.1 Gaudier's * multinomials

Classical (commutative) Witt vectors give rise to examples involving interesting integer sequences. Let us recall for example the construction of *-multinomial coefficients and *-factorials given by Gaudier in [6]. If R is a

\mathbb{Q} -algebra, there is a commutative diagram

$$\begin{array}{ccc} W(R) & \xrightarrow{e} & \Lambda(R) = 1 + tR[[t]] \\ w \downarrow & & \downarrow \partial \\ R^{\mathbb{N}^*} & \xrightarrow{\iota} & R[[t]] \end{array}$$

where

$$\begin{aligned} \partial &= \frac{d}{dt} \ln, \\ l(c_1, \dots, c_n, \dots) &= \sum_{n \geq 1} c_n t^{n-1}, \\ w &= (w_1, \dots, w_n, \dots), \\ e(a_1, \dots, a_n, \dots) &= \prod_{\vec{n}} \frac{1}{1 - a_n t^n}. \end{aligned}$$

These maps are all isomorphisms. Let i_1, i_2, \dots, i_k be k positive integers and $n = i_1 + \dots + i_k$. In [6] Gaudier has defined the $*$ -multinomial coefficient $*\binom{n}{i_1, \dots, i_k} \in W(\mathbb{Q})$ as the Witt vector such that $w_p = \binom{np}{pi_1, \dots, pi_k}$. In the same paper, he has defined the $*$ -factorial $*n!/n!$ by $w_p(*n!/n!) = \frac{1}{n!} \frac{(pn)!}{p^n}$. In particular, he has computed $e(*2!/2!) = e(\frac{1}{2} * \binom{2}{1})$ in closed form and related it to the Catalan numbers

$$e(*2!/2!) = \left(\frac{1 + \sqrt{1 - 4t}}{2} \right)^{-1} \quad (26)$$

This raised the question whether it was possible to find similar expressions for other $*$ -factorials and $*$ -multinomials, and to give combinatorial interpretations.

5.2 Non-commutative analogues of $\frac{1}{n} * \binom{n}{1}$

Consider the Dyck code of example 3. The free monoid D^* is a submonoid of $\{aa, ab, bb\}^*$, graded by $\rho(aa) = \rho(bb) = \rho(ab) = 1$. Under the D -specialization,

$$\Lambda_n = (-1)^{n+1} \sum_{\substack{w \in D \\ \rho(w)=n}} w. \quad (27)$$

Recall that the noncommutative power sums of the second kind Φ_n are defined by

$$\sum_{n \geq 1} \Phi_n \frac{t^n}{n} = \log \sigma_t. \quad (28)$$

If one interprets Dyck words as binary trees, the Φ_n can be regarded as sums over forests

$$\Phi_n = n \sum_k \frac{1}{k} \sum_{\substack{w_1, \dots, w_k \in D \\ \rho(w_1) + \dots + \rho(w_k) = n}} w_1 \cdots w_k. \quad (29)$$

Example 5.1 The first values of Φ_n are

$$\Phi_1 = ab$$

$$\Phi_2 = 2aabb + ab.ab$$

$$\Phi_3 = 3aaabbb + 3aababb + \frac{3}{2}aabb.ab + \frac{3}{2}ab.aabb + ab.ab.ab$$

$$\begin{aligned} \Phi_4 = & 4aaaabbbb + 4aaababb + 4aabaabbb + 4aabababb + 4aaababbb \\ & + 2aaaabbb.ab + 2aababb.ab + 2ab.aaabbb + 2ab.aababb + \frac{4}{3}aabb.ab.ab \\ & + \frac{4}{3}ab.aabb.ab + \frac{4}{3}ab.ab.aabb + ab.ab.ab.ab \end{aligned}$$

See example 3 for the first values of Q_n .

Setting $\pi(w) = 1$ for all words w , one has

$$\pi(\Phi_n) = w_n \left(\frac{1}{2!} * 2! \right) = w_n \left(\frac{1}{2} * \binom{2}{1} \right) \quad (30)$$

and one recovers (26).

More generally, for $*\frac{1}{n} \binom{n}{1}$, an easy application of the Lagrange inversion formula gives that

$$e \left(\frac{1}{n} * \binom{n}{1} \right) = \sum_{k \geq 0} \frac{\binom{nk}{k}}{(n-1)k+1} t^k \quad (31)$$

is also the generating series of n -ary trees. Consider the free monoid $F_n = T_n^*$ over the alphabet $T_n = \{a_{\mathbf{t}}\}$ whose letters are labelled by n -ary trees and weighted by $\rho(a_{\mathbf{t}}) = E(\mathbf{t})/n$, where $E(\mathbf{t})$ is the number of edges of \mathbf{t} . Instead of specializing Λ_k as in the case $n = 2$, we set $S_k = \sum_{\rho(\mathbf{t})=k} a_{\mathbf{t}}$. Then, we obtain

$$\Phi_k = k \sum_p \frac{(-1)^{p+1}}{p} \sum_{\substack{\mathbf{t}_1, \dots, \mathbf{t}_p \\ \rho(\mathbf{t}_1) + \dots + \rho(\mathbf{t}_p) = k}} a_{\mathbf{t}_1} \cdots a_{\mathbf{t}_p}. \quad (32)$$

Again, applying the morphism $\pi(w) = 1$, we obtain

$$\pi(\Phi_k) = w_k \left(\frac{1}{n} * \binom{n}{1} \right), \quad (33)$$

that is, we recover (31). Remark that this implies the identity

$$\sum_p \frac{(-1)^{p+1}}{p} \sum_{i_1+\dots+i_p=k} \prod_m \frac{\binom{ni_m}{i_m}}{(n-1)i_m+1} = \frac{1}{nk} \binom{nk}{k}. \quad (34)$$

5.3 Combinatorial interpretations of some $*$ -binomial coefficients

Remark that, as in section 5.2, the specialization $S_k = \sum_{\rho(\mathbf{t})=k+1} a_{\mathbf{t}}$ gives a non-commutative analogue of

$$e\left(* \binom{n}{1}\right)(z) = \sum_{k=1}^{\infty} \frac{\binom{nk}{k}}{(n-1)k+1} t^{k-1} = \frac{e\left(\frac{1}{n} * \binom{n}{1}\right) - 1}{z}. \quad (35)$$

This last equality can be generalized as follows.

Proposition 5.2 *Let $p \geq 1$ and $\omega = e^{2i\pi/p}$.*

$$e\left(* \binom{np}{p}\right) = \frac{-1}{z} \prod_{k=0}^{p-1} \left(1 - e\left(\frac{1}{n} * \binom{n}{1}\right) \right) (\omega^k z^{\frac{1}{p}}) \quad (36)$$

$$= \prod_{k=0}^{p-1} e\left(* \binom{n}{1}\right) (\omega^k z^{\frac{1}{p}}) \quad (37)$$

Proof We first prove (37), by computing

$$\begin{aligned} \ln \left(\prod_{k=0}^{p-1} e\left(* \binom{n}{1}\right) (\omega^k z^{\frac{1}{p}}) \right) &= \\ \sum_{k=0}^{p-1} \ln \left(e\left(* \binom{n}{1}\right) (\omega^k z^{\frac{1}{p}}) \right) &= \\ \sum_{k=0}^{p-1} \sum_{j \geq 1} \frac{\binom{jn}{j}}{j} \omega^{jk} z^{\frac{j}{p}} &= \sum_{k \geq 1} \frac{\binom{kn p}{kp}}{k} z^k \\ &= \ln \left(e\left(* \binom{np}{p}\right) \right). \end{aligned} \quad (38)$$

Next (36) follows from the equality

$$e\left(*\binom{np}{p}\right) = \prod_{k=0}^{p-1} \frac{e\left(\frac{1}{n} * \binom{n}{1}\right) (\omega^k z^{\frac{1}{p}}) - 1}{\omega^k z^{\frac{1}{p}}} \quad (39)$$

□

From [13] (Theorem 1, p.7), the series $e\left(*\binom{np}{p}\right)$ is the generating series of the number q_m of lattice paths from $(0, 0)$ to $((n-1)p, pm)$ that never go above the path $(\uparrow^{(n-1)p} \rightarrow^p)^m$ (lattice paths are represented by words over the alphabet $F = \{\rightarrow, \uparrow\}$, where \rightarrow means the elementary horizontal path $(1, 0)$ and \uparrow the elementary vertical path $(0, 1)$).

Example 5.3 The series

$$\begin{aligned} e\left(*\binom{4}{2}\right)(z) &= 1 + 6z + 53z^2 + 554z^3 + 6362z^4 + \dots \\ &= e\left(*\binom{2}{1}\right)(z^{\frac{1}{2}})e\left(*\binom{2}{1}\right)(-z^{\frac{1}{2}}) \end{aligned}$$

is also the generating series of ordered trees on $2n$ nodes with every subtree at the root having an even number of edges (see Sloane [16] ID number : A066357). See also [1] for another enumeration.

$$\begin{aligned} e\left(*\binom{6}{2}\right)(z) &= 1 + 15z + 360z^2 + 10463z^3 + \dots \\ &= e\left(*\binom{3}{1}\right)(z^{\frac{1}{2}})e\left(*\binom{3}{1}\right)(-z^{\frac{1}{2}}) \\ e\left(*\binom{6}{3}\right)(z) &= 1 + 20z + 662z^2 + 26780z^3 + \dots \\ &= e\left(*\binom{2}{1}\right)(z^{\frac{1}{3}})e\left(*\binom{2}{1}\right)(\exp\{\frac{2i\pi}{3}\}z^{\frac{1}{3}}) \times \\ &\quad \times e\left(*\binom{2}{1}\right)(\exp\{\frac{4i\pi}{3}\}z^{\frac{1}{3}}) \\ e\left(*\binom{12}{3}\right)(z) &= 1 + 220z + 91498z^2 + 47961320z^3 + \dots \\ &= e\left(*\binom{4}{1}\right)(z^{\frac{1}{3}})e\left(*\binom{4}{1}\right)(\exp\{\frac{2i\pi}{3}\}z^{\frac{1}{3}}) \times \\ &\quad \times e\left(*\binom{4}{1}\right)(\exp\{\frac{4i\pi}{3}\}z^{\frac{1}{3}}) \end{aligned}$$

One can construct noncommutative analogues of the $*\binom{np}{p}$ by specializing S_m to the sum of the words coding the lattice paths from $(0, 0)$ to $((n-1)p, mp)$ that never go above the path $(\uparrow^{(n-1)p} \rightarrow^p)^m$.

Example 5.4 Let us consider the non-commutative analogue of $*\binom{2}{1}$. Under this specialization, the first values of the S_n are

$$\begin{aligned} S_1 &= \uparrow \rightarrow + \rightarrow \uparrow \\ S_2 &= \rightarrow^2 \uparrow^2 + (\rightarrow \uparrow)^2 + \rightarrow \uparrow^2 \rightarrow + \uparrow \rightarrow^2 \uparrow + (\uparrow \rightarrow)^2 \end{aligned}$$

$$\begin{aligned}
S_3 = & \rightarrow^3\uparrow^3 + \rightarrow^2\uparrow\rightarrow\uparrow^2 + \rightarrow^2\uparrow^2\rightarrow\uparrow + \rightarrow^2\uparrow^3\rightarrow \\
& + \rightarrow\uparrow\rightarrow^2\uparrow^2 + (\rightarrow\uparrow)^3 + (\rightarrow\uparrow)^2\uparrow\rightarrow + \rightarrow\uparrow^2\rightarrow^2\uparrow \\
& + \rightarrow\uparrow^2\rightarrow\uparrow\rightarrow + \uparrow\rightarrow^3\uparrow^2 + \uparrow\rightarrow^2\uparrow\rightarrow\uparrow + \uparrow\rightarrow^2\uparrow^2\rightarrow + \uparrow\rightarrow\uparrow\rightarrow^2\uparrow + (\uparrow\rightarrow)^3
\end{aligned}$$

and the first values of the Λ_n are

$$\begin{aligned}
\Lambda_1 &= \uparrow\rightarrow + \rightarrow\uparrow \\
\Lambda_2 &= -\rightarrow^2\uparrow^2 \\
\Lambda_3 &= \rightarrow^2\uparrow\rightarrow\uparrow^2 + \rightarrow^3\uparrow^3
\end{aligned}$$

More generally, from

$$\Lambda_m = (-1)^{m+1} \sum_{k=1}^m (-1)^k \sum_{i_1+\dots+i_k=m} S_{i_1} \cdots S_{i_k}$$

one obtains

$$\Lambda_m = (-1)^{m+1} \sum_w w \tag{40}$$

where the sum is over all lattice paths from $(0, 0)$ to $((n-1)mp, mp)$ that never go above $(\uparrow^{(n-1)p}\rightarrow^p)^m$ and which avoid the points $((n-1)kp, kp)$ for $0 < k < m$. Hence,

$$\Phi_m = m \sum_w \frac{w}{c(w) + 1} \tag{41}$$

where the sum is over the lattice paths from $(0, 0)$ to $((n-1)mp, mp)$ below $(\uparrow^{(n-1)p}\rightarrow^p)^m$, and $c(w)$ is the number of points $((n-1)kp, kp)$ belonging to the path w , with $0 < k < m$.

Example 5.5 Let us consider the noncommutative analogue of $\ast \binom{4}{2}$. The first values of the Λ_n are

$$\begin{aligned}
\Lambda_1 &= \rightarrow^2\uparrow^2 + (\rightarrow\uparrow)^2 + \rightarrow\uparrow^2\rightarrow + \uparrow\rightarrow^2\uparrow + \uparrow\rightarrow\uparrow\rightarrow + \uparrow^2\rightarrow^2 \\
\Lambda_2 &= -(\rightarrow^4\uparrow^4 + \rightarrow^3\uparrow\rightarrow\uparrow^3 + \rightarrow^3\uparrow^2\rightarrow\uparrow^2 + \rightarrow^3\uparrow^3\rightarrow\uparrow + \rightarrow^3\uparrow^4\rightarrow \\
& + \rightarrow^2\uparrow\rightarrow^2\uparrow^3 + \rightarrow^2\uparrow(\rightarrow\uparrow)^2\uparrow + \rightarrow^2\uparrow\rightarrow\uparrow^2\rightarrow\uparrow + \rightarrow^2\uparrow\rightarrow\uparrow^3\rightarrow \\
& + \rightarrow\uparrow\rightarrow^3\uparrow^3 + \rightarrow\uparrow\rightarrow^2\uparrow\rightarrow\uparrow^2 + \rightarrow\uparrow\rightarrow\uparrow^2\rightarrow\uparrow + \rightarrow\uparrow\rightarrow\uparrow^3\rightarrow \\
& + \uparrow\rightarrow^4\uparrow^3 + \uparrow\rightarrow^3\uparrow\rightarrow\uparrow^2 + \uparrow\rightarrow^2\uparrow^2\rightarrow\uparrow + \uparrow\rightarrow^2\uparrow^3\rightarrow)
\end{aligned}$$

A natural question is whether it is possible to find similar interpretations for other \ast -binomials.

5.4 Shuffle analogues of *-multinomials in non commutative Witt vectors

We shall now construct another noncommutative analogue of *-multinomials. Let A be an alphabet and w_1, \dots, w_k be k words of respective lengths m_1, \dots, m_k . Let $w = w_1 \cdot w_2 \cdots w_k$ and $m = m_1 + \cdots + m_k$. The shuffle $w_1 \sqcup \cdots \sqcup w_k$ contains $\binom{m}{m_1, \dots, m_k}$ terms, and we can denote it by $\binom{w}{w_1, \dots, w_k}$ to emphasize this point. Then, we introduce the noncommutative Witt vector analogue

$$*\binom{w}{w_1, \dots, w_k} = (Q_1, Q_2, \dots, Q_n, \dots) \quad (42)$$

given by the sequence of Witt symmetric functions under the specialization

$$\Phi_n = w_1^n \sqcup w_2^n \sqcup \cdots \sqcup w_k^n. \quad (43)$$

Example 5.6 If $w_i = a^{p_i}$, one has

$$\begin{aligned} * \binom{w}{w_1, \dots, w_k} &= (w_1(* \binom{p_1 + \cdots + p_k}{p_1, \dots, p_k}) a^{p_1 + \cdots + p_k}, \\ &\dots, w_n(* \binom{p_1 + \cdots + p_k}{p_1, \dots, p_k}) a^{n(p_1 + \cdots + p_k)}, \dots). \end{aligned} \quad (44)$$

Sending a to 1, one recovers *-multinomials.

Acknowledgements

The authors are grateful to Henri Gaudier for fruitful discussions and comments.

This project has been partially supported by EC's IHRP Programme, grant HPRN-CT-2001-00272, "Algebraic Combinatorics in Europe".

References

- [1] C. BANDERIER, D. MERLINI, Lattice paths with an infinite set of jumps, *Proc. Formal Power Series and Algebraic Combinatorics 2002*, The University of Melbourne, Australia.

- [2] F. BASSINO, M.-P. BÉAL, D. PERRIN, Length distributions and regular sequences, in Codes, Systems and Graphical Models, *IMA Volumes in Mathematics and its Applications*, Springer-Verlag, p. 415-437, 2001.
- [3] J. BERSTEL, D. PERRIN, *Theory of Codes*, Academic Press, New-York, 1985.
- [4] P. CARTIER, Groupes formels associés aux anneaux de Witt généralisés, *C.R. Acad. Sci. Paris* **265** (1967), 49-52.
- [5] J. DIEUDONNÉ, Schur functions and group representations, *Astérisque* **87-88** (1983), 7-19.
- [6] H. GAUDIER, Relèvement des coefficients binômiaux dans les vecteurs de Witt, *Séminaire Lotharingien de Combinatoire* **16** (1988), 93-108.
- [7] I.M. GELFAND, D. KROB, A. LASCoux, B. LECLERC, V.S. RETAKH, J.-Y. THIBON, Noncommutative symmetric functions, *Adv. Math.* **112** (1995), 218-348.
- [8] M. HAZEWINKEL, *Formal Groups and Applications*, Academic Press, New York, 1978.
- [9] S. LANG, *Algebra*. Addison-Wesley Publishing Co., Inc., Reading, Mass. 1965.
- [10] M. LAZARD, *Groupes, anneaux de Lie et problème de Burnside*, Istituto Matematico dell'Università di Roma, 1960.
- [11] I.G. MACDONALD, *Symmetric functions and Hall polynomials*, Oxford Math. Monographs, Oxford, 1979; 2nd ed. 1995.
- [12] N. METROPOLIS, G.-C. ROTA, Witt vectors and the algebra of necklaces, *Adv. Math.* **50** (1983), 95-125.
- [13] A. DE MIER, M. NOY, A solution to the tennis ball problem, *Proc. Formal Power Series and Algebraic Combinatorics 2004*, University of British Columbia (Vancouver B.C., Canada).
- [14] C. REUTENAUER, On symmetric functions related to Witt vectors and the free Lie algebra, *Adv. Math.* **110** (1995), 234-246.

- [15] C. REUTENAUER, *Free Lie Algebras*. Clarendon Press, Oxford, 1993.
- [16] N. J. A. SLOANE, The On-Line Encyclopedia of Integer Sequences, <http://www.research.att.com/~njas/sequences/>
- [17] T. SCHARF, J.-Y. THIBON, On Witt vectors and symmetric functions, *Algebra Colloq.* **3** (1996), 231–238.
- [18] G. VIENNOT, *Algèbres de Lie libres et monoïdes libres*, Thèse de Doctorat d'Etat, Université Paris VII, 1974.
- [19] E. WITT, Zyklische Körper und Algebren der Charakteristik p vom Grade p^m , *J. reine angew. Math.* **176** (1937), 126-140.
- [20] E. WITT, Treue Darstellung Liescher Ringe, *J. reine angew. Math.* **177** (1937), 152-160.