

# Asymptotics of the Stirling numbers of the first kind revisited: A saddle point approach

Guy Louchard

Université Libre de Bruxelles, Dép. d'Informatique, CP 212, B-1050 Bruxelles, Belgium. [louchard@ulb.ac.be](mailto:louchard@ulb.ac.be)

received April 11, 2009, revised Dec. 21, 2009, accepted Feb. 18, 2010.

---

Using the saddle point method, we obtain from the generating function of the Stirling numbers of the first kind  $\left[ \begin{smallmatrix} n \\ j \end{smallmatrix} \right]$  and Cauchy's integral formula, asymptotic results in central and non-central regions. In the central region, we revisit the celebrated Goncharov theorem with more precision. In the region  $j = n - n^\alpha$ ,  $\alpha > 1/2$ , we analyze the dependence of  $\left[ \begin{smallmatrix} n \\ j \end{smallmatrix} \right]$  on  $\alpha$ .

**Keywords:** Stirling numbers, saddle point method.

---

To Philippe

## 1 Introduction

Let  $\left[ \begin{smallmatrix} n \\ j \end{smallmatrix} \right]$  be the Stirling number of the first kind (unsigned version). Their generating function is given by

$$\phi_n(z) = \prod_0^{n-1} (z+i) = \frac{\Gamma(z+n)}{\Gamma(z)}, \quad \phi_n(1) = n!.$$

In the sequel all asymptotics are meant for  $n \rightarrow \infty$ .

An asymptotic expansion for  $j = \mathcal{O}(1)$  is given in Wilf [14], which has been extended to the range  $j = \mathcal{O}(\ln n)$  by Hwang [6]. The generalized Stirling numbers have been considered by Tsylova [13] and Chelluri et al. [2]. The  $q$ -Stirling numbers are studied in Kyriakoussis and Vamvakari [9].

In Sec.2, we revisit the asymptotic expansions in the central region and in Sec.3, we analyse the non-central region  $j = n - n^\alpha$ ,  $\alpha > 1/2$ . We use Cauchy's integral formula and the saddle point method.

## 2 Central region

Consider the random variable  $J_n$ , with probability distribution

$$\mathbb{P}[J_n = j] = Z_n(j),$$

$$Z_n(j) := \frac{\begin{bmatrix} n \\ j \end{bmatrix}}{n!}.$$

The mean and variance are given by

$$M := \mathbb{E}(J_n) = \sum_0^{n-1} \frac{1}{1+i} = H_n = \psi(n+1) + \gamma,$$

$$\sigma^2 := \mathbb{V}(J_n) = \sum_0^{n-1} \frac{i}{(1+i)^2} = \psi(1, n+1) + \psi(n+1) - \frac{\pi^2}{6} + \gamma,$$

where  $\psi(x)$  is the digamma function,  $\psi(k, x)$  is the  $k$ th polygamma function, and

$$M \sim \ln(n) + \gamma + \frac{1}{2n} + \mathcal{O}\left(\frac{1}{n^2}\right),$$

$$\sigma^2 \sim \ln(n) - \frac{\pi^2}{6} + \gamma + \frac{3}{2n} + \mathcal{O}\left(\frac{1}{n^2}\right).$$

It is convenient to set

$$A_n := \ln(n) - \frac{\pi^2}{6} + \gamma = \ln\left(ne^{\gamma - \pi^2/6}\right),$$

and to consider all our next asymptotics ( $n \rightarrow \infty$ ) as functions of  $A_n$ . Of course, all asymptotics can be reformulated in terms of  $\ln(n)$ .

We have

$$M \sim A_n + \frac{\pi^2}{6} + \mathcal{O}\left(\frac{1}{n}\right),$$

$$\sigma^2 \sim A_n + \mathcal{O}\left(\frac{1}{n}\right).$$

A celebrated central limit theorem of Goncharov says that

$$J_n \sim \mathcal{N}(M, \sigma),$$

where  $\mathcal{N}$  is the Gaussian distribution, with a rate of convergence  $\mathcal{O}(1/\sqrt{\ln(n)})$ . This can also be deduced from the Quasi-Power theorem of Hwang [7],[8].

In this Section, we want to obtain a more precise local limit theorem for  $J_n$  in terms of  $x := \frac{J_n - M}{\sigma}$  and  $A_n$ . Actually, we obtain the following theorem, where we use  $B_n := \sqrt{A_n}$  to simplify the expressions.

**Theorem 2.1**

$$Z_n(j) \sim \frac{1}{\sqrt{2\pi}B_n} e^{-x^2/2} \cdot \left[ 1 + \frac{x^3/6 - x/2}{B_n} + \frac{3x^2/8 - x^4/6 - 1/12 + x^6/72}{B_n^2} + \frac{-\pi^2 x^3/18 + 37x^5/240 - 355x^3/144 + x/8 - x^7/48 + x^9/1296 + \pi^2 x/6 - \zeta(3)x + \zeta(3)x^3/3}{B_n^3} + \dots \right].$$

**Proof:** By Cauchy's theorem,

$$\begin{aligned} Z_n(j) &= \frac{1}{2\pi i} \int_{\Omega} \frac{\phi_n(z)}{z^{j+1}n!} dz \\ &= \frac{1}{2\pi i} \int_{\Omega} e^{S(z)} dz, \end{aligned}$$

where  $\Omega$  is inside the analyticity domain of the integrand and encircles the origin and

$$S(z) = S_1(z) + S_2(z), \quad S_1(z) = \sum_{i=0}^{n-1} \ln(z+i) - \ln(n!), \quad S_2(z) = -(j+1)\ln(z).$$

We will use the Saddle point method (for a good introduction to this method, see Flajolet and Sedgewick [3], ch.VIII). Set

$$S^{(i)} := \frac{d^i S}{dz^i}.$$

These derivatives can be expressed in terms of  $\psi(k, z+n)$  and  $\psi(k, z)$ .

First we must find the solution of

$$S^{(1)}(\tilde{z}) = 0 \tag{1}$$

with smallest module.

Set  $\tilde{z} := z^* - \varepsilon$ , where  $z^* = \lim_{n \rightarrow \infty} \tilde{z}$ . Here, it is easy to check that  $z^* = 1$ . Set  $j = M + x\sigma$ ,  $x$  fixed and  $B_n := \sqrt{A_n}$ .

This leads, to first order (keeping only the  $\varepsilon$  term in (1)), to

$$\varepsilon := \frac{-x}{B_n} + \frac{x^2 - 1}{B_n^2} + \dots + \frac{1}{n} \left( \frac{3x}{4B_n^3} + \dots \right) + \mathcal{O}\left(\frac{1}{n^2 B_n^4}\right).$$

This shows that, asymptotically,  $\varepsilon$  is given by a series of powers of  $n^{-1}$ , where each coefficient is given by a series of powers of  $B_n^{-1}$ . To obtain more precision, we set again  $j = M + x\sigma$ , expand in powers of  $n^{-1}$ , and equate each coefficient to 0. . This leads to (here and in the following, we provide only a few terms but Maple knows more)

$$\varepsilon = \frac{-x}{B_n} - \frac{1}{B_n^2} + \frac{0}{B_n^3} + \dots + \frac{1}{n} \left( \frac{3x}{4B_n^3} + \frac{x^2 + 3/2}{B_n^4} + \dots \right) + \mathcal{O}\left(\frac{1}{n^2 B_n^4}\right).$$

We have, with  $\tilde{z} := z^* - \varepsilon = 1 - \varepsilon$ ,

$$Z_n(j) = \frac{1}{2\pi\mathbf{i}} \int_{\Omega} \exp \left[ S(\tilde{z}) + S^{(2)}(\tilde{z})(z - \tilde{z})^2/2! + \sum_{l=3}^{\infty} S^{(l)}(\tilde{z})(z - \tilde{z})^l/l! \right] dz.$$

Note that the linear term vanishes. Set  $z = \tilde{z} + \mathbf{i}\tau$ . This gives

$$Z_n(j) = \frac{1}{2\pi} \exp[S(\tilde{z})] \int_{-\infty}^{\infty} \exp \left[ S^{(2)}(\tilde{z})(\mathbf{i}\tau)^2/2! + \sum_{l=3}^{\infty} S^{(l)}(\tilde{z})(\mathbf{i}\tau)^l/l! \right] d\tau. \quad (2)$$

The justification of the integration procedure is given in the Appendix. Let us first analyze  $S(\tilde{z})$ . We obtain

$$\begin{aligned} S(\tilde{z}) &= -x^2/2 + \frac{x^3/6 - x}{B_n} + \frac{-x^4/12 + x^2/2 - 1/2}{B_n^2} \\ &+ \frac{-x^3/3 + x^5/20 + x/2 - \pi^2 x^3/18 + \zeta(3)x^3/3}{B_n^3} + \dots + \mathcal{O}\left(\frac{1}{nB_n^3}\right). \end{aligned}$$

Also,

$$\begin{aligned} S^{(2)}(\tilde{z}) &= B_n^2 - B_n x - 1 + x^2 + \dots, \\ S^{(3)}(\tilde{z}) &= -2B_n^2 + 4B_n x - \pi^2/3 + 2\zeta(3) - 6x^2 + 4 + \dots, \\ S^{(4)}(\tilde{z}) &= 6B_n^2 - 18B_n x + 36x^2 - 18 + \pi^2 - \pi^4/15 + \dots, \\ S^{(l)}(\tilde{z}) &= \mathcal{O}(B_n^2), \quad l \geq 5. \end{aligned}$$

We need these many terms in the following. Note that, with  $z = \tilde{z}e^{i\theta}$ , this leads to

$$S^{(2)}(\tilde{z}) \frac{(z - \tilde{z})^2}{2} \sim -\frac{1}{2} \ln(n)\theta^2. \quad (3)$$

We can now compute (2), for instance by using the classical trick of setting


$$S^{(2)}(\tilde{z})(i\tau)^2/2! + \sum_{l=3}^{\infty} S^{(l)}(\tilde{z})(i\tau)^l/l! = -u^2/2.$$

Computing  $\tau$  as a truncated series in  $u$ , this gives, by inversion,

$$\tau = \left[ u(1 + x/(2B_n) + \dots) + u^2(\mathbf{i}/(3B_n) + \dots) + u^3(-1/(36B_n^2) + \dots) \right] / B_n + \dots$$

Setting  $d\tau = \frac{d\tau}{du} du$ , expanding w.r.t.  $B_n$  and integrating on  $[u = -\infty.. \infty]$ , this gives

$$\frac{1}{\sqrt{2\pi}B_n} \left[ 1 + \frac{x}{2B_n} + \frac{5/12 - x^2/8}{B_n^2} + \frac{x(8\pi^2 - 10 - 93x^2 - 48\zeta(3))}{48B_n^3} + \dots \right].$$



**Fig. 1:** Comparison between  $Z_n(j)$  and  $\frac{1}{\sqrt{2\pi}\sigma} \exp\left[-\left(\frac{j-M}{\sigma}\right)^2/2\right]$ ,  $n = 3000$

Finally (2) leads to

$$Z_n(j) \sim \frac{1}{\sqrt{2\pi}B_n} e^{-x^2/2} \cdot \exp\left[\frac{x^3/6 - x}{B_n} + \frac{-x^4/12 + x^2/2 - 1/2}{B_n^2} + \frac{-x^3/3 + x^5/20 + x/2 - \pi^2 x^3/18 + \zeta(3)x^3/3}{B_n^3} + \dots\right] \cdot \left[1 + \frac{x}{2B_n} + \frac{5/12 - x^2/8}{B_n^2} + \frac{x(8\pi^2 - 10 - 93x^2 - 48\zeta(3))}{48B_n^3} + \dots\right],$$


or

$$Z_n(j) \sim R_1,$$

$$R_1 = \frac{1}{\sqrt{2\pi}B_n} e^{-x^2/2} \cdot \left[1 + \frac{x^3/6 - x/2}{B_n} + \frac{3x^2/8 - x^4/6 - 1/12 + x^6/72}{B_n^2} + \frac{-\pi^2 x^3/18 + 37x^5/240 - 355x^3/144 + x/8 - x^7/48 + x^9/1296 + \pi^2 x/6 - \zeta(3)x + \zeta(3)x^3/3}{B_n^3} + \dots\right].$$

□

For  $n = 3000$ , a comparison between  $Z_n(j)$  and  $\frac{1}{\sqrt{2\pi}\sigma} \exp\left[-\left(\frac{j-M}{\sigma}\right)^2/2\right]$  is given in Figure 1.



**Fig. 2:**  $Z_n(j) / \left[ \frac{1}{\sqrt{2\pi\sigma}} \exp \left[ - \left( \frac{j-M}{\sigma} \right)^2 / 2 \right] \right]$ , color=red,  $Z_n(j)/R_1$ , color=blue,  $n = 3000$

Of course, only few values of  $j$  are significant and also the quality of the Gaussian is low, all asymptotic expressions depend actually on powers of  $A_n^{-1}$ , but  $A_n$  is not large.

A comparison of  $Z_n(j) / \left[ \frac{1}{\sqrt{2\pi\sigma}} \exp \left[ - \left( \frac{j-M}{\sigma} \right)^2 / 2 \right] \right]$  with  $Z_n(j)/R_1$ , with 2 terms in  $R_1$ , is given in Figure 2.

The precision of  $R_1$  is of order  $10^{-2}$ . Using 3 terms in  $R_1$  leads to a less good result:  $A_n$  is not large enough to take advantage of the  $A_n^{-3/2}$  term:  $A_n = 6.94$  here, we deal with asymptotic series, not necessarily convergent ones. More terms can be computed in  $R_1$  (which is almost automatic with Maple).

### 3 Large deviation, $j = n - n^\alpha$ , $1 > \alpha > 1/2$

The case  $j = \mathcal{O}(n)$  is analyzed in Timashev [12]. But he obtains a series of powers of  $n^{-1}$ , determined by a power series of a certain function that depends on the solution of a given non-linear differential equation of the first order. The coefficients obey some linear recurrence relations in the complex plane. The case  $j = n - c$ ,  $c$  constant, is considered in Grünberg [5]. As previous work for the case  $j = n - n^\alpha$ , let us mention Bender [1], Temme [11], Moser and Wyman [10] (see also the comments by Odlyzko in [4], p.1182). They all use, explicitly or not, the Saddle point method. For  $\alpha < 1/2$ , Moser and Wyman (6.9) give an explicit asymptotic expression. For  $\alpha > 1/2$ , they first compute in (4.52) the numerical solution  $zn$  of  $S'(zn) = 0$  and give in (4.51) an asymptotic expression. This is rather precise: for  $n = 50$ , this gives a precision of order  $10^{-4}$ . [1] and [11] also compute numerically  $zn$ .

However, all these results do not shed light on the dependence of  $[z^j]\phi(z)$  on  $n^\alpha$ . This is what we want

to explicit in this Section. It appears that the range  $\alpha > 1/2$  is more delicate than the other range.

Recall that

$$\phi_n(z) = \prod_0^{n-1} (z+i) = \frac{\Gamma(z+n)}{\Gamma(z)}.$$

We have

$$G_n(z) := \frac{\Gamma(z+n)}{\Gamma(z)z^{j+1}} = \exp[S(z)],$$

with

$$S(z) = S_1(z) + S_2(z), S_1(z) = \sum_0^{n-1} \ln(z+i), S_2(z) = -(j+1) \ln(z).$$

We first compute  $\tilde{z}$  such that

$$S'(\tilde{z}) = 0. \quad (4)$$

We have

$$S'(z) = \psi(z+n) - \psi(z) - \frac{j+1}{z}.$$

Similarly (we need these expressions later on)

$$S^{(2)}(z) = \psi(1, z+n) - \psi(1, z) + \frac{j+1}{z^2},$$

$$S^{(k)}(z) = \psi(k-1, z+n) - \psi(k-1, z) + (-1)^k (k-1)! \frac{j+1}{z^k}.$$

Some experiments with some values for  $\alpha$  ( $\alpha = 5/8$  is a good choice) show that  $\tilde{z}$  must be a combination of  $x = n^\alpha$  and  $y = n^{1-\alpha}$  and  $x \gg y \gg 1$ . Note that both  $x$  and  $y$  are large. We will derive series of powers of  $x^{-1}$ , where each coefficient is a series of powers of  $y^{-1}$ . We obtain the following theorem

**Theorem 3.1**

$$[z^j]\phi_n(z) \sim \frac{1}{\sqrt{2\pi}} \frac{y^2 \sqrt{x}}{2} \exp \left[ x \left[ 1 - \ln(2) + 2 \ln(y) + \ln(x) - \frac{2}{3y} - \frac{2}{9y^2} - \frac{44}{405y^3} - \frac{26}{405y^4} + \frac{40}{27y^5} \right. \right. \\ \left. \left. + \frac{179968}{18225y^6} + \frac{4727552}{127575y^7} + \frac{3436796}{32805y^8} + \frac{5492621728}{22143375y^9} + \dots \right] + \ln(2) - 2 \ln(y) - \ln(x) \right] \cdot \\ \cdot \left[ 1 - \frac{3}{3y} - \frac{1}{18y^2} - \frac{1}{30y^3} + \frac{17207}{3240y^4} + \dots + \frac{1}{x} \left( -\frac{1}{12} + \frac{1}{36y} - \frac{35}{216y^2} + \frac{15029}{3240y^3} + \dots \right) \right. \\ \left. + \frac{1}{x^2} \left( \frac{1}{288} - \frac{1}{864y} + \frac{3527}{5184y^2} + \dots \right) + \mathcal{O} \left( \frac{1}{x^3} \right) \right].$$

**Proof:** Let us summarize the different steps of the proof. First we compute  $\tilde{z}$  and  $S(\tilde{z})$  as  $S(\tilde{z}) = T_1 T_2$ , where  $T_1$  is the dominant term and  $T_2$  is a series of powers of  $x^{-1}$ , where each coefficient is a series of powers of  $y^{-1}$ . We expand  $T_3 := e^{T_2}$ . Next the integration procedure leads to  $\frac{y^2 \sqrt{x}}{2} T_4$ , where  $T_4$  is again a series of powers of  $x^{-1}$ , where each coefficient is a series of powers of  $y^{-1}$ . We set  $T_5 := \frac{1}{\sqrt{2\pi}} \frac{y^2 \sqrt{x}}{2} e^{T_1}$ . Finally, we obtain

$$[z^j]\phi_n(z) \sim T_5 T_3 T_4. \quad (5)$$

The first terms in the asymptotics of  $\tilde{z}$  are easy to compute: set  $\tilde{z} = n\beta$ . Equation (4) leads to

$$\psi(n(1 + \beta)) - \psi(n\beta) = \frac{1}{\beta} - \frac{1}{y\beta} + \frac{1}{n\beta}.$$

But  $\psi(n) \sim \ln(n)$ . So we have

$$\ln\left(1 + \frac{1}{\beta}\right) \sim \frac{1}{\beta} - \frac{1}{y\beta} + \frac{1}{n\beta},$$

or

$$\frac{1}{\beta} - \frac{1}{2\beta^2} \sim \frac{1}{\beta} - \frac{1}{y\beta},$$

or  $\beta \sim \frac{y}{2}$ .

More generally, we have

$$\beta = \frac{y}{2} \left[ 1 + \frac{a_1}{y} + \frac{a_2}{y^2} + \frac{a_3}{y^3} + \dots + \frac{1}{x} \left( 1 + \frac{b_1}{y} + \frac{b_2}{y^2} + \dots \right) + \frac{1}{x^2} \left( 1 + \frac{c_1}{y} + \frac{c_2}{y^2} + \dots \right) + \mathcal{O}\left(\frac{1}{x^3}\right) \right].$$

Note that  $\frac{1}{y^3}$  can be of the same order than  $\frac{1}{x}$ , see below.

By bootstrapping, we obtain (we give the first terms)

$$\begin{aligned} \tilde{z} &= \frac{ny}{2} \left[ 1 - \frac{4}{3y} + \frac{2}{9y^2} + \frac{8}{135y^3} + \frac{8}{405y^4} + \frac{16}{1701y^5} + \frac{232}{45525y^6} + \frac{64}{18225y^7} + \dots \right. \\ &\quad + \frac{1}{x} \left[ 1 - \frac{1}{y} + \frac{4}{9y^2} - \frac{16}{135y^3} + \dots \right] \\ &\quad + \frac{1}{x^2} \left[ 1 - \frac{1}{y} + \frac{0}{y^2} + \dots \right] \\ &\quad + \frac{1}{x^3} [1 + \dots] \\ &\quad \left. + \mathcal{O}\left(\frac{1}{x^4}\right) \right]. \end{aligned} \tag{6}$$


Note that the choice of dominant terms in the bracket of (6) depends on  $\alpha$ . For instance, for  $\alpha = 3/4$ , the dominant terms (in decreasing order) are

$$1, \frac{1}{y}, \frac{1}{y^2}, \left\{ \frac{1}{x}, \frac{1}{y^3} \right\}, \left\{ \frac{1}{xy}, \frac{1}{y^4} \right\}, \left\{ \frac{1}{xy^2}, \frac{1}{y^5} \right\}, \left\{ \frac{1}{x^2}, \frac{1}{xy^3}, \frac{1}{y^6} \right\}, \dots$$


Now we must compute  $S(\tilde{z})$  and its asymptotics. First we compute  $\ln(\tilde{z} + i)$ , take the asymptotics wrt  $x$ , sum on  $i$ , and again take the asymptotics wrt  $x$  (recall that  $n = xy$ ). this leads to

$$\begin{aligned} S_1(\tilde{z}) &= x \left[ (-\ln(2) + 2\ln(y) + \ln(x))y - \frac{1}{3} + \frac{4}{405y^2} + \frac{2}{405y^3} + \dots \right] + y - \frac{2}{3} - \frac{2}{3y} - \frac{49}{135y^2} + \dots \\ &\quad + \frac{1}{x} \left( \frac{y}{2} + \frac{1}{6y} + \dots \right) + \frac{1}{x^2} \left( \frac{y}{3} + \dots \right) + \mathcal{O}\left(\frac{y}{x^3}\right). \end{aligned}$$





**Fig. 3:**  $zn/\bar{z}, n = 500$ , as function of  $j$ , full range



**Fig. 4:**  $zn/\bar{z}, n = 500$ , as function of  $j$ , restricted range

Here we provide only a few terms but Maple knows more. Next

$$\begin{aligned}
S_2(\tilde{z}) &= x \left[ (\ln(2) - 2\ln(y) - \ln(x))y + \frac{4}{3} - \ln(2) + 2\ln(y) + \ln(x) - \frac{2}{3y} - \frac{94}{405y^2} + \dots \right] \\
&\quad - y + \frac{2}{3} + \ln(2) - 2\ln(y) - \ln(x) + \frac{1}{y} + \frac{94}{135y^2} + \dots \\
&\quad + \frac{1}{x} \left( \frac{y}{2} + \frac{1}{6y} + \dots \right) \\
&\quad + \frac{1}{x^2} \left( \frac{y}{3} + \dots \right) \\
&\quad + \mathcal{O}\left(\frac{y}{x^3}\right).
\end{aligned}$$

So, finally

$$\begin{aligned}
S(\tilde{z}) &= S_1(\tilde{z}) + S_2(\tilde{z}) \sim x \left[ 1 - \ln(2) + 2\ln(y) + \ln(x) - \frac{2}{3y} + \dots \right] \\
&\quad + \ln(2) - 2\ln(y) - \ln(x) + \frac{1}{3y} + \frac{1}{3y^2} + \dots \\
&\quad + \frac{1}{x} \left( -\frac{1}{2} + \frac{1}{3y} - \frac{1}{2y^2} + \dots \right) \\
&\quad + \frac{1}{x^2} \left( -\frac{1}{6} + \frac{19}{18y^2} \dots \right) \\
&\quad + \mathcal{O}\left(\frac{1}{x^3}\right).
\end{aligned}$$

Now we split  $S(\tilde{z})$  into two parts:

$$\begin{aligned}
T_1 &= x \left[ 1 - \ln(2) + 2\ln(y) + \ln(x) - \frac{2}{3y} + \dots \right] + \ln(2) - 2\ln(y) - \ln(x), \\
T_2 &= \frac{1}{3y} + \frac{1}{3y^2} + \dots \\
&\quad + \frac{1}{x} \left( -\frac{1}{2} + \frac{1}{3y} - \frac{1}{2y^2} + \dots \right) \\
&\quad + \frac{1}{x^2} \left( -\frac{1}{6} - \frac{17}{18y^2} \dots \right) \\
&\quad + \mathcal{O}\left(\frac{1}{x^3}\right).
\end{aligned}$$

Note that the dominant term of  $T_1$  is given by

$$T_1 \sim (2 - \alpha)n^\alpha \ln(n). \quad (7)$$

We obtain

$$\exp(S(\tilde{z})) = e^{T_1} e^{T_2} = e^{T_1} T_3,$$

with

$$\begin{aligned} T_3 = e^{T_2} &= 1 + \frac{1}{3y} + \frac{7}{18y^2} + \frac{89}{270y^3} + \frac{18263}{3240y^4} + \frac{98009}{3240y^5} + \frac{9517337}{97200y^6} + \frac{491504273}{2041200y^7} + \dots \\ &+ \frac{1}{x} \left( -\frac{1}{2} + \frac{1}{6y} - \frac{7}{12y^2} + \frac{2311}{540y^3} + \frac{112469}{6480y^4} + \frac{5137}{144y^5} + \dots \right) \\ &+ \frac{1}{x^2} \left( -\frac{1}{24} - \frac{13}{72y} - \frac{557}{932y^2} + \dots \right) \\ &+ \mathcal{O}\left(\frac{1}{x^3}\right). \end{aligned}$$

Here we have given all terms compatible with the expansion (6). Also, with more precision,


$$\begin{aligned} T_1 = x &\left[ 1 - \ln(2) + 2 \ln(y) + \ln(x) - \frac{2}{3y} - \frac{2}{9y^2} - \frac{44}{405y^3} - \frac{26}{405y^4} + \frac{40}{27y^5} \right. \\ &\left. + \frac{179968}{18225y^6} + \frac{4727552}{127575y^7} + \frac{3436796}{32805y^8} + \frac{5492621728}{22143375y^9} + \dots \right] \\ &+ \ln(2) - 2 \ln(y) - \ln(x). \end{aligned}$$

Now we must consider  $S^{(k)}(\tilde{z})$ . By direct expansion, this gives the following expressions (again we provide only the first few terms). We must use up to six derivatives to get a sufficient precision (of order  $x^{-2}$ ) in the Saddle integrals.


$$\begin{aligned} S^{(2)}(\tilde{z}) &= \frac{1}{x} \left[ \frac{4}{y^4} + \frac{16}{3y^5} + \dots \right] \\ &+ \frac{1}{x^2} \left[ -\frac{12}{y^4} - \frac{40}{3y^5} + \dots \right] \\ &+ \frac{1}{x^3} \left[ \frac{12}{y^4} + \frac{8}{y^5} + \dots \right] \\ &+ \frac{1}{x^4} \left[ \frac{-4}{y^4} + \dots \right] \\ &+ \mathcal{O}\left(\frac{1}{x^5 y^4}\right), \end{aligned} \tag{8}$$

Note that, with  $z = \tilde{z}e^{i\theta}$ , this leads to

$$S^{(2)}(\tilde{z}) \frac{(z - \tilde{z})^2}{2} \sim -\frac{1}{2} n^\alpha \theta^2. \tag{9}$$



**Fig. 5:**  $G_n(zn)/G_n(\tilde{z})$ ,  $n = 500$ , as function of  $j$



**Fig. 6:** The quotient of the expression (8) and  $S^{(2)}(\tilde{z})$  as function of  $j$ ,  $n = 500$

$$\begin{aligned}
S^{(3)}(\tilde{z}) &= \frac{1}{x^2} \left[ -\frac{32}{y^6} + \dots \right] \\
&+ \frac{1}{x^3} \left[ \frac{128}{y^6} + \dots \right] \\
&+ \frac{1}{x^4} \left[ -\frac{192}{y^6} + \dots \right] \\
&+ \frac{1}{x^5} \left[ \frac{128}{y^6} + \dots \right] \\
&+ \mathcal{O}\left(\frac{1}{x^6 y^6}\right),
\end{aligned}$$


$$\begin{aligned}
S^{(4)}(\tilde{z}) &= \frac{1}{x^3} \left[ \frac{288}{y^8} + \dots \right] \\
&+ \frac{1}{x^4} \left[ -\frac{1440}{y^8} + \dots \right] \\
&+ \frac{1}{x^5} \left[ \frac{2880}{y^8} + \dots \right] \\
&+ \frac{1}{x^6} \left[ -\frac{2880}{y^8} + \dots \right] \\
&+ \mathcal{O}\left(\frac{1}{x^7 y^8}\right),
\end{aligned}$$

$$\begin{aligned}
S^{(5)}(\tilde{z}) &= \frac{1}{x^4} \left[ -\frac{3072}{y^{10}} + \dots \right] \\
&+ \frac{1}{x^5} \left[ \frac{18432}{y^{10}} + \dots \right] \\
&+ \mathcal{O}\left(\frac{1}{x^6 y^{10}}\right),
\end{aligned}$$

$$\begin{aligned}
S^{(6)}(\tilde{z}) &= \frac{1}{x^5} \left[ \frac{38400}{y^{12}} + \dots \right] \\
&+ \frac{1}{x^6} \left[ \frac{268800}{y^{12}} + \dots \right] \\
&+ \mathcal{O}\left(\frac{1}{x^7 y^{12}}\right).
\end{aligned}$$

We proceed now as in Section 2. Again, the justification of the integration procedure is given in the Appendix. This leads to

$$\tau = \frac{y^2 \sqrt{x}}{2} \left[ u a_1 + \frac{u^2 a_2}{x^{1/2}} + \frac{u^3 a_3}{x} + \frac{u^4 a_4}{x^{3/2}} + \frac{u^5 a_5}{x^2} + \mathcal{O}\left(\frac{u^6}{x^{5/2}}\right) \right].$$



**Fig. 7:** The quotient of the expression (8) and  $S^{(2)}(\tilde{z})$ , as function of  $j$ ,  $n = 500$ . Restricted range,  $\alpha \leq .84$

We give only  $a_1$ :

$$a_1 = 1 - \frac{2}{3y} - \frac{2}{9y^2} + \dots + \frac{1}{x} \left( \frac{3}{2} - \frac{4}{3y} + \dots \right) + \frac{1}{x^2} \left( \frac{15}{8} - \frac{7}{4y} + \dots \right) + \mathcal{O} \left( \frac{1}{x^3} \right).$$

This leads to

$$\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-u^2/2} \tau'(u) du = \frac{y^2 \sqrt{x}}{2} T_4,$$

with

$$T_4 = 1 - \frac{2}{3y} - \frac{2}{9y^2} + \dots + \frac{1}{x} \left( \frac{5}{12} - \frac{11}{18y} + \dots \right) + \frac{1}{x^2} \left( \frac{73}{288} - \frac{133}{432y} + \dots \right) + \frac{1}{x^3} \left( \frac{721}{576} + \dots \right) + \mathcal{O} \left( \frac{1}{x^4} \right).$$

Set

$$T_5 := \frac{1}{\sqrt{2\pi}} \frac{y^2 \sqrt{x}}{2} e^{T_1}.$$


This leads to

$$[z^j] \phi_n(z) \sim T_5 T_3 T_4. \quad (10)$$


We can of course combine  $T_3$  and  $T_4$ :

$$T_6 := T_3 T_4 = 1 - \frac{3}{3y} - \frac{1}{18y^2} - \frac{1}{30y^3} + \frac{17207}{3240y^4} + \dots + \frac{1}{x} \left( -\frac{1}{12} + \frac{1}{36y} - \frac{35}{216y^2} + \frac{15029}{3240y^3} + \dots \right) + \frac{1}{x^2} \left( \frac{1}{288} - \frac{1}{864y} + \frac{3527}{5184y^2} + \dots \right) + \mathcal{O} \left( \frac{1}{x^3} \right).$$

□



**Fig. 8:** The quotient  $[z^j]\phi_n(z)/T_8$ , two terms in  $T_4$ , as function of  $j$ ,  $n = 500$



**Fig. 9:** The quotient  $[z^j]\phi_n(z)/T_8$ , three terms in  $T_4$ , as function of  $j$ ,  $n = 500$

Let us consider the precision of our asymptotics.

The quality of asymptotic (6) is given in Figure 3 and 4, for  $n = 500$ , and  $x \in [\sqrt{n}, n^{0.9}]$  (first range) so that  $y \in [n^{0.1}, \sqrt{n}]$ . For some values of  $j = n - x$ , we show  $\tilde{z}/zn$ , where, as mentioned,  $zn$  is the numerical solution of  $S'(zn) = 0$ . In the full range  $j \in [n - n^{0.9}, n - \sqrt{n}]$ , the precision is of order  $10^{-5}$ , in a restricted range, the precision is of order  $10^{-6}$ .

Also a comparison of  $G_n(\tilde{z})$  and  $G_n(zn)$  is given in Figure 5, showing again a precision of order  $10^{-6}$ .

To check the quality of asymptotic (6), we give in Figure 6 the comparison between the expression (8) and  $S^{(2)}(\tilde{z})$ . The precision is of order  $10^{-2}$ .

In a restricted range, given in Figure 7, the precision is of order  $10^{-4}$ .  $\alpha \leq 0.84$  in this range.

We have made several experiments with (10), with  $n$  up to 500. The result is unsatisfactory, only values of  $x$  of order  $\sqrt{n}$  give reasonable results. Also using  $e^{T_2}$  instead of  $T_3$  does not improve the precision. Actually, only very large values of  $n$  lead to good precision. So we turn to another formulation: instead of using  $e^{T_1 T_3}$  for  $e^{S(\tilde{z})}$ , we plug directly  $\tilde{z}$  into  $G_n(z)$ , ie we set

$$T_7 = G_n(\tilde{z}),$$

leading to

$$[z^j]\phi_n(z) \sim \frac{1}{\sqrt{2\pi}} \frac{y^2 \sqrt{x}}{2} T_7 T_4 =: T_8 \text{ say.}$$

For  $n = 500$ , using two and three terms in  $T_4$ , we give in Figures 8 and 9, the quotient  $[z^j]\phi_n(z)/T_8$ . The precision is of order  $10^{-5}$ .

## 4 Conclusion

Using an almost mechanized program in Maple, we have obtained some asymptotic expressions for Stirling numbers in central and non-central regions. We intend to use these techniques in other non-central ranges.

### Acknowledgement

The pertinent comments of the referee led to improvements in the presentation.

## A Appendix. Justification of the integration procedure

### A.1 The central region

We proceed as in Flajolet and Sedgewick [3], ch.VIII. We can choose here  $\tilde{z} = 1$ . This leads, with  $z = e^{i\theta}$ , to

$$S(z) \sim S_0(z) + \mathcal{O}\left(\sqrt{\ln(n)\theta}\right) + \text{constant term,}$$



with

$$\begin{aligned} S_0(z) &= \sum_{k=0}^{n-1} \ln[e^{i\theta} + k] - H_n i\theta \\ &\sim \sum_{k=0}^{n-1} \frac{1}{1+k} [e^{i\theta} - 1] - \frac{1}{2} \sum_{k=0}^{n-1} \left[ \frac{1}{1+k} [e^{i\theta} - 1] \right]^2 - H_n i\theta + \mathcal{O}(\theta^3) \\ &\sim H_n [e^{i\theta} - 1 - i\theta] + \mathcal{O}(\theta^2). \end{aligned}$$

Set

$$h(\theta) := e^{i\theta} - 1 - i\theta.$$

We have

$$h(\theta) \sim -\frac{\theta^2}{2},$$

which conforms to (3).

The function  $h(\theta)$  is the same as in [3], Ex.VIII.3, which proves the validity of our integration procedure: we use here  $H_n \sim \ln(n)$  instead of  $n$ . The complete asymptotic expansion is justified as in [3], Ex.VIII.4.

## A.2 The non-central region

We choose here  $\tilde{z} = \frac{ny}{2} = \frac{n^{2-\alpha}}{2} := \delta$ , say. We have

$$\begin{aligned} \frac{1}{2} &< \alpha < 1, \\ n^\alpha &= \frac{n^2}{2\delta}, \\ n^2 &\gg \delta \gg n \gg n^\alpha. \end{aligned}$$

Set  $z = \delta e^{i\theta}$ , this leads, with Euler-Maclaurin formula, with the first correction (the other corrections are negligible), to

$$\begin{aligned} S(z) &\sim \sum_{k=0}^{n-1} \ln [\delta e^{i\theta} + k] - (n - n^\alpha) i\theta - (n - n^\alpha) \ln(\delta) \\ &\sim \ln [n + \delta e^{i\theta}] [n + \delta e^{i\theta}] - n - \delta e^{i\theta} \ln [\delta e^{i\theta}] - \left[ n - \frac{n^2}{2\delta} \right] (i\theta + \ln(\delta)) - \frac{1}{2} \ln [n + \delta e^{i\theta}] + \frac{1}{2} \ln [\delta e^{i\theta}]. \end{aligned}$$

Set now  $n = \rho\delta$ ,  $\rho = 2n^{\alpha-1} \ll 1$  and expand wrt  $\rho$ . This gives

$$\begin{aligned} S(z) &\sim \rho \left[ -\frac{1}{2} e^{-i\theta} \right] \\ &\quad + \rho^2 \left[ \delta \frac{1 + i\theta e^{i\theta}}{2e^{i\theta}} + \frac{1}{4} e^{-2i\theta} + \frac{1}{2} \delta \ln(\delta) \right] \\ &\quad + \rho^3 \left[ -\frac{\delta}{6} e^{-2i\theta} - \frac{1}{6} e^{-3i\theta} \right] \\ &\quad + \mathcal{O}(\delta\rho^4). \end{aligned}$$

Note that the dominant constant contribution is given by  $\frac{1}{2}\rho^2\delta\ln(\delta) = (2 - \alpha)n^\alpha\ln(n)$ , which conforms to (7). The first term gives a variable part  $\mathcal{O}(\rho)$ . The second term gives a variable part  $2n^\alpha h(\theta) + \mathcal{O}(\rho^2)$ , with

$$h(\theta) := \frac{1 + i\theta e^{i\theta}}{2e^{i\theta}}.$$

The third term give  $\mathcal{O}(n^{2\alpha-1}) \ll n^\alpha$ . Note that  $2n^\alpha h(\theta) \sim -\frac{1}{2}n^\alpha\theta^2$ , which conforms to (9). The function  $|e^{h(\theta)}| = e^{\cos(\theta)/2}$  is unimodal with peak at 0 and  $h(0) = 1/2$ . Let us introduce a splitting value  $\theta_0$  such that  $n^\alpha\theta_0^2 \rightarrow \infty, n^\alpha\theta_0^3 \rightarrow 0, n \rightarrow \infty$ . For instance, we choose  $\theta_0 = n^\beta, \beta = -\frac{5\alpha}{12}$ . By unimodality property of the cosine, the tail integral

$$K_n^{(1)} := \int_{\theta_0}^{2\pi-\theta_0} e^{2n^\alpha(h(\theta)-1/2)} d\theta$$

is such that

$$|K_n^{(1)}| = \mathcal{O}\left(e^{n^\alpha[\cos(\theta_0)-1]}\right) = \mathcal{O}\left(e^{-Cn^{\alpha/6}}\right)$$

for some  $C > 0$ . The tail integral is exponentially small.

As  $h(\theta) \sim -\frac{\theta^2}{4}$ , the central approximation and the tail completion are immediate.

## References

- [1] E.A. Bender. Central and local limit theorems applied to asymptotics enumeration. *Journal of Combinatorial Theory*, Series A, 15:91–111, 1973.
- [2] R. Chelluri, L. B. Richmond, and N. M. Temme. Asymptotic estimates for generalized Stirling numbers. Technical report, CWI, MAS-R9923, 1999.
- [3] P. Flajolet and R. Sedgewick. *Analytic combinatorics*. Cambridge University press, 2009.
- [4] R. L. Graham, M. Grötschel, and L. Lovász. *Handbook of Combinatorics*, vol. 2. Elsevier, 1995.
- [5] D. B. Grünberg. On asymptotics, Stirling numbers, Gamma function and polylogs. *Results in Mathematics*, 49(1):89–125, 2006.
- [6] H. K. Hwang. Asymptotic expansions for the Stirling numbers of the first kind. *Journal of Combinatorial Theory*, Series A, 71(2):343–351, 1995.
- [7] H.K. Hwang. Théorèmes limites pour les structures aléatoires et les fonctions arithmétiques. 1994. Thèse, Ecole Polytechnique de Palaiseau.
- [8] H.K. Hwang. On convergence rates in the central limit theorems for combinatorial structures. *European Journal of Combinatorics*, 19:329–343, 1998.
- [9] A. Kyriakoussis and M. G. Vamvakari. On asymptotic for the signless noncentral q-Stirling numbers of the first kind. *Studies in Applied Mathematics*, 117:191–213, 2006.
- [10] L. Moser and M. Wyman. Asymptotic development of the Stirling numbers of the first kind. *Journal of the London Mathematical Society*, 33:133–146, 1958.
- [11] N.M. Temme. Asymptotic estimates of Stirling numbers. *Studies in Applied Mathematics*, 89:233–243, 1993.
- [12] A. N. Timashev. On asymptotic expansions of Stirling numbers of the first and second kind. *Discrete Mathematics and Applications*, 8(5):533–544, 1998.
- [13] E. G. Tsylova. Probabilistic methods for obtaining asymptotic formulas for generalized Stirling numbers. *Discrete Mathematics and Applications*, 75(2):1607–1614, 1995.
- [14] H. S. Wilf. The asymptotic behavior of the Stirling numbers of the first kind. *Journal of Combinatorial Theory*, Series A, 64:344–349, 1993.